

Full wwPDB X-ray Structure Validation Report (i)

Oct 3, 2023 – 04:30 AM EDT

PDB ID : 6OHA

Title : Yeast Guanine Deaminase Authors : Shek, R.S.; French, J.B.

Deposited on : 2019-04-05

Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)


There are 6 unique types of molecules in this entry. The entry contains 7357 atoms, of which 3476 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Probable guanine deaminase.

Mo	l Chain	Residues			Atom	ıs			ZeroOcc	AltConf	Trace
1	A	451	Total 6999	C 2241	H 3463	N 602	O 674	S 19	0	7	0

• Molecule 2 is XANTHINE (three-letter code: XAN) (formula: $C_5H_4N_4O_2$).




ľ	Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
	າ	Λ	1	Total	С	Н	N	О	0	0
	2	А	1	12	5	1	4	2	0	U

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0


• Molecule 4 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C₁₀H₂₂O₆).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C H O 28 10 12 6	0	0
4	A	1	Total C O 14 10 4	0	0
4	A	1	Total C O 13 8 5	0	0

 \bullet Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Ato	ms		ZeroOcc	AltConf
5	A	1	Total 5	O 4	S 1	0	0

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	275	Total O 275 275	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants	106.81Å 106.81Å 114.18Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.07 - 2.21	Depositor
% Data completeness	99.1 (29.07-2.21)	Depositor
(in resolution range)	,	-
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.47 (at 2.22Å)	Xtriage
Refinement program	PHENIX 1.14_3260, PHENIX 1.14_3260	Depositor
R, R_{free}	0.156 , 0.203	Depositor
Wilson B-factor (\mathring{A}^2)	30.0	Xtriage
Anisotropy	0.442	Xtriage
L-test for twinning ²	$< L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7357	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type C		Res	Link	Bo	ond leng	ths	В	ond ang	les
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	A	507	-	4,4,4	0.22	0	6,6,6	0.11	0
5	SO4	A	508	-	4,4,4	0.14	0	6,6,6	0.15	0
4	1PE	A	503	-	15,15,15	0.54	0	14,14,14	0.26	0
4	1PE	A	504	-	13,13,15	0.61	0	12,12,14	0.66	0
2	XAN	A	501	-	8,12,12	2.56	3 (37%)	4,17,17	2.60	3 (75%)
5	SO4	A	506	-	4,4,4	0.20	0	6,6,6	0.25	0
4	1PE	A	505	-	12,12,15	0.56	0	11,11,14	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	1PE	A	504	-	-	5/11/11/13	-
2	XAN	A	501	-	-	-	0/2/2/2
4	1PE	A	503	-	-	8/13/13/13	-
4	1PE	A	505	-	-	6/10/10/13	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
2	A	501	XAN	O6-C6	5.86	1.39	1.24
2	A	501	XAN	C6-N1	2.61	1.37	1.33
2	A	501	XAN	C4-N9	2.15	1.38	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	A	501	XAN	C2-N1-C6	3.34	117.97	115.14
2	A	501	XAN	C4-C5-C6	-2.76	118.16	120.80
2	A	501	XAN	C4-C5-N7	-2.36	106.94	109.40

There are no chirality outliers.

All (19) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	A	503	1PE	OH5-C14-C24-OH4
4	A	504	1PE	ОН4-С13-С23-ОН3
4	A	503	1PE	ОН6-С15-С25-ОН5
4	A	504	1PE	OH5-C14-C24-OH4
4	A	503	1PE	ОН7-С16-С26-ОН6
4	A	505	1PE	OH6-C15-C25-OH5
4	A	503	1PE	ОН4-С13-С23-ОН3
4	A	505	1PE	ОН7-С16-С26-ОН6
4	A	504	1PE	OH6-C15-C25-OH5
4	A	505	1PE	ОН4-С13-С23-ОН3
4	A	505	1PE	C23-C13-OH4-C24
4	A	503	1PE	OH2-C12-C22-OH3
4	A	505	1PE	C16-C26-OH6-C15
4	A	503	1PE	C24-C14-OH5-C25
4	A	503	1PE	C16-C26-OH6-C15
4	A	505	1PE	C25-C15-OH6-C26
4	A	504	1PE	C25-C15-OH6-C26
4	A	503	1PE	C14-C24-OH4-C13
4	A	504	1PE	C15-C25-OH5-C14

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

