



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 23, 2020 – 06:11 AM BST

PDB ID : 6TB1
Title : Crystal structure of thermostable omega transaminase 6-fold mutant from *Pseudomonas jessenii*
Authors : Capra, N.; Rozeboom, H.J.; Thunnissen, A.M.W.H.; Janssen, D.B.
Deposited on : 2019-10-31
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	458	 6% 97%
1	B	458	 3% 98%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate aminotransferase family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	451	3467	2218	600	633	16	0	0	0
1	B	453	3517	2246	610	644	17	0	5	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	ALA	PRO	engineered mutation	UNP A0A2D8IND4
A	38	GLN	GLU	engineered mutation	UNP A0A2D8IND4
A	60	VAL	ALA	engineered mutation	UNP A0A2D8IND4
A	87	ASN	SER	engineered mutation	UNP A0A2D8IND4
A	128	PHE	MET	engineered mutation	UNP A0A2D8IND4
A	154	VAL	ILE	engineered mutation	UNP A0A2D8IND4
A	456	PRO	-	expression tag	UNP A0A2D8IND4
A	457	GLY	-	expression tag	UNP A0A2D8IND4
A	458	GLY	-	expression tag	UNP A0A2D8IND4
B	9	ALA	PRO	engineered mutation	UNP A0A2D8IND4
B	38	GLN	GLU	engineered mutation	UNP A0A2D8IND4
B	60	VAL	ALA	engineered mutation	UNP A0A2D8IND4
B	87	ASN	SER	engineered mutation	UNP A0A2D8IND4
B	128	PHE	MET	engineered mutation	UNP A0A2D8IND4
B	154	VAL	ILE	engineered mutation	UNP A0A2D8IND4
B	456	PRO	-	expression tag	UNP A0A2D8IND4
B	457	GLY	-	expression tag	UNP A0A2D8IND4
B	458	GLY	-	expression tag	UNP A0A2D8IND4

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	15	8	1	5	1	0	0
2	B	1	15	8	1	5	1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SUCCINIC ACID (three-letter code: SIN) (formula: C₄H₆O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			8	4	4		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Na	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	245	Total	O	0	5
			250	250		
6	B	279	Total	O	0	5
			284	284		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aspartate aminotransferase family protein



- Molecule 1: Aspartate aminotransferase family protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	98.92Å 98.92Å 120.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.74 – 1.85 45.74 – 1.85	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.74-1.85) 100.0 (45.74-1.85)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.156 , 0.187 0.170 , 0.195	Depositor DCC
R_{free} test set	5035 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7582	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SIN, NA, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/3549	0.75	0/4811
1	B	0.73	0/3599	0.74	0/4876
All	All	0.73	0/7148	0.75	0/9687

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3467	0	3440	4	0
1	B	3517	0	3486	3	0
2	A	15	0	6	0	0
2	B	15	0	6	0	0
3	A	24	0	32	1	0
4	B	8	0	4	0	0
5	B	2	0	0	0	0
6	A	250	0	0	1	0
6	B	284	0	0	1	0
All	All	7582	0	6974	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 0.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:385[B]:LEU:HD11	1:B:405:MET:HG3	1.94	0.50
1:A:230:ALA:HB1	3:A:1003:GOL:H11	1.95	0.48
1:B:385[B]:LEU:CD1	1:B:405:MET:HG3	2.46	0.46
1:A:385:LEU:HD11	1:A:405:MET:HG3	1.99	0.44
1:B:367[B]:ASP:OD2	6:B:1101:HOH:O	2.22	0.42
1:A:75:GLN:NE2	6:A:1109:HOH:O	2.53	0.41
1:A:435:GLU:HA	1:A:438:MET:CE	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	449/458 (98%)	430 (96%)	19 (4%)	0	100	100
1	B	456/458 (100%)	434 (95%)	22 (5%)	0	100	100
All	All	905/916 (99%)	864 (96%)	41 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/365 (98%)	358 (100%)	1 (0%)	92	91
1	B	365/365 (100%)	363 (100%)	2 (0%)	88	86
All	All	724/730 (99%)	721 (100%)	3 (0%)	91	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	GLU
1	B	52	GLU
1	B	304	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	312	GLN
1	B	356	GLN
1	B	396	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SIN	B	1002	-	1,7,7	0.07	0	2,8,8	1.32	0
3	GOL	A	1003	-	5,5,5	0.11	0	5,5,5	0.39	0
3	GOL	A	1002	-	5,5,5	0.12	0	5,5,5	0.24	0
2	PLP	B	1001	1	15,15,16	0.89	0	20,22,23	1.55	3 (15%)
3	GOL	A	1005	-	5,5,5	0.11	0	5,5,5	0.37	0
3	GOL	A	1004	-	5,5,5	0.10	0	5,5,5	0.19	0
2	PLP	A	1001	1	15,15,16	0.84	0	20,22,23	1.35	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SIN	B	1002	-	-	0/1/5/5	-
3	GOL	A	1003	-	-	2/4/4/4	-
3	GOL	A	1002	-	-	2/4/4/4	-
2	PLP	B	1001	1	-	0/6/6/8	0/1/1/1
3	GOL	A	1005	-	-	2/4/4/4	-
3	GOL	A	1004	-	-	2/4/4/4	-
2	PLP	A	1001	1	-	1/6/6/8	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1001	PLP	O4P-C5A-C5	4.48	117.89	109.35
2	A	1001	PLP	O4P-C5A-C5	3.80	116.60	109.35
2	B	1001	PLP	C4A-C4-C5	3.45	124.49	120.94
2	A	1001	PLP	C4A-C4-C5	2.63	123.64	120.94
2	B	1001	PLP	O4P-P-O1P	-2.21	100.28	106.47
2	A	1001	PLP	O3P-P-O4P	-2.16	100.99	106.73

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1003	GOL	C1-C2-C3-O3
3	A	1003	GOL	O2-C2-C3-O3
3	A	1004	GOL	C1-C2-C3-O3
3	A	1004	GOL	O2-C2-C3-O3
3	A	1005	GOL	O2-C2-C3-O3
3	A	1002	GOL	O1-C1-C2-C3
3	A	1005	GOL	C1-C2-C3-O3
2	A	1001	PLP	C5A-O4P-P-O1P
3	A	1002	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1003	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	451/458 (98%)	0.12	26 (5%) 23 22	22, 31, 50, 95	0
1	B	453/458 (98%)	-0.08	13 (2%) 51 50	20, 28, 43, 64	0
All	All	904/916 (98%)	0.02	39 (4%) 35 33	20, 29, 48, 95	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	456	PRO	7.9
1	A	192	ALA	5.8
1	A	6	SER	5.3
1	A	191	LEU	4.4
1	B	6	SER	4.2
1	A	323	PHE	3.2
1	A	198	HIS	3.0
1	A	251	TYR	3.0
1	A	193	GLY	2.9
1	A	154	VAL	2.8
1	B	260	VAL	2.8
1	B	209	GLN	2.7
1	B	154	VAL	2.7
1	B	323	PHE	2.5
1	A	252	ASP	2.5
1	A	324	THR	2.5
1	B	458	GLY	2.5
1	B	115	THR	2.4
1	A	121	ALA	2.4
1	A	238	ARG	2.4
1	B	117	SER	2.4
1	B	119	SER	2.4
1	A	115	THR	2.4
1	A	246	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	297	ALA	2.4
1	A	85	LEU	2.3
1	A	296	ILE	2.3
1	A	239	THR	2.3
1	B	57	LEU	2.3
1	A	137	GLY	2.2
1	A	260	VAL	2.2
1	A	390	VAL	2.2
1	A	297	ALA	2.2
1	B	284	VAL	2.1
1	A	155	THR	2.1
1	A	388	ASP	2.1
1	B	121	ALA	2.0
1	A	119	SER	2.0
1	A	117	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	1002	6/6	0.81	0.23	45,58,61,68	0
3	GOL	A	1003	6/6	0.82	0.28	42,52,60,70	0
5	NA	B	1003	1/1	0.90	0.33	45,45,45,45	0
3	GOL	A	1004	6/6	0.92	0.10	52,53,54,66	0
4	SIN	B	1002	8/8	0.93	0.18	39,44,50,52	0
3	GOL	A	1005	6/6	0.94	0.11	30,43,56,66	0
2	PLP	B	1001	15/16	0.96	0.14	21,26,32,32	0
2	PLP	A	1001	15/16	0.97	0.12	23,29,32,36	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NA	B	1004	1/1	0.97	0.14	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.