



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 03:27 pm BST

PDB ID : 6U36
Title : PCSK9 in complex with a Fab and compound 14
Authors : Lu, J.; Soisson, S.
Deposited on : 2019-08-21
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	
1	B	707	
2	H	246	
3	L	215	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	92	712	460	122	128	2	0	0	0
1	B	481	3463	2141	630	662	30	0	0	0

There are 94 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	474	ILE	VAL	engineered mutation	UNP Q8NBP7
A	670	GLU	GLY	engineered mutation	UNP Q8NBP7
A	693	LYS	-	expression tag	UNP Q8NBP7
A	694	GLY	-	expression tag	UNP Q8NBP7
A	695	ASN	-	expression tag	UNP Q8NBP7
A	696	SER	-	expression tag	UNP Q8NBP7
A	697	ALA	-	expression tag	UNP Q8NBP7
A	698	ASP	-	expression tag	UNP Q8NBP7
A	699	ILE	-	expression tag	UNP Q8NBP7
A	700	GLN	-	expression tag	UNP Q8NBP7
A	701	HIS	-	expression tag	UNP Q8NBP7
A	702	SER	-	expression tag	UNP Q8NBP7
A	703	GLY	-	expression tag	UNP Q8NBP7
A	704	GLY	-	expression tag	UNP Q8NBP7
A	705	ARG	-	expression tag	UNP Q8NBP7
A	706	SER	-	expression tag	UNP Q8NBP7
A	707	SER	-	expression tag	UNP Q8NBP7
A	708	LEU	-	expression tag	UNP Q8NBP7
A	709	GLU	-	expression tag	UNP Q8NBP7
A	710	GLY	-	expression tag	UNP Q8NBP7
A	711	PRO	-	expression tag	UNP Q8NBP7
A	712	ARG	-	expression tag	UNP Q8NBP7
A	713	PHE	-	expression tag	UNP Q8NBP7
A	714	GLU	-	expression tag	UNP Q8NBP7
A	715	GLY	-	expression tag	UNP Q8NBP7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	716	LYS	-	expression tag	UNP Q8NBP7
A	717	PRO	-	expression tag	UNP Q8NBP7
A	718	ILE	-	expression tag	UNP Q8NBP7
A	719	PRO	-	expression tag	UNP Q8NBP7
A	720	ASN	-	expression tag	UNP Q8NBP7
A	721	PRO	-	expression tag	UNP Q8NBP7
A	722	LEU	-	expression tag	UNP Q8NBP7
A	723	LEU	-	expression tag	UNP Q8NBP7
A	724	GLY	-	expression tag	UNP Q8NBP7
A	725	LEU	-	expression tag	UNP Q8NBP7
A	726	ASP	-	expression tag	UNP Q8NBP7
A	727	SER	-	expression tag	UNP Q8NBP7
A	728	THR	-	expression tag	UNP Q8NBP7
A	729	ARG	-	expression tag	UNP Q8NBP7
A	730	THR	-	expression tag	UNP Q8NBP7
A	731	GLY	-	expression tag	UNP Q8NBP7
A	732	HIS	-	expression tag	UNP Q8NBP7
A	733	HIS	-	expression tag	UNP Q8NBP7
A	734	HIS	-	expression tag	UNP Q8NBP7
A	735	HIS	-	expression tag	UNP Q8NBP7
A	736	HIS	-	expression tag	UNP Q8NBP7
A	737	HIS	-	expression tag	UNP Q8NBP7
B	474	ILE	VAL	engineered mutation	UNP Q8NBP7
B	670	GLU	GLY	engineered mutation	UNP Q8NBP7
B	693	LYS	-	expression tag	UNP Q8NBP7
B	694	GLY	-	expression tag	UNP Q8NBP7
B	695	ASN	-	expression tag	UNP Q8NBP7
B	696	SER	-	expression tag	UNP Q8NBP7
B	697	ALA	-	expression tag	UNP Q8NBP7
B	698	ASP	-	expression tag	UNP Q8NBP7
B	699	ILE	-	expression tag	UNP Q8NBP7
B	700	GLN	-	expression tag	UNP Q8NBP7
B	701	HIS	-	expression tag	UNP Q8NBP7
B	702	SER	-	expression tag	UNP Q8NBP7
B	703	GLY	-	expression tag	UNP Q8NBP7
B	704	GLY	-	expression tag	UNP Q8NBP7
B	705	ARG	-	expression tag	UNP Q8NBP7
B	706	SER	-	expression tag	UNP Q8NBP7
B	707	SER	-	expression tag	UNP Q8NBP7
B	708	LEU	-	expression tag	UNP Q8NBP7
B	709	GLU	-	expression tag	UNP Q8NBP7
B	710	GLY	-	expression tag	UNP Q8NBP7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	711	PRO	-	expression tag	UNP Q8NBP7
B	712	ARG	-	expression tag	UNP Q8NBP7
B	713	PHE	-	expression tag	UNP Q8NBP7
B	714	GLU	-	expression tag	UNP Q8NBP7
B	715	GLY	-	expression tag	UNP Q8NBP7
B	716	LYS	-	expression tag	UNP Q8NBP7
B	717	PRO	-	expression tag	UNP Q8NBP7
B	718	ILE	-	expression tag	UNP Q8NBP7
B	719	PRO	-	expression tag	UNP Q8NBP7
B	720	ASN	-	expression tag	UNP Q8NBP7
B	721	PRO	-	expression tag	UNP Q8NBP7
B	722	LEU	-	expression tag	UNP Q8NBP7
B	723	LEU	-	expression tag	UNP Q8NBP7
B	724	GLY	-	expression tag	UNP Q8NBP7
B	725	LEU	-	expression tag	UNP Q8NBP7
B	726	ASP	-	expression tag	UNP Q8NBP7
B	727	SER	-	expression tag	UNP Q8NBP7
B	728	THR	-	expression tag	UNP Q8NBP7
B	729	ARG	-	expression tag	UNP Q8NBP7
B	730	THR	-	expression tag	UNP Q8NBP7
B	731	GLY	-	expression tag	UNP Q8NBP7
B	732	HIS	-	expression tag	UNP Q8NBP7
B	733	HIS	-	expression tag	UNP Q8NBP7
B	734	HIS	-	expression tag	UNP Q8NBP7
B	735	HIS	-	expression tag	UNP Q8NBP7
B	736	HIS	-	expression tag	UNP Q8NBP7
B	737	HIS	-	expression tag	UNP Q8NBP7

- Molecule 2 is a protein called Fab Heavy Chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	213	1583	1005	263	309	6	0	0	0

- Molecule 3 is a protein called Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	210	1520	955	249	312	4	0	0	0

- Molecule 4 is 2-fluoro-4-([(1R)-6-(2-{4-[1-(4-methoxyphenyl)-5-methyl-6-oxo-1,6-dihydropyridazin-3-yl]-1H-1,2,3-triazol-1-yl}ethoxy)-1-methyl-1-{2-oxo-2-[(1,3-thiazol-2-yl)amino]eth

yl}-1,2,3,4-tetrahydroisoquinolin-7-yl]oxy}benzoic acid (three-letter code: PVM) (formula: $C_{38}H_{35}FN_8O_7S$) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
4	B	1	55	38	1	8	7	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			1	1		
5	B	19	Total	O	0	0
			19	19		
5	H	7	Total	O	0	0
			7	7		
5	L	1	Total	O	0	0
			1	1		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proprotein convertase subtilisin/kexin type 9





- Molecule 1: Proprotein convertase subtilisin/kexin type 9







• Molecule 2: Fab Heavy Chain



• Molecule 3: Fab Light Chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	155.60Å 155.60Å 152.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.67 – 2.70 36.67 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.67-2.70) 99.8 (36.67-2.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 2.68Å)	Xtrriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.240 , 0.251 0.234 , 0.249	Depositor DCC
R_{free} test set	2925 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	69.7	Xtrriage
Anisotropy	0.127	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.029 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7361	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PVM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/728	0.64	0/989
1	B	0.41	0/3525	0.69	0/4803
2	H	0.38	0/1625	0.73	0/2217
3	L	0.38	0/1558	0.68	0/2140
All	All	0.40	0/7436	0.69	0/10149

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	712	0	708	1	0
1	B	3463	0	3299	24	0
2	H	1583	0	1500	13	0
3	L	1520	0	1363	9	0
4	B	55	0	0	0	0
5	A	1	0	0	0	0
5	B	19	0	0	0	0
5	H	7	0	0	0	0
5	L	1	0	0	0	0
All	All	7361	0	6870	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:344:GLN:HE22	1:B:425:ASN:H	1.38	0.69
1:B:185:LEU:HD11	1:B:271:ILE:HD11	1.77	0.67
3:L:6:GLN:HE22	3:L:87:TYR:HA	1.64	0.63
1:B:468:THR:HB	1:B:471:ALA:HB2	1.80	0.62
1:B:200:VAL:HG22	1:B:247:MET:HB2	1.83	0.60
3:L:6:GLN:HE21	3:L:100:GLY:HA3	1.68	0.59
1:B:183:TYR:HE1	1:B:248:ARG:HD3	1.69	0.58
3:L:34:ASN:HD22	3:L:89:GLN:NE2	2.04	0.56
2:H:12:VAL:HG11	2:H:86:LEU:HD12	1.88	0.55
3:L:34:ASN:HD22	3:L:89:GLN:HE22	1.52	0.55
3:L:149:TRP:HE1	3:L:160:SER:HG	1.56	0.54
1:B:193:HIS:HD2	1:B:195:GLU:H	1.56	0.54
2:H:38:ARG:HB3	2:H:48:ILE:HD11	1.90	0.53
1:B:157:ASN:ND2	1:B:368:ILE:HG23	2.24	0.53
2:H:124:PRO:HB3	2:H:150:TYR:HB3	1.91	0.53
1:B:156:TRP:CH2	1:B:364:PRO:HB3	2.45	0.51
2:H:30:SER:HA	2:H:53:PRO:HB2	1.91	0.51
2:H:22:CYS:HB3	2:H:79:LEU:HB3	1.93	0.50
1:B:157:ASN:HD22	1:B:368:ILE:HG23	1.76	0.49
1:B:155:PRO:HB2	1:B:157:ASN:OD1	2.13	0.49
2:H:186:VAL:HG11	3:L:136:LEU:HD22	1.96	0.47
1:A:65:HIS:CE1	1:B:295:ARG:HH11	2.33	0.47
1:B:484:LEU:HB3	1:B:560:THR:HB	1.97	0.47
1:B:472:THR:HG21	1:B:510:ARG:NH1	2.29	0.46
1:B:537:HIS:CE1	1:B:554:GLN:HE22	2.33	0.46
1:B:612:GLU:HG2	1:B:675:VAL:HG22	1.98	0.46
1:B:616:PRO:HA	1:B:671:ALA:HA	1.98	0.46
2:H:200:ILE:HG22	2:H:215:LYS:HA	1.98	0.46
2:H:173:ALA:HB2	2:H:183:LEU:HD12	1.99	0.44
1:B:345:PRO:HD3	1:B:424:ILE:HG23	2.00	0.43
3:L:123:ASP:HA	3:L:126:LEU:HD12	2.01	0.43
2:H:36:TRP:CE2	2:H:81:LEU:HB2	2.54	0.42
2:H:34:MET:HB3	2:H:79:LEU:HD22	2.02	0.42
1:B:230:LEU:HD11	1:B:386:SER:HB3	2.02	0.42
1:B:410:GLU:HA	1:B:528:LEU:HD11	2.01	0.41
1:B:155:PRO:HD2	1:B:158:LEU:HD12	2.01	0.41
2:H:100:ARG:HD2	2:H:103:TYR:HB2	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:468:THR:HG22	1:B:470:MET:H	1.85	0.41
3:L:117:PHE:HD2	3:L:136:LEU:HD23	1.86	0.41
3:L:92:ASP:HB3	3:L:95:PRO:O	2.21	0.41
1:B:195:GLU:HG3	1:B:236:GLY:HA2	2.03	0.40
2:H:91:THR:HG23	2:H:115:THR:HA	2.03	0.40
2:H:153:GLU:HG2	2:H:154:PRO:HA	2.02	0.40
1:B:183:TYR:HB2	1:B:285:VAL:HG22	2.04	0.40
1:B:330:ALA:O	1:B:333:VAL:HG22	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	90/707 (13%)	88 (98%)	2 (2%)	0	100	100
1	B	467/707 (66%)	452 (97%)	15 (3%)	0	100	100
2	H	209/246 (85%)	204 (98%)	5 (2%)	0	100	100
3	L	208/215 (97%)	197 (95%)	10 (5%)	1 (0%)	29	54
All	All	974/1875 (52%)	941 (97%)	32 (3%)	1 (0%)	51	78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	L	51	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	74/571 (13%)	74 (100%)	0	100	100
1	B	358/571 (63%)	357 (100%)	1 (0%)	92	98
2	H	168/203 (83%)	165 (98%)	3 (2%)	59	83
3	L	157/186 (84%)	156 (99%)	1 (1%)	86	95
All	All	757/1531 (49%)	752 (99%)	5 (1%)	84	94

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	421	LYS
2	H	72	ARG
2	H	100	ARG
2	H	209	ASN
3	L	109	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	65	HIS
1	A	113	HIS
1	B	193	HIS
1	B	278	GLN
1	B	344	GLN
1	B	464	HIS
1	B	503	GLN
1	B	554	GLN
2	H	209	ASN
3	L	6	GLN
3	L	89	GLN
3	L	139	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PVM	B	801	-	53,61,61	1.34	6 (11%)	55,88,88	1.61	10 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PVM	B	801	-	-	6/25/46/46	0/7/7/7

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	801	PVM	C33-C32	4.67	1.47	1.39
4	B	801	PVM	C32-N37	-3.72	1.29	1.33
4	B	801	PVM	C35-C34	3.53	1.49	1.41
4	B	801	PVM	C18-C1	-2.40	1.51	1.53
4	B	801	PVM	C32-C29	-2.04	1.44	1.49
4	B	801	PVM	C17-C10	-2.01	1.49	1.51

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	B	801	PVM	C47-O46-C43	-4.48	107.80	117.51
4	B	801	PVM	C33-C32-C29	-3.96	115.82	120.98
4	B	801	PVM	C32-C33-C34	-3.74	116.78	121.16
4	B	801	PVM	C33-C34-C35	3.47	119.50	115.15
4	B	801	PVM	O-C25-C26	-2.77	104.45	107.28
4	B	801	PVM	C32-C29-N30	2.39	124.84	120.88
4	B	801	PVM	C4-C3-C8	-2.39	115.13	119.91
4	B	801	PVM	C-C4-C3	2.21	114.63	110.69
4	B	801	PVM	C21-C22-C23	2.17	119.23	116.10
4	B	801	PVM	F-C23-C22	2.08	122.03	118.89

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	801	PVM	C28-C29-C32-C33
4	B	801	PVM	C28-C29-C32-N37
4	B	801	PVM	C42-C43-O46-C47
4	B	801	PVM	C44-C43-O46-C47
4	B	801	PVM	N-C1-C17-C10
4	B	801	PVM	C18-C1-C17-C10

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	92/707 (13%)	0.18	1 (1%) 80 82	49, 69, 91, 97	0
1	B	481/707 (68%)	0.37	16 (3%) 46 46	48, 63, 85, 97	0
2	H	213/246 (86%)	0.23	13 (6%) 21 20	58, 72, 111, 130	0
3	L	210/215 (97%)	0.61	32 (15%) 2 1	61, 91, 122, 127	0
All	All	996/1875 (53%)	0.37	62 (6%) 20 19	48, 69, 114, 130	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	L	195	CYS	5.8
2	H	143	LEU	5.5
1	B	547	GLY	4.9
3	L	154	ALA	4.9
3	L	118	ILE	4.6
3	L	197	VAL	4.0
3	L	153	ASN	3.9
1	B	548	THR	3.9
2	H	144	GLY	3.8
3	L	155	LEU	3.7
1	B	313	THR	3.6
3	L	77	SER	3.4
2	H	131	PRO	3.3
3	L	194	ALA	3.3
3	L	192	VAL	3.3
3	L	147	VAL	3.2
2	H	200	ILE	3.2
2	H	199	TYR	3.2
1	B	221	SER	3.1
1	B	545	SER	3.1
3	L	185	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	139	HIS	2.9
1	B	642	SER	2.8
3	L	201	GLY	2.7
3	L	149	TRP	2.7
3	L	152	ASP	2.7
3	L	80	PRO	2.7
2	H	140	THR	2.6
3	L	109	ARG	2.6
2	H	196	THR	2.6
3	L	117	PHE	2.6
1	B	543	GLU	2.5
3	L	198	THR	2.5
1	B	333	VAL	2.5
3	L	193	TYR	2.5
3	L	116	VAL	2.5
3	L	137	LEU	2.4
3	L	78	LEU	2.4
2	H	139	GLY	2.4
2	H	198	THR	2.3
1	B	328	ALA	2.3
3	L	16	GLY	2.3
3	L	150	LYS	2.3
1	B	348	LEU	2.3
1	B	641	THR	2.3
1	B	544	ALA	2.2
1	B	183	TYR	2.2
3	L	204	SER	2.2
3	L	107	ILE	2.2
1	B	361	LEU	2.2
3	L	112	ALA	2.2
1	B	335	THR	2.2
3	L	190	HIS	2.2
3	L	19	ALA	2.2
1	B	359	VAL	2.1
3	L	209	SER	2.1
2	H	160	ASN	2.1
3	L	105	VAL	2.1
3	L	186	ASP	2.1
2	H	216	VAL	2.0
2	H	218	PRO	2.0
2	H	138	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PVM	B	801	55/55	0.88	0.24	49,55,70,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers

There are no such residues in this entry.