



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2023 – 06:51 AM EDT

PDB ID : 6VGE  
Title : Crystal structure of the DNA binding domains of human transcription factor ERG, human Runx2 bound to core binding factor beta (Cbfb), in complex with 16mer DNA CAGAGGATGTGGCTTC  
Authors : Hou, C.; Tsodikov, O.V.  
Deposited on : 2020-01-07  
Resolution : 4.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**  
Xtrriage (Phenix) : 1.13  
EDS : **FAILED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 4.25 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 3361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator ERG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	98	806	512	143	146	5	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	302	GLY	-	expression tag	UNP P11308
A	303	PRO	-	expression tag	UNP P11308
A	304	HIS	-	expression tag	UNP P11308
A	305	MET	-	expression tag	UNP P11308
A	420	SER	-	expression tag	UNP P11308
A	421	TYR	-	expression tag	UNP P11308
A	422	HIS	-	expression tag	UNP P11308
A	423	ALA	-	expression tag	UNP P11308
A	424	HIS	-	expression tag	UNP P11308
A	425	PRO	-	expression tag	UNP P11308
A	426	GLN	-	expression tag	UNP P11308
A	427	LYS	-	expression tag	UNP P11308
A	428	MET	-	expression tag	UNP P11308
A	429	ASN	-	expression tag	UNP P11308

- Molecule 2 is a DNA chain called DNA (5'-D(P\*CP\*AP\*GP\*AP\*GP\*GP\*AP\*TP\*GP\*TP\*GP\*GP\*CP\*TP\*TP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	16	332	157	62	97	16	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(P\*GP\*AP\*AP\*GP\*CP\*CP\*AP\*CP\*AP\*TP\*CP\*CP\*TP\*CP\*TP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	16	Total	C	N	O	P	0	0	0
			324	154	59	95	16			

- Molecule 4 is a protein called Runt-related transcription factor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	117	Total	C	N	O	S	0	0	0
			901	567	164	166	4			

- Molecule 5 is a protein called Core-binding factor subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	G	121	Total	C	N	O	S	0	0	0
			998	622	181	189	6			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	-13	MET	-	initiating methionine	UNP Q13951
G	-12	GLY	-	expression tag	UNP Q13951
G	-11	SER	-	expression tag	UNP Q13951
G	-10	SER	-	expression tag	UNP Q13951
G	-9	HIS	-	expression tag	UNP Q13951
G	-8	HIS	-	expression tag	UNP Q13951
G	-7	HIS	-	expression tag	UNP Q13951
G	-6	HIS	-	expression tag	UNP Q13951
G	-5	HIS	-	expression tag	UNP Q13951
G	-4	HIS	-	expression tag	UNP Q13951
G	-3	SER	-	expression tag	UNP Q13951
G	-2	GLN	-	expression tag	UNP Q13951
G	-1	ASP	-	expression tag	UNP Q13951
G	0	PRO	-	expression tag	UNP Q13951

MolProbity and EDS failed to run properly - this section is therefore empty.

### 3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.92Å 103.92Å 322.91Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.00 – 4.25	Depositor
% Data completeness (in resolution range)	95.0 (35.00-4.25)	Depositor
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.70 (at 4.28Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.312 , 0.317	Depositor
Wilson B-factor (Å <sup>2</sup> )	194.8	Xtrriage
Anisotropy	0.477	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3361	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	303.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 4 Model quality [i](#)

### 4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles [i](#)

#### 4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 4.7 Other polymers [i](#)

There are no such residues in this entry.

## 4.8 Polymer linkage issues

There are no chain breaks in this entry.

## 5 Fit of model and data

### 5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

### 5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers

EDS failed to run properly - this section is therefore empty.