

wwPDB EM Validation Summary Report (i)

Dec 18, 2022 – 02:14 am GMT

PDB ID : 7A02

EMDB ID : EMD-11591

Title : Bacillus endospore appendages form a novel family of disulfide-linked pili Authors : Pradhan, B.; Liedtke, J.; Sleutel, M.; Lindback, T.; Llarena, A.K.; Brynild-

srud, O.; Aspholm, M.; Remaut, H.

Deposited on : 2020-08-06

Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

MolProbity : 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	EM structures	
Metric	$(\# \mathrm{Entries})$	$(\# \mathrm{Entries})$	
Clashscore	158937	4297	
Ramachandran outliers	154571	4023	
Sidechain outliers	154315	3826	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	117	82%	14%	•
1	В	117	79%	16%	·
1	С	117	79%	17%	•
1	D	117	82%	14%	•
1	Е	117	78%	18%	•
1	F	117	84%	12%	.
1	G	117	80%	15%	.
1	Н	117	80%	15%	.

Continued on next page...



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	I	117	•	
т	1	111		•
1	J	117	82% 14%	·
1	K	117	79% 16%	•
1	L	117	79% 16%	
1	M	117	81% 15%	-
	111	111	•	•
1	N	117	81% 15%	•
1	О	117	81% 15%	
1	Р	117	79% 16%	•
1	Q	117	79% 17%	<u>.</u>
			- 15% - 17%	
1	R	117	79% 16%	•
1	S	117	78% 18%	
1	Т	117	80% 15%	.
1	U		9%	
1	U	117	81% 15% 7%_	•
1	V	117	81% 15%	•
1	W	117	81% 15%	·



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 18699 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DUF3992 domain-containing protein.

1		Residues		At	\mathbf{oms}			AltConf	Trace	
	٨	110	Total	С	N	О	S	0	0	
1	A	112	813	512	130	166	5	0	0	
1	D	110	Total	С	N	О	S	0	0	
1	В	112	813	512	130	166	5	0	0	
1	C	110	Total	С	N	О	S	0	0	
1	С	112	813	512	130	166	5	0	0	
1	D	110	Total	С	N	О	S	0	0	
1	D	112	813	512	130	166	5	U	0	
1	Е	110	Total	С	N	О	S	0	0	
1	Ŀ	112	813	512	130	166	5	U	0	
1	F	110	Total	С	N	О	S	0	0	
1	Г	112	813	512	130	166	5	U		
1	G	110	Total	С	N	О	S	0	0	
1	G	112	813	512	130	166	5	0	0	
1	Н	110	Total	С	N	О	S	0	0	
1	11	112	813	512	130	166	5	U		
1	I	110	Total	С	N	О	S	0	0	
1	1	112	813	512	130	166	5	U		
1	J	112	Total	С	N	О	S	0	0	
1	J	112	813	512	130	166	5	U		
1	K	110	Total	С	N	О	S	0	0	
1	N	112	813	512	130	166	5	U	0	
1	L	112	Total	С	N	О	S	0	0	
1	Ъ	112	813	512	130	166	5	U		
1	M	112	Total	С	N	О	S	0	0	
1	IVI	112	813	512	130	166	5	0		
1	N	112	Total	С	N	О	S	0	0	
1	IN	112	813	512	130	166	5	0		
1	O 112	119	Total	С	N	O	S	0	0	
1		112	813	512	130	166	5	0		
1	P 112	112	Total	С	N	О	S	0	0	
1	1	114	813	512	130	166	5	U	0	
1	Q	112	Total	С	N	О	S	0	0	
1	Q	112	813	512	130	166	5			



Continued from previous page...


Mol	Chain	Residues		At	oms			AltConf	Trace										
1	R	112	Total	С	N	О	S	0	0										
1	11	112	813	512	130	166	5	0	0										
1	S	112	Total	С	N	О	S	0	0										
1	b	112	813	512	130	166	5	0	0										
1	Т	112	Total	С	N	О	S	0	0										
1	1	112	813	512	130	166	5	U	0										
1	IJ	112	Total	С	N	О	S	0	0										
1		112	813	512	130	166	5	0	0										
1	V	17	1/	1/	V	17	V	V	17	V	V	V 112	Total	С	N	О	S	0	0
1	v	112	813	512	130	166	5	0	0										
1	W	112	Total	С	N	О	S	0	0										
1	V V	112	813	512	130	166	5												




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DUF3992 domain-containing protein












• Molecule 1: DUF3992 domain-containing protein







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=3.43721°, rise=32.3504 Å,	Depositor
	axial sym=C1	
Number of segments used	65466	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	JEOL CRYO ARM 300	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	64.66	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	60000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	8.824	Depositor
Minimum map value	-5.849	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	0.65	Depositor
Map size (Å)	119.168, 119.168, 130.928	wwPDB
Map dimensions	167, 152, 152	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.784, 0.784, 0.784	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.44	0/824	0.55	0/1126	
1	В	0.44	0/824	0.55	0/1126	
1	С	0.44	0/824	0.55	0/1126	
1	D	0.44	0/824	0.55	0/1126	
1	Е	0.44	0/824	0.55	0/1126	
1	F	0.44	0/824	0.55	0/1126	
1	G	0.44	0/824	0.55	0/1126	
1	Н	0.44	0/824	0.55	0/1126	
1	I	0.44	0/824	0.55	0/1126	
1	J	0.44	0/824	0.55	0/1126	
1	K	0.44	0/824	0.55	0/1126	
1	L	0.44	0/824	0.55	0/1126	
1	M	0.44	0/824	0.55	0/1126	
1	N	0.44	0/824	0.55	0/1126	
1	О	0.44	0/824	0.55	0/1126	
1	Р	0.44	0/824	0.55	0/1126	
1	Q	0.44	0/824	0.55	0/1126	
1	R	0.44	0/824	0.55	0/1126	
1	S	0.44	0/824	0.55	0/1126	
1	Т	0.44	0/824	0.55	0/1126	
1	U	0.44	0/824	0.55	0/1126	
1	V	0.44	0/824	0.55	0/1126	
1	W	0.44	0/824	0.55	0/1126	
All	All	0.44	0/18952	0.55	0/25898	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	813	0	801	12	0
1	В	813	0	802	22	0
1	С	813	0	801	18	0
1	D	813	0	801	17	0
1	Ε	813	0	801	19	0
1	F	813	0	801	13	0
1	G	813	0	801	15	0
1	Н	813	0	801	13	0
1	I	813	0	801	14	0
1	J	813	0	800	14	0
1	K	813	0	800	15	0
1	L	813	0	799	19	0
1	M	813	0	799	15	0
1	N	813	0	800	17	0
1	О	813	0	801	13	0
1	Р	813	0	801	18	0
1	Q	813	0	801	19	0
1	R	813	0	801	21	0
1	S	813	0	801	21	0
1	Т	813	0	801	15	0
1	U	813	0	801	12	0
1	V	813	0	801	14	0
1	W	813	0	801	15	0
All	All	18699	0	18417	300	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 300 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:R:90:ARG:HH22	1:S:43:GLN:HE22	1.19	0.88
1:R:90:ARG:HH22	1:S:43:GLN:NE2	1.71	0.87
1:D:90:ARG:HH22	1:E:43:GLN:NE2	1.79	0.81
1:F:90:ARG:HH22	1:G:43:GLN:NE2	1.80	0.80



Continued from previous page...

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:B:90:ARG:HH22	1:C:43:GLN:NE2	1.84	0.74

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	В	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	С	110/117~(94%)	100 (91%)	10 (9%)	0	100	100
1	D	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	E	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	F	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	G	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	Н	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	I	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	J	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	K	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	L	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	M	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	N	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	О	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	Р	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	Q	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	R	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	S	110/117 (94%)	101 (92%)	9 (8%)	0	100	100



I 'omtamalod	trom	mmonia	maaa
Continued	11 0116	DICUIUUS	Daue
	.,	10	1

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	${ m T}$	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	U	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	V	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
1	W	110/117 (94%)	101 (92%)	9 (8%)	0	100	100
All	All	2530/2691 (94%)	2322 (92%)	208 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	$87/92\ (95\%)$	87 (100%)	0	100	100
1	В	$87/92\ (95\%)$	87 (100%)	0	100	100
1	С	$87/92\ (95\%)$	87 (100%)	0	100	100
1	D	$87/92\ (95\%)$	87 (100%)	0	100	100
1	E	$87/92\ (95\%)$	87 (100%)	0	100	100
1	F	$87/92\ (95\%)$	87 (100%)	0	100	100
1	G	$87/92\ (95\%)$	87 (100%)	0	100	100
1	Н	$87/92\ (95\%)$	87 (100%)	0	100	100
1	I	87/92 (95%)	87 (100%)	0	100	100
1	J	$87/92\ (95\%)$	87 (100%)	0	100	100
1	K	$87/92\ (95\%)$	87 (100%)	0	100	100
1	L	$87/92\ (95\%)$	87 (100%)	0	100	100
1	M	87/92 (95%)	87 (100%)	0	100	100
1	N	$87/92\ (95\%)$	87 (100%)	0	100	100
1	О	87/92 (95%)	87 (100%)	0	100	100
1	Р	87/92 (95%)	87 (100%)	0	100	100
1	Q	87/92 (95%)	87 (100%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	R	$87/92\ (95\%)$	87 (100%)	0	100	100
1	S	87/92 (95%)	87 (100%)	0	100	100
1	Т	87/92 (95%)	87 (100%)	0	100	100
1	U	87/92 (95%)	87 (100%)	0	100	100
1	V	87/92 (95%)	87 (100%)	0	100	100
1	W	87/92 (95%)	87 (100%)	0	100	100
All	All	2001/2116 (95%)	2001 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	S	43	GLN
1	Т	43	GLN
1	N	43	GLN
1	О	43	GLN
1	Q	43	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-11591. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map








The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map







X Index: 49


Y Index: 53


Z Index: 92

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map







The images above show the 3D surface view of the map at the recommended contour level 0.65. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is $205~\mathrm{nm}^3$; this corresponds to an approximate mass of $185~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)


This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.333 $\rm \AA^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
rtesolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	3.00	-	-	
Author-provided FSC curve	2.98	3.55	3.06	
Unmasked-calculated*	-	-	-	


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-11591 and PDB model 7A02. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.65 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.65).



9.4 Atom inclusion (i)



At the recommended contour level, 94% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.65) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8805	0.5360
A	0.8876	0.5430
В	0.8764	0.5340
С	0.8714	0.5320
D	0.8976	0.5390
Е	0.8939	0.5400
F	0.9051	0.5510
G	0.8926	0.5450
Н	0.8801	0.5260
I	0.8939	0.5500
J	0.8926	0.5450
K	0.9014	0.5560
L	0.9026	0.5490
M	0.8926	0.5420
N	0.8851	0.5440
О	0.8914	0.5350
Р	0.8814	0.5330
Q	0.8826	0.5370
R	0.8801	0.5440
S	0.8801	0.5420
Т	0.8427	0.5180
U	0.8165	0.5010
V	0.8277	0.4970
W	0.8764	0.5270



