



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 20, 2023 – 11:30 PM JST

PDB ID : 7ELX  
Title : The crystal structure of CTLA-4 and Fab  
Authors : Yu, X.J.; Wang, L.; Yu, C.F.  
Deposited on : 2021-04-12  
Resolution : 2.14 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2523 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	467	 2% 45% 55%
1	h	467	 4% 44% 55%
2	L	235	 89% 10%
2	l	235	 12% 87% 10%
3	C	126	 17% 91% 9%
3	c	126	 12% 90% 9%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8836 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called heavy chain of Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	211	Total 1608	C 1021	N 272	O 309	S 6	2	0	0
1	h	212	Total 1614	C 1024	N 273	O 311	S 6	2	0	0

- Molecule 2 is a protein called light chain of Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	212	Total 1633	C 1026	N 276	O 327	S 4	1	0	0
2	l	212	Total 1633	C 1026	N 276	O 327	S 4	1	0	0

- Molecule 3 is a protein called Cytotoxic T-lymphocyte protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	c	115	Total 858	C 541	N 142	O 166	S 9	1	0	0
3	C	115	Total 858	C 541	N 142	O 166	S 9	0	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	141	Total 141	O 141	0	0
4	L	178	Total 178	O 178	0	0
4	h	105	Total 105	O 105	0	0
4	l	133	Total 133	O 133	0	0

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
4	c	40	Total O 40 40	0	0
4	C	35	Total O 35 35	0	0









• Molecule 2: light chain of Fab



• Molecule 3: Cytotoxic T-lymphocyte protein 4



• Molecule 3: Cytotoxic T-lymphocyte protein 4



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.92Å 196.92Å 148.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.20 – 2.14 45.64 – 2.14	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.20-2.14) 99.8 (45.64-2.14)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.19 (at 2.14Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.177 , 0.217 0.178 , 0.180	Depositor DCC
$R_{free}$ test set	3853 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.8	Xtrriage
Anisotropy	0.639	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 51.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8836	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	H	0.64	1/1650 (0.1%)	0.75	4/2250 (0.2%)
1	h	0.63	2/1656 (0.1%)	0.79	4/2258 (0.2%)
2	L	0.63	1/1671 (0.1%)	0.74	2/2270 (0.1%)
2	l	0.55	0/1671	1.01	8/2270 (0.4%)
3	C	0.48	0/871	0.73	0/1183
3	c	0.47	0/871	0.70	0/1183
All	All	0.59	4/8390 (0.0%)	0.81	18/11414 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	h	0	1
3	c	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	206	LYS	CD-CE	8.43	1.72	1.51
1	h	211	LYS	CD-CE	6.73	1.68	1.51
1	h	211	LYS	CE-NZ	6.61	1.65	1.49
2	L	195	CYS	CB-SG	-6.56	1.71	1.82

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	l	78	ARG	NE-CZ-NH1	-24.18	108.21	120.30
2	l	78	ARG	NE-CZ-NH2	19.05	129.83	120.30
1	h	214	LYS	N-CA-CB	9.35	127.43	110.60
1	h	214	LYS	CA-CB-CG	7.61	130.14	113.40

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	l	155	LEU	CA-CB-CG	7.21	131.88	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	c	83	GLY	Peptide
1	h	131	PRO	Peptide

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	207/467 (44%)	207 (100%)	0	0	100	100
1	h	208/467 (44%)	204 (98%)	4 (2%)	0	100	100
2	L	210/235 (89%)	206 (98%)	4 (2%)	0	100	100
2	l	210/235 (89%)	205 (98%)	5 (2%)	0	100	100
3	C	111/126 (88%)	104 (94%)	7 (6%)	0	100	100
3	c	111/126 (88%)	107 (96%)	4 (4%)	0	100	100
All	All	1057/1656 (64%)	1033 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	179/413 (43%)	178 (99%)	1 (1%)	86	89
1	h	180/413 (44%)	179 (99%)	1 (1%)	86	89
2	L	183/203 (90%)	182 (100%)	1 (0%)	88	91
2	l	183/203 (90%)	179 (98%)	4 (2%)	52	53
3	C	94/105 (90%)	94 (100%)	0	100	100
3	c	94/105 (90%)	94 (100%)	0	100	100
All	All	913/1442 (63%)	906 (99%)	7 (1%)	81	85

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	l	53	PHE
2	l	78	ARG
2	l	155	LEU
2	l	153	ASN
1	h	182	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	41	GLN
3	C	75	ASN
3	C	76	GLN
2	l	159	ASN
1	H	110	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	H	211/467 (45%)	0.18	9 (4%) 35 43	28, 40, 64, 83	1 (0%)
1	h	212/467 (45%)	0.41	19 (8%) 9 12	30, 45, 83, 110	1 (0%)
2	L	212/235 (90%)	0.06	1 (0%) 91 93	24, 35, 52, 71	2 (0%)
2	l	212/235 (90%)	0.62	29 (13%) 3 3	26, 46, 96, 112	2 (0%)
3	C	115/126 (91%)	0.96	21 (18%) 1 1	33, 56, 105, 133	0
3	c	115/126 (91%)	0.69	15 (13%) 3 4	35, 55, 94, 104	1 (0%)
All	All	1077/1656 (65%)	0.43	94 (8%) 10 13	24, 44, 87, 133	7 (0%)

The worst 5 of 94 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	57	GLY	9.1
3	c	84	LEU	9.0
1	h	195	GLY	8.2
1	h	132	SER	7.9
2	l	155	LEU	7.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.