



wwPDB EM Validation Summary Report ⓘ

Jun 5, 2021 – 06:55 am BST

PDB ID : 7NW0
EMDB ID : EMD-12619
Title : RNA polymerase II pre-initiation complex with open promoter DNA
Authors : Aibara, S.; Schilbach, S.; Cramer, P.
Deposited on : 2021-03-16
Resolution : 6.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.0.dev75
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.19

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 6.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	0	760	
2	1	548	
3	2	462	
4	3	309	
5	4	308	
6	5	71	
7	6	395	
8	7	782	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	A	1970	11% 63% 9% 28%
10	B	1174	16% 87% 10%
11	C	275	17% 82% 12% 7%
12	D	142	58% 86% 10%
13	E	210	14% 87% 13%
14	F	127	6% 57% 5% 38%
15	G	172	28% 88% 11%
16	H	150	25% 90% 9%
17	I	125	9% 78% 14% 9%
18	J	67	19% 87% 9%
19	K	117	24% 86% 12%
20	L	58	24% 67% 9% 24%
21	M	316	19% 87% 7% 6%
22	N	106	7% 53% 9% 38%
23	O	339	21% 47% 5% 47%
24	Q	517	12% 24% 73%
25	R	249	36% 80% 10% 11%
26	T	106	10% 54% 10% 36%
27	U	376	30% 28% 70%
28	V	109	90% 82% 9% 9%
29	W	439	23% 39% 7% 54%
30	X	291	31% 52% 7% 41%
31	Y	16	81% 81% 19%
32	Z	8	100% 100%

2 Entry composition [i](#)

There are 37 unique types of molecules in this entry. The entry contains 66935 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TFIID basal transcription factor complex helicase XPD subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	0	714	5751	3683	999	1040	29	0	0

- Molecule 2 is a protein called General transcription factor IID subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	1	265	2167	1382	378	395	12	0	0

- Molecule 3 is a protein called General transcription factor IID subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	2	390	3158	2050	545	551	12	0	0

- Molecule 4 is a protein called CDK-activating kinase assembly factor MAT1.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
4	3	44	382	245	61	76	0	0

- Molecule 5 is a protein called General transcription factor IID subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	4	263	2066	1323	344	380	19	0	0

- Molecule 6 is a protein called General transcription factor IID subunit 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	5	66	523	337	83	100	3	0	0

- Molecule 7 is a protein called General transcription factor IIIH subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	6	329	2567	1621	440	479	27	0	0

- Molecule 8 is a protein called General transcription and DNA repair factor IIIH helicase subunit XPB.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	7	605	4890	3127	848	885	30	0	0

- Molecule 9 is a protein called RPB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	A	1423	11274	7092	2016	2094	72	0	0

- Molecule 10 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	B	1136	9076	5739	1597	1676	64	0	0

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	C	257	2059	1294	351	408	6	0	0

- Molecule 12 is a protein called RPOL4c domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	D	128	1050	656	178	212	4	0	0

- Molecule 13 is a protein called DNA-directed RNA polymerase II subunit E.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	E	209	1721	1089	300	324	8	0	0

- Molecule 14 is a protein called DNA-directed RNA polymerase II subunit F.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	F	79	Total	C	N	O	S	0	0
			636	406	108	117	5		

- Molecule 15 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	G	171	Total	C	N	O	S	0	0
			1351	875	219	249	8		

- Molecule 16 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	H	148	Total	C	N	O	S	0	0
			1186	750	194	237	5		

- Molecule 17 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	I	114	Total	C	N	O	S	0	0
			928	571	166	180	11		

- Molecule 18 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	J	64	Total	C	N	O	S	0	0
			507	328	86	87	6		

- Molecule 19 is a protein called RNA_pol_L_2 domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	K	115	Total	C	N	O	S	0	0
			920	593	152	173	2		

- Molecule 20 is a protein called RNA polymerase II subunit K.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	L	44	Total	C	N	O	S	0	0
			373	231	72	64	6		

- Molecule 21 is a protein called Transcription initiation factor IIB.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
21	M	298	2301	1435	409	439	18	0	0

- Molecule 22 is a DNA chain called Non-template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
22	N	66	1369	646	260	398	65	0	0

- Molecule 23 is a protein called TATA-box-binding protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
23	O	179	1422	923	251	241	7	0	0

- Molecule 24 is a protein called General transcription factor IIF subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
24	Q	138	1138	719	208	208	3	0	0

- Molecule 25 is a protein called General transcription factor IIF subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
25	R	222	1788	1127	320	338	3	0	0

- Molecule 26 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
26	T	68	1358	642	246	402	68	0	0

- Molecule 27 is a protein called Transcription initiation factor IIA subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
27	U	113	930	585	152	189	4	0	0

- Molecule 28 is a protein called Transcription initiation factor IIA subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
28	V	99	806	510	142	151	3	0	0

- Molecule 29 is a protein called General transcription factor IIE subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
29	W	202	1659	1042	299	307	11	0	0

- Molecule 30 is a protein called Transcription initiation factor IIE subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
30	X	171	1403	895	243	261	4	0	0


- Molecule 31 is a protein called Unassigned peptide, likely TFIIIE-Beta.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
31	Y	16	80	48	16	16	0	0

- Molecule 32 is a protein called Unassigned peptide, likely XPB.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
32	Z	8	40	24	8	8	0	0

- Molecule 33 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms		AltConf
33	0	1	Total	Fe S	0
			8	4 4	

- Molecule 34 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
34	4	2	Total	Zn	0
			2	2	
34	6	3	Total	Zn	0
			3	3	
34	A	2	Total	Zn	0
			2	2	
34	B	1	Total	Zn	0
			1	1	
34	C	1	Total	Zn	0
			1	1	
34	I	2	Total	Zn	0
			2	2	
34	J	1	Total	Zn	0
			1	1	
34	L	1	Total	Zn	0
			1	1	
34	M	1	Total	Zn	0
			1	1	
34	W	1	Total	Zn	0
			1	1	

- Molecule 35 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:

C₁₀H₁₅N₅O₁₀P₂).




Mol	Chain	Residues	Atoms				AltConf	
35	7	1	Total	C	N	O	P	0
			27	10	5	10	2	

- Molecule 36 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
36	7	1	Total	Mg	0
			1	1	
36	A	1	Total	Mg	0
			1	1	

- Molecule 37 is BERYLLIUM TRIFLUORIDE ION (three-letter code: BEF) (formula: BeF₃).




Mol	Chain	Residues	Atoms			AltConf
			Total	Be	F	
37	7	1	4	1	3	0

3 Residue-property plots



These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TFIID basal transcription factor complex helicase XPD subunit







• Molecule 2: General transcription factor IIH subunit 1




• Molecule 3: General transcription factor IIH subunit 4









• Molecule 4: CDK-activating kinase assembly factor MAT1





• Molecule 5: General transcription factor IIH subunit 3











• Molecule 6: General transcription factor IIH subunit 5




• Molecule 7: General transcription factor IIH subunit 2





• Molecule 8: General transcription and DNA repair factor IIH helicase subunit XPB








• Molecule 9: RPB1







- Molecule 10: DNA-directed RNA polymerase subunit beta







• Molecule 11: DNA-directed RNA polymerase II subunit RPB3





• Molecule 12: RPOL4c domain-containing protein








• Molecule 13: DNA-directed RNA polymerase II subunit E




• Molecule 14: DNA-directed RNA polymerase II subunit F




• Molecule 15: DNA-directed RNA polymerase II subunit RPB7





• Molecule 16: DNA-directed RNA polymerases I, II, and III subunit RPABC3









• Molecule 17: DNA-directed RNA polymerase II subunit RPB9




• Molecule 18: DNA-directed RNA polymerases I, II, and III subunit RPABC5





• Molecule 19: RNA_pol_L_2 domain-containing protein




• Molecule 20: RNA polymerase II subunit K




• Molecule 21: Transcription initiation factor IIB







• Molecule 22: Non-template DNA




• Molecule 23: TATA-box-binding protein





• Molecule 24: General transcription factor IIF subunit 1







• Molecule 25: General transcription factor IIF subunit 2



• Molecule 26: Template DNA



• Molecule 27: Transcription initiation factor IIA subunit 1



PRO ASP
SER SER
LYS LYS
LEU LEU
ILE ILE
GLN GLN
HIS HIS
MET MET
ALA ALA
ASN ASN
THR THR
ALA ALA
SER SER
ASN ASN
MET MET
SER SER
GLY GLY
ALA ALA
VAL VAL
LEU LEU
PRO PRO


SER VAL
VAL VAL
LEU LEU
ALA ALA
GLN GLN
GLN GLN
VAL VAL
ALA ALA
VAL VAL
LEU LEU
PRO PRO
THR THR
ILE ILE
GLN GLN
ALA ALA
LEU LEU
VAL VAL
PRO PRO

ALA ALA
ALA PRO
THR THR
PRO PRO
LEU LEU
GLN GLN
ALA ALA
GLN GLN
VAL VAL
LEU LEU
GLN GLN
ALA ALA
THR THR
ALA ALA
GLN GLN
VAL VAL
LEU LEU
PRO PRO

LYS ASP
ASP GLY
ALA ALA
GLU GLU
ASP ASP
G307
Q308
V309
E310
E311
E312
P313
L314
N315
S316
E317
D318
D319
V320
S321
D322
E323
E324
G325
Q326
E327
L328
F329
D330
T331
E332
N333
V334
V335
V336
C337
Q338
Y339
D340
K341
I342
H343
R344
S345
K346
N347
K348
N349
K350
F351
H352
L353
K354
D355
G356
I357
N358
N359
L360


N361
G362
R363
D364
Y365
Y366
F367
S368
K369
A370
I371
G372
D373
A374
E375
I376

• Molecule 28: Transcription initiation factor IIA subunit 2



M1 A2
Y3 Q4
L5 Y6
R7 N8
T9 T10
L11 M12
M13 S14
L15 L16
Q16 E17
S18 L19
D20 E21
L22 E22
I23 Q24
S25 E26
E27 L28
I29 T29
P30 Q31
L32 R32
A33 V33
K34 I34
Q35 V36
L37 L38
Q39 F40
D41 K42
A43 I44
M45 A46
A47 L48
A49 O50
R51 V52
R53 R54
R55 M56
M57 F58
R59 G60

• Molecule 29: General transcription factor IIE subunit 1



MET ALA
ALA PRO
ASP ASP
VAL VAL
LEU LEU
THR THR
V10 P11
H109 M110
A12 A13
R16 L17
F25 Y26
E30 S31
A33 I36
L37 I38
R39 M40
A41 C42
D47 R56
K57 K70
K73 C74
R75 M76
E79 T80
A81 A82
D83 G84
K85 T86
T87 R88
H89 Y92
F93 I94
M95

Y96 R97
T98 V103
K104 L107
D108 H109
M110 R111
I114 E115
E118 R119
D120 S121
T122 N123
R124 A125
S126 F127
K128 C129
P130 V131
C132 S133
L134 T135
F136 T137
D138 L139
E140 L144
M148 T149
G150 F151
F152 R153
C154 T155
F156 C157
H158 T159
E160 V161
E162 D163
D164 E165


S166 A167
M168 P169
K170 K171
D172 A173
R174 T175
L176 L177
A178 E182
Q183 I184
I187 Y188
R192 E193
T194 E195
D196 VAL
ASN ASN
LEU LEU
ALA ALA
THR THR
GLU GLU
ILE ILE
GLN GLN
VAL VAL
MET MET
MET MET
ASP ASP
ASP ASP
GLN GLN
GLU GLU
ASP ASP
LEU LEU
LEU LEU
PHE PHE
GLU GLU
GLU GLU
VAL VAL
VAL VAL
LYS LYS
HIS HIS
ASP ASP
ARG ARG
VAL VAL
ILE ILE
GLU GLU
ASP ASP
GLU VAL
GLU MET
GLU MET
GLU MET
ASP ASP
ASP ASP
LEU LEU
LEU LEU
PHE PHE
GLU GLU
VAL VAL
VAL VAL
LYS LYS
HIS HIS
ASP ASP
ARG ARG
VAL VAL
ILE ILE
MET MET
VAL VAL
VAL VAL
GLY GLY
SER SER
GLY GLY
ALA ALA
PRO PRO
ALA ALA
PHE PHE
SER SER
TYR TYR
SER SER
GLU GLU
VAL VAL
SER SER
GLN GLN
ARG ARG
PRO PRO
GLU GLU

ALA GLY
HIS HIS
ARG ARG
GLU GLU
ALA ALA
TRP TRP
ASP ASP
ALA ALA
THR THR
ALA ALA
GLY GLY
PRO PRO
SER SER
SER SER
GLU GLU
HIS HIS
THR THR
LEU LEU
THR THR
VAL VAL
ASN ASN
VAL VAL
ILE ILE
GLU GLU
ASP ASP
VAL VAL
MET MET
MET MET
ASP ASP
ASP ASP
GLN GLN
GLU GLU
ASP ASP
LEU LEU
LEU LEU
PHE PHE
GLU GLU
GLU GLU
VAL VAL
VAL VAL
LYS LYS
HIS HIS
ASP ASP
ARG ARG
VAL VAL
ILE ILE
MET MET
VAL VAL
VAL VAL
GLY GLY
SER SER
GLY GLY
ALA ALA
PRO PRO
ALA ALA
PHE PHE
SER SER
TYR TYR
SER SER
GLU GLU
VAL VAL
SER SER
ALA ALA
ASN ASN
GLY GLY
SER SER
GLU GLU
ASP ASP


ASP MET
GLU GLU
THR THR
SER SER
GLU GLU
ASP ASP
MET MET
ASP ASP
ALA ALA
SER SER
PHE PHE
SER SER
PRO PRO
GLU GLU
ARG ARG
GLU GLU
ALA ALA
VAL VAL
VAL VAL
VAL VAL
HIS HIS
PRO PRO
HIS HIS
ASP ASP
VAL VAL
ASN ASN
GLU GLU
ASP ASP
GLU VAL
GLU MET
GLU MET
GLU MET
ASP ASP
ASP ASP
LEU LEU
LEU LEU
PHE PHE
GLU GLU
GLU GLU
VAL VAL
VAL VAL
LYS LYS
HIS HIS
ASP ASP
ARG ARG
VAL VAL
ILE ILE
MET MET
VAL VAL
VAL VAL
GLY GLY
SER SER
GLY GLY
ALA ALA
PRO PRO
ALA ALA
PHE PHE
SER SER
TYR TYR
SER SER
GLU GLU
VAL VAL
SER SER
ALA ALA
ASN ASN
GLY GLY
SER SER
GLU GLU
ASP ASP

LEU VAL
VAL VAL
GLN GLN
MET MET
THR THR
PRO PRO
GLU GLU
ASP ASP
MET MET
LYS LYS
GLU GLU
ALA ALA
THR THR
ILE ILE
ALA ALA
MET MET
GLN GLN
ARG ARG
PHE PHE
ASP ASP
LEU LEU
PHE PHE
GLU GLU


• Molecule 30: Transcription initiation factor IIE subunit beta



• Molecule 31: Unassigned peptide, likely TFIIE-Beta



• Molecule 32: Unassigned peptide, likely XPB



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	26146	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	43	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	31.233	Depositor
Minimum map value	-16.532	Depositor
Average map value	-0.005	Depositor
Map value standard deviation	0.967	Depositor
Recommended contour level	6.4	Depositor
Map size (\AA)	472.49997, 472.49997, 472.49997	wwPDB
Map dimensions	450, 450, 450	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.05, 1.05, 1.05	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP, BEF, ZN, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	0	0.24	0/5875	0.39	0/7955
2	1	0.24	0/2210	0.38	0/2975
3	2	0.24	0/3230	0.37	0/4376
4	3	0.24	0/387	0.36	0/519
5	4	0.25	0/2103	0.43	0/2846
6	5	0.24	0/529	0.36	0/714
7	6	0.24	0/2624	0.40	0/3555
8	7	0.24	0/4994	0.39	0/6745
9	A	0.24	0/11479	0.40	0/15496
10	B	0.25	0/9257	0.41	0/12493
11	C	0.24	0/2102	0.41	0/2857
12	D	0.24	0/1064	0.34	0/1428
13	E	0.25	0/1752	0.40	0/2366
14	F	0.24	0/646	0.39	0/871
15	G	0.25	0/1382	0.41	0/1874
16	H	0.24	0/1207	0.42	0/1628
17	I	0.24	0/949	0.42	0/1284
18	J	0.25	0/516	0.39	0/696
19	K	0.25	0/939	0.38	0/1271
20	L	0.24	0/378	0.41	0/500
21	M	0.24	0/2337	0.39	0/3154
22	N	0.47	0/1537	0.86	0/2374
23	O	0.24	0/1448	0.41	0/1948
24	Q	0.24	0/1167	0.39	0/1576
25	R	0.23	0/1817	0.39	0/2445
26	T	0.50	0/1518	0.87	0/2333
27	U	0.23	0/945	0.39	0/1274
28	V	0.24	0/816	0.39	0/1105
29	W	0.23	0/1686	0.37	0/2266
30	X	0.23	0/1427	0.37	0/1916
All	All	0.26	0/68321	0.43	0/92840

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	5751	0	5794	73	0
2	1	2167	0	2175	27	0
3	2	3158	0	3213	42	0
4	3	382	0	380	4	0
5	4	2066	0	2098	44	0
6	5	523	0	530	5	0
7	6	2567	0	2539	45	0
8	7	4890	0	4949	59	0
9	A	11274	0	11406	129	0
10	B	9076	0	9116	82	0
11	C	2059	0	2007	26	0
12	D	1050	0	1033	4	0
13	E	1721	0	1737	17	0
14	F	636	0	665	5	0
15	G	1351	0	1358	13	0
16	H	1186	0	1147	9	0
17	I	928	0	859	12	0
18	J	507	0	523	5	0
19	K	920	0	942	12	0
20	L	373	0	378	5	0
21	M	2301	0	2316	17	0
22	N	1369	0	743	7	0
23	O	1422	0	1514	14	0
24	Q	1138	0	1103	7	0
25	R	1788	0	1819	16	0
26	T	1358	0	749	8	0
27	U	930	0	888	5	0
28	V	806	0	818	7	0
29	W	1659	0	1666	17	0
30	X	1403	0	1428	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
31	Y	80	0	19	2	0
32	Z	40	0	11	0	0
33	0	8	0	0	0	0
34	4	2	0	0	0	0
34	6	3	0	0	0	0
34	A	2	0	0	0	0
34	B	1	0	0	0	0
34	C	1	0	0	0	0
34	I	2	0	0	0	0
34	J	1	0	0	0	0
34	L	1	0	0	0	0
34	M	1	0	0	0	0
34	W	1	0	0	0	0
35	7	27	0	12	1	0
36	7	1	0	0	0	0
36	A	1	0	0	0	0
37	7	4	0	0	0	0
All	All	66935	0	65935	647	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 647 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:4:255:CYS:SG	5:4:258:HIS:ND1	2.35	0.99
11:C:67:ARG:NH2	18:J:3:ILE:O	1.97	0.97
7:6:109:THR:HG1	7:6:148:SER:HG	1.18	0.89
16:H:8:ASP:OD2	16:H:32:SER:OG	1.95	0.85
9:A:1208:SER:O	9:A:1260:ARG:NH1	2.10	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	0	710/760 (93%)	680 (96%)	30 (4%)	0	100	100
2	1	253/548 (46%)	243 (96%)	10 (4%)	0	100	100
3	2	380/462 (82%)	369 (97%)	10 (3%)	1 (0%)	41	76
4	3	42/309 (14%)	41 (98%)	1 (2%)	0	100	100
5	4	259/308 (84%)	251 (97%)	8 (3%)	0	100	100
6	5	64/71 (90%)	64 (100%)	0	0	100	100
7	6	325/395 (82%)	312 (96%)	13 (4%)	0	100	100
8	7	601/782 (77%)	578 (96%)	23 (4%)	0	100	100
9	A	1413/1970 (72%)	1378 (98%)	35 (2%)	0	100	100
10	B	1130/1174 (96%)	1101 (97%)	29 (3%)	0	100	100
11	C	253/275 (92%)	247 (98%)	6 (2%)	0	100	100
12	D	126/142 (89%)	124 (98%)	2 (2%)	0	100	100
13	E	207/210 (99%)	204 (99%)	3 (1%)	0	100	100
14	F	77/127 (61%)	77 (100%)	0	0	100	100
15	G	169/172 (98%)	168 (99%)	1 (1%)	0	100	100
16	H	146/150 (97%)	141 (97%)	5 (3%)	0	100	100
17	I	112/125 (90%)	106 (95%)	6 (5%)	0	100	100
18	J	62/67 (92%)	61 (98%)	1 (2%)	0	100	100
19	K	113/117 (97%)	111 (98%)	2 (2%)	0	100	100
20	L	42/58 (72%)	39 (93%)	3 (7%)	0	100	100
21	M	294/316 (93%)	291 (99%)	3 (1%)	0	100	100
23	O	177/339 (52%)	176 (99%)	1 (1%)	0	100	100
24	Q	134/517 (26%)	132 (98%)	2 (2%)	0	100	100
25	R	218/249 (88%)	215 (99%)	3 (1%)	0	100	100
27	U	109/376 (29%)	102 (94%)	7 (6%)	0	100	100
28	V	97/109 (89%)	95 (98%)	2 (2%)	0	100	100
29	W	198/439 (45%)	195 (98%)	3 (2%)	0	100	100
30	X	169/291 (58%)	166 (98%)	3 (2%)	0	100	100
All	All	7880/10858 (73%)	7667 (97%)	212 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	2	49	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	0	624/664 (94%)	623 (100%)	1 (0%)	93	96
2	1	241/484 (50%)	241 (100%)	0	100	100
3	2	342/399 (86%)	342 (100%)	0	100	100
4	3	44/283 (16%)	44 (100%)	0	100	100
5	4	234/272 (86%)	234 (100%)	0	100	100
6	5	59/64 (92%)	59 (100%)	0	100	100
7	6	295/352 (84%)	294 (100%)	1 (0%)	92	95
8	7	536/688 (78%)	535 (100%)	1 (0%)	93	96
9	A	1254/1749 (72%)	1253 (100%)	1 (0%)	93	97
10	B	994/1027 (97%)	994 (100%)	0	100	100
11	C	234/252 (93%)	234 (100%)	0	100	100
12	D	118/126 (94%)	117 (99%)	1 (1%)	81	89
13	E	191/192 (100%)	191 (100%)	0	100	100
14	F	69/111 (62%)	69 (100%)	0	100	100
15	G	152/153 (99%)	152 (100%)	0	100	100
16	H	129/131 (98%)	129 (100%)	0	100	100
17	I	103/112 (92%)	103 (100%)	0	100	100
18	J	53/56 (95%)	53 (100%)	0	100	100
19	K	104/106 (98%)	104 (100%)	0	100	100
20	L	41/55 (74%)	41 (100%)	0	100	100
21	M	253/268 (94%)	252 (100%)	1 (0%)	91	94
23	O	154/293 (53%)	154 (100%)	0	100	100
24	Q	121/448 (27%)	119 (98%)	2 (2%)	60	78

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
25	R	196/218 (90%)	195 (100%)	1 (0%)	88	93
27	U	105/324 (32%)	105 (100%)	0	100	100
28	V	90/98 (92%)	89 (99%)	1 (1%)	73	84
29	W	182/373 (49%)	180 (99%)	2 (1%)	73	84
30	X	154/261 (59%)	154 (100%)	0	100	100
All	All	7072/9559 (74%)	7060 (100%)	12 (0%)	93	96

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
24	Q	151	ARG
25	R	230	LYS
29	W	153	ARG
28	V	82	ARG
9	A	483	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 17 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
33	SF4	0	1000	1	0,12,12	0.00	-	-		
35	ADP	7	901	36,37	24,29,29	0.69	0	29,45,45	0.78	1 (3%)
37	BEF	7	903	35	0,3,3	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
33	SF4	0	1000	1	-	-	0/6/5/5
35	ADP	7	901	36,37	-	1/12/32/32	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
35	7	901	ADP	C5-C6-N6	2.24	123.76	120.35

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
35	7	901	ADP	O4'-C4'-C5'-O5'


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
35	7	901	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-12619. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 225




Y Index: 225




Z Index: 225


6.2.2 Raw map



X Index: 225



Y Index: 225




Z Index: 225


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 168




Y Index: 199




Z Index: 213


6.3.2 Raw map



X Index: 168



Y Index: 198



Z Index: 215

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 6.4. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.5.1 emd_12619_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 567 nm^3 ; this corresponds to an approximate mass of 512 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)




*Reported resolution corresponds to spatial frequency of 0.152 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.152 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	6.60	-	-
Author-provided FSC curve	6.48	8.55	6.90
Calculated*	7.89	10.35	8.25

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.89 differs from the reported value 6.6 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-12619 and PDB model 7NW0. Per-residue inclusion information can be found in section [3](#) on page [12](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 6.4 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Atom inclusion [i](#)



At the recommended contour level, 71% of all backbone atoms, 54% of all non-hydrogen atoms, are inside the map.