



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 15, 2022 – 12:08 PM JST

PDB ID : 7WWQ
Title : Crystal structure of human Ufd1-Npl4 complex
Authors : Nguyen, T.Q.; Le, L.T.M.; Kim, D.H.; Ko, K.S.; Lee, H.T.; Nguyen, Y.T.K.;
Kim, H.S.; Han, B.W.; Kang, W.; Yang, J.K.
Deposited on : 2022-02-14
Resolution : 2.72 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.30
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.30


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3359 (2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622 (2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)
RSRZ outliers	127900	3276 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	465	
2	B	16	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3273 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear protein localization protein 4 homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	386	3084	1958	520	594	12	0	0	0

- Molecule 2 is a protein called Ubiquitin recognition factor in ER-associated degradation protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	16	138	93	23	22	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	51	51	51	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nuclear protein localization protein 4 homolog



- Molecule 2: Ubiquitin recognition factor in ER-associated degradation protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	152.96Å 165.42Å 192.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.24 – 2.72 30.24 – 2.72	Depositor EDS
% Data completeness (in resolution range)	99.1 (30.24-2.72) 99.1 (30.24-2.72)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.71 (at 2.72Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.241 , 0.277 0.240 , 0.275	Depositor DCC
R_{free} test set	1703 reflections (5.24%)	wwPDB-VP
Wilson B-factor (Å ²)	51.6	Xtrriage
Anisotropy	0.726	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3273	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/3153	0.65	1/4276 (0.0%)
2	B	0.45	0/141	0.86	0/188
All	All	0.45	0/3294	0.66	1/4464 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	156	GLU	C-N-CD	-5.01	109.57	120.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	421	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3084	0	2964	68	1
2	B	138	0	144	7	0
3	A	51	0	0	5	0
All	All	3273	0	3108	71	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:260:ASN:OD1	2:B:260:ASN:N	2.19	0.75
2:B:258:ILE:HB	2:B:259:PRO:HD3	1.69	0.74
1:A:294:GLU:H	1:A:294:GLU:CD	1.93	0.71
1:A:548:GLN:OE1	3:A:602:HOH:O	2.09	0.69
1:A:403:CYS:HA	2:B:260:ASN:HA	1.76	0.68
1:A:254:GLY:HA2	1:A:305:LEU:HD22	1.79	0.64
1:A:409:LEU:HD13	1:A:442:PRO:HG3	1.80	0.64
1:A:407:PRO:HG3	2:B:271:ILE:HB	1.82	0.61
1:A:329:ARG:HD2	1:A:373:THR:HG21	1.83	0.61
1:A:321:ARG:HD2	3:A:639:HOH:O	2.02	0.60
1:A:473:ILE:HD12	1:A:508:HIS:CD2	2.38	0.59
1:A:545:ARG:HH11	1:A:545:ARG:HG2	1.67	0.59
1:A:321:ARG:NH2	3:A:601:HOH:O	2.07	0.59
1:A:314:ASP:O	1:A:373:THR:HG22	2.04	0.57
1:A:519:MET:HE3	1:A:548:GLN:HG2	1.86	0.57
1:A:405:ASP:CG	2:B:259:PRO:HD2	2.26	0.56
1:A:319:ASP:OD1	1:A:322:LYS:HG2	2.06	0.56
1:A:112:GLN:O	1:A:116:LYS:HE3	2.06	0.56
1:A:411:TYR:CZ	1:A:441:ARG:HG3	2.41	0.55
2:B:258:ILE:HB	2:B:259:PRO:CD	2.37	0.53
1:A:314:ASP:HB3	1:A:373:THR:HG23	1.90	0.53
1:A:257:THR:HG21	1:A:269:GLU:OE2	2.09	0.53
1:A:409:LEU:HB3	1:A:442:PRO:HB2	1.91	0.53
1:A:344:ALA:O	1:A:348:GLN:HG3	2.09	0.52
1:A:292:LYS:NZ	1:A:515:THR:OG1	2.42	0.52
1:A:342:ILE:HA	1:A:395:VAL:HG11	1.92	0.50
1:A:414:GLU:C	1:A:421:VAL:HG22	2.32	0.50
1:A:477:ASP:N	1:A:477:ASP:OD1	2.46	0.49
1:A:382:PHE:O	1:A:454:THR:HG21	2.13	0.48
1:A:390:GLN:O	1:A:394:LEU:HD12	2.13	0.48
1:A:358:SER:O	1:A:360:ASP:N	2.44	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:GLN:O	1:A:169:ARG:NH1	2.45	0.47
1:A:353:ASN:ND2	1:A:363:PHE:O	2.38	0.47
1:A:538:GLU:O	1:A:542:THR:HG23	2.16	0.46
1:A:258:GLU:HG2	3:A:610:HOH:O	2.16	0.46
1:A:223:VAL:O	1:A:452:ILE:HG12	2.16	0.46
1:A:348:GLN:HB3	1:A:365:SER:HB3	1.98	0.46
1:A:299:ILE:HD11	1:A:514:VAL:HG11	1.97	0.46
1:A:160:LYS:N	1:A:160:LYS:HD3	2.30	0.46
1:A:160:LYS:HZ1	1:A:221:ARG:HH12	1.64	0.45
1:A:156:GLU:HA	1:A:157:PRO:O	2.17	0.45
1:A:322:LYS:HB3	1:A:324:THR:HG23	1.97	0.45
1:A:430:ASP:OD2	1:A:430:ASP:N	2.50	0.45
1:A:411:TYR:CE1	1:A:441:ARG:HG3	2.52	0.45
1:A:506:ASP:OD1	1:A:508:HIS:N	2.50	0.45
1:A:187:SER:HA	1:A:396:ARG:HH22	1.83	0.44
1:A:401:LEU:HD22	1:A:413:LYS:HD3	2.00	0.43
1:A:216:ASN:O	1:A:390:GLN:NE2	2.51	0.43
1:A:436:ILE:HD12	1:A:436:ILE:HA	1.80	0.43
1:A:364:GLY:HA2	1:A:392:MET:CE	2.49	0.42
1:A:409:LEU:HB3	1:A:442:PRO:CB	2.49	0.42
1:A:219:LYS:O	1:A:219:LYS:HE3	2.19	0.42
1:A:493:SER:O	1:A:496:THR:HG23	2.20	0.42
1:A:406:ALA:HA	1:A:407:PRO:HD3	1.96	0.42
1:A:488:LEU:HD13	1:A:509:LEU:HD11	2.02	0.42
1:A:519:MET:O	1:A:519:MET:HG3	2.20	0.42
1:A:249:PHE:HB3	1:A:285:LEU:HD21	2.01	0.42
1:A:428:ASP:C	1:A:436:ILE:HG22	2.41	0.41
1:A:292:LYS:O	1:A:295:VAL:HG12	2.20	0.41
1:A:231:HIS:HD2	3:A:649:HOH:O	2.03	0.41
1:A:409:LEU:HD22	1:A:442:PRO:HG2	2.03	0.41
1:A:333:THR:OG1	1:A:334:TYR:N	2.48	0.41
1:A:413:LYS:HD2	1:A:413:LYS:HA	1.92	0.41
1:A:458:LYS:O	1:A:460:PRO:HD3	2.21	0.41
1:A:281:THR:HG22	1:A:283:ASN:H	1.86	0.41
1:A:405:ASP:OD1	1:A:405:ASP:N	2.53	0.41
1:A:281:THR:HG22	1:A:284:SER:N	2.36	0.40
1:A:421:VAL:HB	1:A:422:PRO:HD3	2.03	0.40
1:A:247:GLN:HG3	2:B:270:PHE:CE2	2.56	0.40
1:A:464:PHE:CE2	1:A:507:PHE:HB2	2.56	0.40
1:A:547:GLU:O	1:A:551:THR:HG23	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:430:ASP:OD1	1:A:438:GLN:NE2[2_655]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/465 (81%)	352 (94%)	23 (6%)	1 (0%)	41	65
2	B	14/16 (88%)	11 (79%)	3 (21%)	0	100	100
All	All	390/481 (81%)	363 (93%)	26 (7%)	1 (0%)	41	65

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	359	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/413 (82%)	316 (94%)	22 (6%)	17	37
2	B	15/15 (100%)	12 (80%)	3 (20%)	1	3
All	All	353/428 (82%)	328 (93%)	25 (7%)	14	33

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	ASP
1	A	153	ASN
1	A	186	ILE
1	A	249	PHE
1	A	281	THR
1	A	284	SER
1	A	313	THR
1	A	321	ARG
1	A	325	VAL
1	A	353	ASN
1	A	373	THR
1	A	381	HIS
1	A	387	VAL
1	A	423	ASP
1	A	425	PHE
1	A	432	PHE
1	A	454	THR
1	A	477	ASP
1	A	487	SER
1	A	493	SER
1	A	498	SER
1	A	547	GLU
2	B	260	ASN
2	B	264	LYS
2	B	267	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	438	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/465 (83%)	0.23	28 (7%) 15 13	33, 61, 132, 158	0
2	B	16/16 (100%)	0.40	1 (6%) 20 19	61, 77, 128, 144	0
All	All	402/481 (83%)	0.24	29 (7%) 15 14	33, 63, 132, 158	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	173	GLY	7.5
2	B	262	GLU	4.4
1	A	213	ILE	4.0
1	A	151	TYR	3.7
1	A	172	THR	3.7
1	A	355	CYS	3.7
1	A	158	PRO	3.3
1	A	432	PHE	3.3
1	A	154	HIS	3.3
1	A	170	LYS	3.3
1	A	359	PRO	3.1
1	A	429	VAL	3.1
1	A	155	LEU	2.9
1	A	150	ASP	2.9
1	A	431	LYS	2.6
1	A	358	SER	2.4
1	A	360	ASP	2.3
1	A	148	ASP	2.3
1	A	497	SER	2.2
1	A	362	HIS	2.2
1	A	169	ARG	2.2
1	A	153	ASN	2.2
1	A	361	GLY	2.2
1	A	423	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	387	VAL	2.1
1	A	428	ASP	2.1
1	A	413	LYS	2.1
1	A	187	SER	2.0
1	A	368	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.