

wwPDB X-ray Structure Validation Summary Report (i)

Oct 15, 2023 – 11:04 PM EDT

PDB ID	:	8E6I
Title	:	X-ray structure of the Deinococcus radiodurans Nramp/MntH divalent tran-
		sition metal transporter M230A mutant in an inward-open, manganese-bound
		state
Authors	:	Ray, S.; Gaudet, R.
Deposited on		
Resolution	:	2.52 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	5743 (2.54-2.50)
Clashscore	141614	6463 (2.54-2.50)
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630(2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			9%		
1	А	414	81%	13%	5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	OLC	А	511	-	-	-	Х
4	OLC	А	514	-	-	-	Х
4	OLC	А	515	-	-	-	Х
4	OLC	А	517	-	-	-	Х
4	OLC	А	523	-	-	-	Х
5	PEG	А	528	-	-	-	Х



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 3451 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Divalent metal cation transporter MntH.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	392	Total 2934	C 1935	N 486	0 497	S 16	0	2	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	MET	-	expression tag	UNP Q9RTP8
А	24	HIS	-	expression tag	UNP Q9RTP8
A	25	HIS	-	expression tag	UNP Q9RTP8
А	26	HIS	-	expression tag	UNP Q9RTP8
А	27	HIS	-	expression tag	UNP Q9RTP8
А	28	HIS	-	expression tag	UNP Q9RTP8
А	29	HIS	-	expression tag	UNP Q9RTP8
А	30	HIS	-	expression tag	UNP Q9RTP8
А	31	HIS	-	expression tag	UNP Q9RTP8
А	230	ALA	MET	engineered mutation	UNP Q9RTP8

• Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	2	Total Mn 2 2	0	0

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	2	Total Cl 2 2	0	0

• Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4$).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total C O 22 18 4	0	0
4	А	1	Total C O 22 18 4	0	0
4	А	1	Total C O 25 21 4	0	0
4	А	1	Total C 18 18	0	0
4	А	1	Total C O 18 14 4	0	0
4	А	1	Total C O 25 21 4	0	0
4	А	1	Total C 17 17	0	0
4	А	1	Total C O 25 21 4	0	0
4	А	1	Total C O 25 21 4	0	0
4	А	1	Total C O 25 21 4	0	0
4	А	1	Total C O 25 21 4	0	0
4	А	1	Total C O 16 12 4	0	0
4	А	1	Total C O 25 21 4	0	0
4	А	1	Total C 14 14	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total C 17 17	0	0
4	А	1	Total C O 25 21 4	0	0
4	А	1	Total C O 23 19 4	0	0
4	А	1	Total C O 25 21 4	0	0
4	А	1	Total C 8 8	0	0
4	А	1	Total C 9 9	0	0
4	А	1	Total C 8 8	0	0
4	А	1	Total C 11 11	0	0
4	А	1	Total C O 12 8 4	0	0

Continued from previous page...

• Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	А	1	Total 4	$\begin{array}{c} \mathrm{C} \\ \mathrm{3} \end{array}$	0 1	0	0

• Molecule 6 is water.




]	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	6	А	69	Total O 69 69	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Divalent metal cation transporter MntH



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants	58.67Å 71.35Å 98.59Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.32 - 2.52	Depositor
Resolution (A)	45.32 - 2.52	EDS
% Data completeness	99.9 (45.32-2.52)	Depositor
(in resolution range)	99.9 (45.32 - 2.52)	EDS
R _{merge}	0.27	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.09 (at 2.51 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
D D.	0.225 , 0.266	Depositor
R, R_{free}	0.223 , 0.263	DCC
R_{free} test set	1454 reflections (10.00%)	wwPDB-VP
Wilson B-factor $(Å^2)$	54.3	Xtriage
Anisotropy	0.268	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.31 , 64.7	EDS
L-test for twinning ²	$ \langle L \rangle = 0.50, \langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3451	wwPDB-VP
Average B, all atoms $(Å^2)$	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.42% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OLC, MN, CL, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	I Chain Bond lengths				angles
Mol Chain		RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.24	0/3003	0.43	0/4111

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	added) Clashes Symm-C	
1	А	2934	0	3058	36	0
2	А	2	0	0	0	0
3	А	2	0	0	1	0
4	А	440	0	692	27	0
5	А	4	0	2	0	0
6	А	69	0	0	3	0
All	All	3451	0	3752	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:LEU:HD21	4:A:527:OLC:H2A	1.66	0.78
1:A:253:LYS:HB3	1:A:405:LEU:HD11	1.76	0.67
3:A:504:CL:CL	6:A:609:HOH:O	2.50	0.67
4:A:505:OLC:H14A	4:A:506:OLC:H11A	1.78	0.66
1:A:355:ILE:HG12	4:A:517:OLC:H18B	1.80	0.63

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	А	392/414~(95%)	385~(98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	298/321~(93%)	295~(99%)	3~(1%)	76 89		

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	56	ASP
1	А	82	ASN



Continued from previous page...

Mol	Chain	Res	Type
1	А	255	ARG

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such side chains are listed below:

Mol	Chain	Res	Type
1	А	82	ASN
1	А	237	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 28 ligands modelled in this entry, 4 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	\mathbf{ths}	В	ond ang	les
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	OLC	А	505	-	21,21,24	0.86	2 (9%)	$22,\!22,\!25$	1.01	1 (4%)
4	OLC	А	510	-	24,24,24	0.82	2 (8%)	$25,\!25,\!25$	0.98	1 (4%)
4	OLC	А	523	-	7,7,24	0.32	0	6,6,25	0.62	0
4	OLC	А	517	-	24,24,24	0.81	2 (8%)	$25,\!25,\!25$	0.96	1 (4%)
4	OLC	А	511	-	16,16,24	0.33	0	$15,\!15,\!25$	0.77	0



8E6I

Mol	Tuno	Chain	Res	Link	Bo	ond leng	$_{\rm ths}$	B	ond ang	les
10101	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
4	OLC	А	513	-	24,24,24	0.81	2 (8%)	25,25,25	0.88	1 (4%)
4	OLC	А	507	-	24,24,24	0.82	2 (8%)	25,25,25	0.91	1 (4%)
4	OLC	А	509	-	17,17,24	0.96	2 (11%)	18,18,25	1.12	1 (5%)
5	PEG	А	528	-	3,3,6	0.33	0	2,2,5	0.11	0
4	OLC	А	521	-	22,22,24	0.84	2 (9%)	23,23,25	0.98	1 (4%)
4	OLC	А	526	-	10,10,24	0.41	0	9,9,25	0.92	0
4	OLC	А	512	-	24,24,24	0.81	1 (4%)	$25,\!25,\!25$	1.01	1 (4%)
4	OLC	А	518	-	13,13,24	0.35	0	12,12,25	0.75	0
4	OLC	А	522	-	24,24,24	0.80	2 (8%)	25,25,25	0.98	1 (4%)
4	OLC	А	516	-	15,15,24	0.99	2 (13%)	16,16,25	1.04	1 (6%)
4	OLC	А	508	-	17,17,24	0.32	0	16,16,25	0.81	0
4	OLC	А	524	-	8,8,24	0.30	0	7,7,25	0.76	0
4	OLC	А	506	-	21,21,24	0.86	2(9%)	$22,\!22,\!25$	0.94	1 (4%)
4	OLC	А	515	-	24,24,24	0.80	2 (8%)	25,25,25	0.96	1 (4%)
4	OLC	А	525	-	7,7,24	0.30	0	6,6,25	0.70	0
4	OLC	А	519	-	16,16,24	0.32	0	$15,\!15,\!25$	0.77	0
4	OLC	А	520	-	24,24,24	0.80	2 (8%)	$25,\!25,\!25$	0.95	1 (4%)
4	OLC	А	514	-	24,24,24	0.81	2 (8%)	25,25,25	0.93	1 (4%)
4	OLC	А	527	-	11,11,24	1.15	2 (18%)	12,12,25	1.17	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLC	А	505	-	-	12/21/21/24	-
4	OLC	А	510	-	-	14/24/24/24	-
4	OLC	А	523	-	-	3/5/5/24	-
4	OLC	А	517	-	-	12/24/24/24	-
4	OLC	А	511	-	-	7/14/14/24	-
4	OLC	А	513	-	-	14/24/24/24	-
4	OLC	А	507	-	-	13/24/24/24	-
4	OLC	А	509	-	-	10/17/17/24	-
5	PEG	А	528	-	-	0/0/1/4	-
4	OLC	А	521	-	-	16/22/22/24	-
4	OLC	А	526	_	-	4/8/8/24	_



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLC	А	512	-	-	11/24/24/24	-
4	OLC	А	518	-	-	7/11/11/24	-
4	OLC	А	522	-	-	10/24/24/24	-
4	OLC	А	516	-	-	11/15/15/24	-
4	OLC	А	508	-	-	8/15/15/24	-
4	OLC	А	524	-	-	4/6/6/24	-
4	OLC	А	506	-	-	10/21/21/24	-
4	OLC	А	515	-	-	9/24/24/24	-
4	OLC	А	525	-	-	3/5/5/24	-
4	OLC	А	519	-	-	4/14/14/24	-
4	OLC	А	520	-	-	13/24/24/24	-
4	OLC	А	514	-	-	14/24/24/24	-
4	OLC	А	527	-	-	6/11/11/24	_

Continued from previous page...

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
4	А	512	OLC	O20-C1	2.56	1.40	1.33
4	А	507	OLC	O20-C1	2.48	1.40	1.33
4	А	509	OLC	O20-C1	2.47	1.40	1.33
4	А	505	OLC	O20-C1	2.47	1.40	1.33
4	А	515	OLC	O20-C1	2.44	1.40	1.33

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
4	А	527	OLC	O20-C1-C2	2.86	120.87	111.91
4	А	512	OLC	O20-C1-C2	2.84	120.83	111.91
4	А	509	OLC	O20-C1-C2	2.84	120.81	111.91
4	А	510	OLC	O20-C1-C2	2.73	120.47	111.91
4	А	505	OLC	O20-C1-C2	2.71	120.42	111.91

There are no chirality outliers.

 $5~{\rm of}~215$ torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	507	OLC	C21-C22-C24-O25
4	А	509	OLC	C21-C22-C24-O25



	Chain	-	1 0	Atoms
4	А	509	OLC	O23-C22-C24-O25
4	А	509	OLC	O20-C21-C22-C24
4	А	509	OLC	O20-C21-C22-O23

Continued from previous page...


There are no ring outliers.

16 monomers are involved in 27 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	А	505	OLC	1	0
4	А	510	OLC	3	0
4	А	517	OLC	2	0
4	А	511	OLC	1	0
4	А	513	OLC	2	0
4	А	507	OLC	1	0
4	А	521	OLC	1	0
4	А	526	OLC	1	0
4	А	512	OLC	1	0
4	А	522	OLC	2	0
4	А	524	OLC	2	0
4	А	506	OLC	3	0
4	А	515	OLC	3	0
4	А	519	OLC	3	0
4	А	514	OLC	3	0
4	А	527	OLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$<$ RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	392/414~(94%)	0.66	38 (9%) 7 7	39, 56, 127, 195	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	46	PRO	10.6
1	А	244	ARG	10.4
1	А	48	VAL	10.4
1	А	247	THR	9.9
1	А	49	ILE	9.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	\mathbf{RSR}	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
4	OLC	А	523	8/25	0.38	0.41	81,92,95,96	0
4	OLC	А	511	17/25	0.50	0.45	$63,\!75,\!86,\!89$	0




	-	m previou		1	Daga	DOD		0.00
Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	OLC	А	520	25/25	0.52	0.33	57,69,83,86	0
4	OLC	А	514	25/25	0.56	0.45	57, 72, 76, 84	0
4	OLC	А	515	25/25	0.59	0.45	$67,\!74,\!78,\!84$	0
4	OLC	А	517	25/25	0.60	0.48	$72,\!78,\!89,\!99$	0
4	OLC	А	512	25/25	0.61	0.27	77,81,104,109	0
4	OLC	А	513	25/25	0.64	0.35	65,94,100,101	0
4	OLC	А	506	22/25	0.66	0.39	62,68,73,76	0
4	OLC	А	519	17/25	0.67	0.31	72,80,81,84	0
5	PEG	А	528	4/7	0.67	0.49	67,69,69,70	0
4	OLC	А	508	18/25	0.68	0.37	55,64,69,71	0
4	OLC	А	526	11/25	0.71	0.40	65,69,73,73	0
4	OLC	А	510	25/25	0.71	0.31	49,78,102,106	0
4	OLC	А	524	9/25	0.72	0.32	71,77,85,86	0
4	OLC	А	518	14/25	0.72	0.29	71,82,85,85	0
4	OLC	А	507	25/25	0.72	0.33	59,71,97,99	0
4	OLC	А	522	25/25	0.73	0.29	62,73,83,92	0
4	OLC	А	525	8/25	0.73	0.35	61,63,66,67	0
4	OLC	А	516	16/25	0.76	0.24	78,84,91,98	0
4	OLC	А	505	22/25	0.78	0.33	52,65,77,78	0
4	OLC	А	521	23/25	0.81	0.24	68,76,84,86	0
4	OLC	А	527	12/25	0.83	0.22	62,65,66,68	0
4	OLC	А	509	18/25	0.84	0.21	88,95,102,103	0
2	MN	А	501	1/1	0.86	0.05	78,78,78,78	1
2	MN	А	502	1/1	0.93	0.11	71,71,71,71	1
3	CL	А	503	1/1	0.93	0.15	62,62,62,62	0
3	CL	А	504	1/1	0.98	0.09	64,64,64,64	0


Continued from previous page...

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



























































6.5 Other polymers (i)

There are no such residues in this entry.

