



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 26, 2023 – 12:09 AM EDT

PDB ID : 1A3X
Title : PYRUVATE KINASE FROM SACCHAROMYCES CEREVISIAE COM-
PLEXED WITH PG, MN²⁺ AND K⁺
Authors : Jurica, M.S.; Mesecar, A.; Heath, P.J.; Shi, W.; Nowak, T.; Stoddard, B.L.
Deposited on : 1998-01-26
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.34


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain			
1	A	500	30%	51%	15%	..
1	B	500	32%	49%	16%	..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PGA	A	1005	-	-	X	-
2	PGA	B	1006	-	X	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PYRUVATE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	487	Total 3725	C 2347	N 642	O 718	S 18	0	0	0
1	B	487	Total 3725	C 2347	N 642	O 718	S 18	0	0	0

- Molecule 2 is 2-PHOSPHOGLYCOLIC ACID (three-letter code: PGA) (formula: C₂H₅O₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	A	1	Total 9	C 2	O 6	P 1	0	0
2	B	1	Total 9	C 2	O 6	P 1	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mn 1 1	0	0
3	B	1	Total Mn 1 1	0	0

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total K 1 1	0	0
4	B	1	Total K 1 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: PYRUVATE KINASE

Chain A: 



- Molecule 1: PYRUVATE KINASE

Chain B: 



PHE	L422	M348	E274	E197	V136
LYS	V423	A349	P272	D198	I137
ALA	T424	E350	A273	L199	S138
GLY	R425	Q357	P274	R200	A139
ALA	C426	A358	E275	F201	G140
GLY	P427	I359	V279	H208	R141
HIS	R428	A360	K280	M209	I142
SER	A429	Y361	K281	V210	I143
M493	A430	L362	K282	F211	Y144
T494	R431	P363	L283	A212	V145
L495	S433	G366	I284	S213	D146
Q496	H434	R369	L289	F214	D147
V497	L435	N370	K292	I215	G148
S498	Y436	C371	P293	R216	V149
T499	R437	T372	V294	L150	L150
V500	V438	P373	L295	S151	F152
	P441	K374	C286	D220	Q153
	F444	P375	T298	V221	V154
	E445	T376	Q299	L222	L155
	E446	S377	M300	T223	E156
	K446	T378	L301	I224	V157
	E447	V382	E302	R225	V158
PRO			S303	E226	D159
VAL			M304	V227	D160
SER			T305	L228	K161
ASP			P308	Q231	T162
TRP			R309	G232	L163
			P310	V235	K164
			T311	V239	V165
			E314	K240	K166
			V315	L241	A167
			S316	E242	M169
			D317	N243	A170
			V318	Q244	G171
			V400	Q245	K172
			L401	G246	I173
			S402	V247	C174
			T403	N248	H176
			S404	G249	G178
				L254	V179
				K255	M180
				V256	L181
				T257	P182
				D258	G183
				M330	T184
				L331	D185
				A336	V186
				K337	D187
				G338	L188
				I342	P189
				M343	R264
				V345	G265
				T346	D266
				I421	L267
					G268
					I269
					E270
					K194
					D195
					K196

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	108.30Å 106.40Å 105.50Å 90.00° 110.80° 90.00°	Depositor
Resolution (Å)	100.00 – 3.00	Depositor
% Data completeness (in resolution range)	85.0 (100.00-3.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.227 , 0.341	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7472	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, MN, PGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	2/3781 (0.1%)	0.88	4/5124 (0.1%)
1	B	0.41	0/3781	1.03	7/5124 (0.1%)
All	All	0.51	2/7562 (0.0%)	0.96	11/10248 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	374	LYS	C-N	-26.09	0.84	1.34
1	A	375	PRO	C-N	5.31	1.46	1.34

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	375	PRO	O-C-N	-31.91	71.64	122.70
1	B	375	PRO	CA-C-N	-31.09	48.81	117.20
1	A	375	PRO	O-C-N	-25.38	82.10	122.70
1	A	375	PRO	CA-C-N	-24.09	64.20	117.20
1	B	374	LYS	O-C-N	-21.78	79.71	121.10

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	374	LYS	Peptide
1	A	375	PRO	Mainchain
1	B	374	LYS	Mainchain,Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3725	0	3804	369	1
1	B	3725	0	3804	375	1
2	A	9	0	2	5	0
2	B	9	0	2	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
All	All	7472	0	7612	744	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 49.

The worst 5 of 744 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:374:LYS:HG2	1:A:375:PRO:HD3	1.25	1.11
1:B:186:VAL:HG23	1:B:216:ARG:HE	1.19	1.06
1:A:272:PRO:HB2	1:A:275:GLU:HG3	1.28	1.06
1:B:272:PRO:HB2	1:B:275:GLU:HG3	1.37	1.05
1:A:390:VAL:HB	1:A:395:ALA:HB3	1.37	1.02

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:ARG:NH2	1:B:268:GLY:O[1_545]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	481/500 (96%)	382 (79%)	65 (14%)	34 (7%)	1	5
1	B	481/500 (96%)	377 (78%)	71 (15%)	33 (7%)	1	6
All	All	962/1000 (96%)	759 (79%)	136 (14%)	67 (7%)	1	6

5 of 67 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	10	LEU
1	A	12	VAL
1	A	96	THR
1	A	99	VAL
1	A	106	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	412/423 (97%)	337 (82%)	75 (18%)	1	9
1	B	412/423 (97%)	339 (82%)	73 (18%)	2	9
All	All	824/846 (97%)	676 (82%)	148 (18%)	1	9

5 of 148 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	225	ARG
1	B	445	GLU
1	B	283	LEU
1	B	376	THR
1	A	283	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	244	GLN
1	B	483	GLN
1	B	280	GLN
1	B	370	ASN
1	B	496	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PGA	A	1005	3,1,4	8,8,8	3.75	3 (37%)	10,11,11	2.88	4 (40%)
2	PGA	B	1006	3,4	8,8,8	2.53	3 (37%)	10,11,11	2.90	4 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PGA	A	1005	3,1,4	-	2/6/6/6	-
2	PGA	B	1006	3,4	-	4/6/6/6	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1005	PGA	O1P-C2	-9.17	1.36	1.43
2	B	1006	PGA	O1P-C2	-4.85	1.39	1.43
2	A	1005	PGA	P-O3P	4.09	1.70	1.54
2	B	1006	PGA	P-O3P	3.97	1.70	1.54
2	B	1006	PGA	P-O2P	2.24	1.57	1.50

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1006	PGA	O1P-P-O2P	6.95	125.98	106.47
2	A	1005	PGA	O1P-P-O2P	6.63	125.08	106.47
2	B	1006	PGA	O1P-C2-C1	3.78	116.24	110.54
2	A	1005	PGA	O1P-C2-C1	3.66	116.06	110.54
2	A	1005	PGA	O3P-P-O1P	2.88	114.39	106.73

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1006	PGA	C2-O1P-P-O3P
2	A	1005	PGA	O2-C1-C2-O1P
2	A	1005	PGA	O1-C1-C2-O1P
2	B	1006	PGA	O2-C1-C2-O1P
2	B	1006	PGA	O1-C1-C2-O1P

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1005	PGA	5	0
2	B	1006	PGA	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	374:LYS	C	375:PRO	N	0.84

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.