



wwPDB NMR Structure Validation Summary Report ⓘ

Feb 7, 2022 – 03:49 PM EST

PDB ID : 1AFF
Title : DNA QUADRUPLEX CONTAINING GGGG TETRADS AND (T.A).A TRI-
ADS, NMR, 8 STRUCTURES
Authors : Kettani, A.; Bouaziz, S.; Wang, W.; Jones, R.A.; Patel, D.J.
Deposited on : 1997-03-06

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.26
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26

2 Ensemble composition and analysis

This entry contains 8 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 516 atoms, of which 188 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called QUADRUPLEX DNA (5'-D(TP*AP*GP*G)-3').


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	4	Total 129	C 40	H 47	N 17	O 22	P 3	0
1	B	4	Total 129	C 40	H 47	N 17	O 22	P 3	0
1	C	4	Total 129	C 40	H 47	N 17	O 22	P 3	0
1	D	4	Total 129	C 40	H 47	N 17	O 22	P 3	0

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: QUADRUPLEX DNA (5'-D(TP*AP*GP*G)-3')




- Molecule 1: QUADRUPLEX DNA (5'-D(TP*AP*GP*G)-3')



- Molecule 1: QUADRUPLEX DNA (5'-D(TP*AP*GP*G)-3')



- Molecule 1: QUADRUPLEX DNA (5'-D(TP*AP*GP*G)-3')



4.2 Residue scores for the first model from the NMR ensemble

No representative models were identified. Colouring as in section 4.1 above.

- Molecule 1: QUADRUPLEX DNA (5'-D(TP*AP*GP*G)-3')



T1
A2
G3
C4

- Molecule 1: QUADRUPLEX DNA (5'-D(TP*AP*GP*G)-3')

Chain B:  100%

T1
A2
G3
C4

- Molecule 1: QUADRUPLEX DNA (5'-D(TP*AP*GP*G)-3')

Chain C:  25%  75%

T1
A2
G3
C4

- Molecule 1: QUADRUPLEX DNA (5'-D(TP*AP*GP*G)-3')

Chain D:  100%

T1
A2
G3
C4

5 Refinement protocol and experimental data overview

The models were refined using the following method: *DISTANCE GEOMETRY AND SIMULATED ANNEALING, DISTANCE RESTRAINED MOLECULAR DYNAMICS REFINEMENT AND RELAXATION MATRIX INTENSITY REFINEMENT.*

Of the 100 calculated structures, 8 were deposited, based on the following criterion: *LEAST ENERGY AND LEAST RESTRAINT VIOLATIONS.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.1
X-PLOR	structure solution	

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	2.11±0.22	4±1/92 (3.8± 1.3%)	2.81±0.13	10±2/141 (7.3± 1.7%)
1	B	2.99±1.38	6±1/92 (6.8± 1.4%)	3.37±0.34	14±4/141 (10.1± 2.6%)
1	C	1.99±0.17	2±1/92 (2.7± 0.9%)	2.68±0.10	9±1/141 (6.1± 1.1%)
1	D	2.68±0.70	6±1/92 (6.9± 1.5%)	3.52±0.38	15±3/141 (10.9± 2.1%)
All	All	2.60	149/2944 (5.1%)	3.13	388/4512 (8.6%)

5 of 45 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	1	DT	C5-C7	-57.62	1.15	1.50	4	6
1	D	1	DT	C5-C7	-32.84	1.30	1.50	4	7
1	A	1	DT	C2'-C1'	11.99	1.64	1.52	2	5
1	D	4	DG	C3'-O3'	-10.67	1.30	1.44	4	1
1	B	2	DA	C5'-C4'	10.40	1.62	1.51	4	8

5 of 118 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	D	2	DA	O4'-C1'-N9	23.53	124.47	108.00	6	8
1	B	2	DA	O4'-C1'-N9	18.45	120.91	108.00	2	8
1	B	1	DT	C6-C5-C7	-17.48	112.42	122.90	8	6
1	B	2	DA	O4'-C4'-C3'	-16.87	95.88	106.00	8	8
1	D	2	DA	O4'-C4'-C3'	-16.52	96.09	106.00	6	8

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	82	47	47	4±1
1	B	82	47	47	6±4
1	C	82	47	47	3±1
1	D	82	47	47	7±3
All	All	2624	1504	1504	134

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

5 of 61 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:D:4:DG:H2'	1:D:4:DG:OP2	0.82	1.75	2	2
1:B:3:DG:N3	1:B:3:DG:H5'	0.79	1.91	1	2
1:D:3:DG:H5'	1:D:3:DG:N3	0.78	1.93	1	3
1:C:1:DT:H4'	1:C:2:DA:OP1	0.73	1.84	2	1
1:A:2:DA:H1'	1:A:3:DG:OP2	0.73	1.83	1	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided