

Full wwPDB X-ray Structure Validation Report (i)

May 21, 2020 – 11:59 pm BST

PDB ID : 1CHO

Title: CRYSTAL AND MOLECULAR STRUCTURES OF THE COMPLEX OF

ALPHA-*CHYMOTRYPSIN WITH ITS INHIBITOR TURKEY OVOMU-

COID THIRD DOMAIN AT 1.8 ANGSTROMS RESOLUTION

Authors: Fujinaga, M.; Sielecki, A.R.; Read, R.J.; Ardelt, W.; Laskowskijunior, M.;

James, M.N.G.

Deposited on : 1988-03-04

Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references 1) were used in the production of this report:

MolProbity: 4.02b-467

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# ext{Entries}, ext{resolution range}(ext{Å}))$
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	\mathbf{Q}	uality of chain		
1	Е	13	38%	31%	8%	23%
2	F	131	66%		27%	5% •
3	G	97	75%		15%	7% •
4	I	56	64%		25%	• • 5%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2371 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ALPHA-CHYMOTRYPSIN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	r.	Ε 10	Total	С	Ν	О	S	0	0	0
1			68	45	11	11	1	U		

• Molecule 2 is a protein called ALPHA-CHYMOTRYPSIN A.

\mathbf{Mol}	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	F	131	Total 980	C 618	N 162	O 196	S 4	0	0	0	ĺ

• Molecule 3 is a protein called ALPHA-CHYMOTRYPSIN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	97	Total 702	C 436	N 123	O 136	S 7	0	0	0

• Molecule 4 is a protein called TURKEY OVOMUCOID THIRD DOMAIN (OMTKY3).

Mol	Chain	Residues	Atoms					ZeroOcc	${f AltConf}$	Trace
4	I	53	Total 400	C 246	N 67	O 81	S 6	0	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	E	8	Total O 8 8	0	0
5	F	101	Total O 101 101	0	0
5	G	68	Total O 68 68	0	0
5	I	44	Total O 44 44	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: ALPHA-CHYMOTRYPSIN A





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	44.92Å 54.52Å 57.18Å	Depositor	
a, b, c, α , β , γ	90.00° 103.90° 90.00°	Depositor	
Resolution (Å)	8.00 - 1.80	Depositor	
% Data completeness	(Not available) (8.00-1.80)	Depositor	
(in resolution range)	, , , , ,	Беровног	
R_{merge}	(Not available)	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	PROLSQ	Depositor	
R, R_{free}	0.168 , (Not available)	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	2371	wwPDB-VP	
Average B, all atoms (Å ²)	19.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles			
10101		RMSZ	# Z >5	RMSZ	# Z > 5		
1	E	0.88	0/69	1.55	0/95		
2	F	0.97	0/1000	1.72	17/1361~(1.2%)		
3	G	1.02	0/715	1.69	7/973~(0.7%)		
4	I	0.97	0/408	1.97	9/551 (1.6%)		
All	All	0.99	0/2192	1.75	33/2980 (1.1%)		

There are no bond length outliers.

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
4	I	21	ARG	NE-CZ-NH2	16.07	128.33	120.30
4	I	21	ARG	CD-NE-CZ	12.71	141.39	123.60
4	I	21	ARG	NE-CZ-NH1	-10.74	114.93	120.30
2	F	48	ASN	CA-CB-CG	10.63	136.78	113.40
4	I	27	ASP	CB-CG-OD2	-8.14	110.97	118.30
3	G	230	ARG	NE-CZ-NH1	7.52	124.06	120.30
2	F	35	ASP	CB-CG-OD2	7.01	124.61	118.30
2	F	128	ASP	CA-CB-CG	-6.87	98.28	113.40
2	F	77	SER	C-N-CA	6.80	138.69	121.70
3	G	216	GLY	O-C-N	6.33	132.83	122.70
2	F	137	VAL	O-C-N	6.23	132.67	122.70
4	I	15	ALA	N-CA-CB	-6.22	101.40	110.10
2	F	145	ARG	NE-CZ-NH2	6.17	123.39	120.30
2	F	42	CYS	CA-CB-SG	6.03	124.86	114.00
2	F	145	ARG	CD-NE-CZ	5.92	131.89	123.60
2	F	97	LEU	CB-CA-C	5.84	121.30	110.20
2	F	48	ASN	N-CA-CB	5.80	121.04	110.60
2	F	127	SER	CB-CA-C	-5.70	99.27	110.10
4	I	34	LYS	N-CA-CB	5.63	120.73	110.60
3	G	230	ARG	O-C-N	5.61	131.68	122.70
3	G	213	VAL	CA-CB-CG1	5.61	119.31	110.90
3	G	195	SER	O-C-N	5.60	132.72	123.20
4	I	27	ASP	CB-CG-OD1	5.48	123.23	118.30



Continued	trom	meaningie	naaa
	110116	$D + C \cup U \cup U \cup S$	DUGGERR
0 0 1000100000	J . \circ \circ	r	r

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
3	G	171	TYR	CB-CG-CD1	-5.46	117.72	121.00
2	F	128	ASP	CB-CG-OD2	-5.44	113.40	118.30
3	G	154	ARG	NE-CZ-NH1	5.38	122.99	120.30
4	I	48	LEU	O-C-N	5.33	131.22	122.70
2	F	55	ALA	N-CA-CB	-5.31	102.66	110.10
2	F	76	SER	C-N-CA	5.29	134.93	121.70
2	F	119	SER	N-CA-CB	5.17	118.25	110.50
2	F	145	ARG	NH1-CZ-NH2	-5.16	113.72	119.40
2	F	145	ARG	NE-CZ-NH1	5.09	122.84	120.30
4	I	16	CYS	O-C-N	5.03	130.74	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	68	0	76	5	0
2	F	980	0	951	27	0
3	G	702	0	698	27	0
4	I	400	0	372	10	0
5	Е	8	0	0	0	0
5	F	101	0	0	5	0
5	G	68	0	0	3	0
5	I	44	0	0	3	0
All	All	2371	0	2097	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{c} ext{Interatomic} \ ext{distance } (ext{Å}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap} \ (ext{\AA}) \end{aligned}$
3:G:154:ARG:HH21	3:G:154:ARG:HG3	1.08	1.15
1:E:10:LEU:HD23	3:G:157:GLN:NE2	1.78	0.99



Continued from previous page...

Continuea from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	$overlap (\AA)$
4:I:44:SER:HB2	4:I:47:THR:HG22	1.48	0.93
3:G:149:ALA:HB3	3:G:151:THR:HG23	1.52	0.90
3:G:154:ARG:HG3	3:G:154:ARG:NH2	1.80	0.89
3:G:149:ALA:CB	3:G:151:THR:CG2	2.54	0.86
3:G:149:ALA:HB1	3:G:151:THR:HG22	1.57	0.83
2:F:84:LYS:HD2	5:F:148:HOH:O	1.79	0.83
2:F:145:ARG:HE	2:F:146:TYR:H	1.35	0.75
4:I:44:SER:HB2	4:I:47:THR:CG2	2.17	0.75
3:G:165:ASN:O	3:G:169:LYS:HG3	1.87	0.74
4:I:44:SER:CB	4:I:47:THR:HG22	2.19	0.73
3:G:149:ALA:HB3	3:G:151:THR:CG2	2.14	0.72
3:G:172:TRP:HB2	3:G:176:ILE:HD11	1.76	0.67
2:F:145:ARG:HE	2:F:145:ARG:HA	1.61	0.66
1:E:10:LEU:HD23	3:G:157:GLN:CD	2.20	0.61
2:F:63:SER:OG	5:F:176:HOH:O	2.17	0.58
3:G:149:ALA:CB	3:G:151:THR:HG22	2.20	0.58
3:G:154:ARG:CG	3:G:154:ARG:HH21	2.00	0.56
2:F:16:ILE:O	2:F:144:THR:HA	2.06	0.55
4:I:23:LEU:N	4:I:23:LEU:HD12	2.21	0.55
2:F:143:LEU:HA	3:G:150:ASN:O	2.07	0.55
4:I:10:GLU:HA	5:I:88:HOH:O	2.07	0.55
4:I:44:SER:O	4:I:47:THR:HG22	2.08	0.54
2:F:83:LEU:HD13	2:F:108:LEU:HD13	1.88	0.54
3:G:149:ALA:HB1	3:G:151:THR:CG2	2.23	0.54
4:I:32:GLY:H	4:I:36:ASN:HD22	1.53	0.54
3:G:172:TRP:CB	3:G:176:ILE:HD11	2.38	0.53
3:G:165:ASN:ND2	5:G:254:HOH:O	2.38	0.52
1:E:10:LEU:CD2	3:G:157:GLN:CD	2.78	0.51
2:F:36:LYS:O	2:F:36:LYS:HD3	2.11	0.50
4:I:28:ASN:ND2	5:I:85:HOH:O	2.43	0.50
2:F:128:ASP:CB	5:F:168:HOH:O	2.59	0.50
2:F:145:ARG:NE	2:F:145:ARG:HA	2.25	0.50
2:F:33:LEU:CD2	2:F:66:VAL:HG22	2.42	0.50
2:F:145:ARG:NE	2:F:146:TYR:H	2.07	0.49
2:F:24:PRO:HG3	2:F:71:PHE:CE2	2.47	0.49
3:G:150:ASN:N	3:G:150:ASN:OD1	2.45	0.49
3:G:170:LYS:HA	3:G:170:LYS:HE3	1.95	0.48
3:G:232:THR:HG23	5:G:249:HOH:O	2.13	0.48
2:F:29:TRP:CG	2:F:121:VAL:HB	2.49	0.47
1:E:4:PRO:HB2	1:E:6:ILE:O	2.15	0.47
4:I:34:LYS:HD2	5:I:64:HOH:O	2.16	0.46



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:E:7:GLN:HA	1:E:8:PRO:HD3	1.73	0.46
2:F:136:CYS:O	3:G:159:SER:HA	2.16	0.46
2:F:57:HIS:CE1	3:G:195:SER:HG	2.33	0.46
4:I:53:PHE:N	4:I:53:PHE:CD2	2.82	0.46
2:F:72:ASP:HB2	3:G:154:ARG:HG2	1.99	0.45
2:F:36:LYS:HA	2:F:36:LYS:HD3	1.41	0.45
2:F:99:ILE:HG22	2:F:99:ILE:O	2.19	0.43
3:G:203:LYS:O	3:G:204:ASN:HB2	2.17	0.43
3:G:170:LYS:HD2	5:G:299:HOH:O	2.19	0.43
2:F:128:ASP:CG	5:F:168:HOH:O	2.57	0.42
2:F:87:LYS:HG2	2:F:89:PHE:CE1	2.54	0.42
2:F:53:VAL:HG21	2:F:103:ILE:HD11	2.01	0.42
2:F:17:VAL:O	2:F:18:ASN:HB2	2.19	0.42
2:F:93:LYS:HB3	2:F:101:ASN:ND2	2.35	0.42
2:F:33:LEU:HD13	2:F:60:VAL:HG21	2.02	0.41
2:F:76:SER:O	2:F:77:SER:HB2	2.19	0.41
5:F:149:HOH:O	3:G:197:GLY:HA3	2.20	0.40
2:F:115:SER:HB2	2:F:116:GLN:H	1.64	0.40
3:G:230:ARG:HD3	3:G:230:ARG:HH21	1.66	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	E	8/13 (62%)	8 (100%)	0	0	100	100
2	F	129/131 (98%)	124 (96%)	3 (2%)	2 (2%)	9	2
3	G	95/97 (98%)	93 (98%)	2 (2%)	0	100	100
4	I	$51/56 \ (91\%)$	49 (96%)	2 (4%)	0	100	100
All	All	283/297 (95%)	274 (97%)	7 (2%)	2 (1%)	22	10



All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	77	SER
2	F	99	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	E	8/10~(80%)	7 (88%)	1 (12%)	4 1
2	F	109/109~(100%)	97 (89%)	12 (11%)	6 1
3	G	77/77~(100%)	65 (84%)	12 (16%)	2 0
4	I	$47/48 \; (98\%)$	41 (87%)	6 (13%)	4 1
All	All	241/244~(99%)	210 (87%)	31 (13%)	4 1

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	7	GLN
2	F	36	LYS
2	F	48	ASN
2	F	76	SER
2	F	77	SER
2	F	78	GLU
2	F	84	LYS
2	F	97	LEU
2	F	113	SER
2	F	116	GLN
2	F	125	SER
2	F	128	ASP
2	F	145	ARG
3	G	150	ASN
3	G	151	THR
3	G	154	ARG
3	G	159	SER
3	G	160	LEU



Continued from previous page...

Mol	Chain	Res	Type
3	G	165	ASN
3	G	170	LYS
3	G	195	SER
3	G	203	LYS
3	G	217	SER
3	G	223	SER
3	G	224	THR
4	I	5	SER
4	I	7	ASP
4	I	9	SER
4	I	10	GLU
4	I	21	ARG
4	I	34	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
2	F	18	ASN
2	F	48	ASN
3	G	165	ASN
3	G	204	ASN
4	I	28	ASN
4	I	36	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

