

Full wwPDB X-ray Structure Validation Report (i)

May 15, 2020 - 11:18 am BST

PDB ID	:	1D6J
Title	:	CRYSTAL STRUCTURE OF ADENOSINE 5'-PHOSPHOSULFATE (APS)
		KINASE FROM PENICILLIUM CHRYSOGENUM
Authors	:	MacRae, I.J.; Segel, I.H.; Fisher, A.J.
Deposited on		
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain					
1	А	211	70%	12%	•	16%		
1	В	211	65%	19%	•	14%		



2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 3006 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ADENOSINE-5'PHOSPHOSULFATE KINASE.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
1	1 1	177	Total	С	Ν	Ο	0	0	0
	111	1378	865	249	264	0	0	0	
1	р	189	Total	С	Ν	Ο	0	0	0
ГБ	182	1420	890	257	273	0	0	U	

• Molecule 2 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: $C_4H_6O_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total C O 10 4 6	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	126	Total O 126 126	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	72	Total O 72 72	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: ADENOSINE-5'PHOSPHOSULFATE KINASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	78.86Å 83.48Å 141.95Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.49 - 2.00	Depositor
% Data completeness	98.6 (36.49-2.00)	Depositor
(in resolution range)	30.0 (30.43-2.00)	Depositor
R_{merge}	0.04	Depositor
R _{sym}	(Not available)	Depositor
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.207 , 0.249	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3006	wwPDB-VP
Average B, all atoms $(Å^2)$	45.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	Cham	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.89	0/1404	0.92	3/1908~(0.2%)	
1	В	0.77	1/1445~(0.1%)	0.87	2/1961~(0.1%)	
All	All	0.83	1/2849~(0.0%)	0.90	5/3869~(0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	В	89	GLU	CG-CD	5.50	1.60	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\mathbf{Ideal}(^{o})$
1	В	148	ARG	NE-CZ-NH2	7.22	123.91	120.30
1	В	10	SER	N-CA-C	-7.09	91.84	111.00
1	А	12	LEU	CA-CB-CG	6.94	131.26	115.30
1	А	113	ARG	NE-CZ-NH2	-6.07	117.27	120.30
1	А	142	VAL	N-CA-C	-5.97	94.89	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	А	109	TYR	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1378	0	1344	26	0
1	В	1420	0	1390	49	0
2	А	10	0	4	0	0
3	А	126	0	0	3	0
3	В	72	0	0	2	0
All	All	3006	0	2738	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:8:HIS:HA	1:B:11:ALA:HB2	1.27	1.14
1:B:209:LYS:H	1:B:209:LYS:HD2	1.05	1.13
1:B:209:LYS:CD	1:B:209:LYS:H	1.71	1.01
1:B:47:HIS:CD2	1:B:51:ARG:HH12	1.80	0.98
1:B:209:LYS:N	1:B:209:LYS:HD2	1.76	0.97
1:A:14:ARG:HH12	1:A:129:THR:HG23	1.34	0.93
1:B:47:HIS:HD2	1:B:51:ARG:HH12	0.94	0.93
1:B:170:ALA:N	1:B:171:PRO:HD3	1.88	0.88
1:A:47:HIS:ND1	1:B:11:ALA:HB1	1.90	0.86
1:A:13:THR:OG1	1:A:16:GLU:HG3	1.79	0.82
1:B:141:PRO:HG2	1:B:144:VAL:HG23	1.62	0.81
1:B:8:HIS:CA	1:B:11:ALA:HB2	2.11	0.79
1:B:47:HIS:HD2	1:B:51:ARG:NH1	1.76	0.79
1:A:119:LEU:HD21	3:A:272:HOH:O	1.85	0.75
1:B:141:PRO:HG2	1:B:144:VAL:CG2	2.17	0.75
1:A:14:ARG:HH12	1:A:129:THR:CG2	2.01	0.73
1:B:23:GLN:OE1	1:B:55:VAL:HG22	1.90	0.71
1:B:118:GLN:HE21	1:B:122:VAL:CG2	2.04	0.71

Continued on next page...



Continued from prev		Interatomic	Clash	
Atom-1	Atom-2	distance (\AA)	overlap (Å)	
1:B:170:ALA:N	1:B:171:PRO:CD	2.53	0.70	
1:B:23:GLN:CD	1:B:55:VAL:HG22	2.13	0.69	
1:B:8:HIS:HA	1:B:11:ALA:CB	2.15	0.68	
1:A:109:TYR:OH	1:A:170:ALA:HA	1.93	0.68	
1:B:118:GLN:NE2	1:B:122:VAL:HG21	2.10	0.66	
1:B:187:LEU:HD12	1:B:188:PRO:HD2	1.78	0.66	
1:B:118:GLN:NE2	1:B:122:VAL:CG2	2.61	0.63	
1:A:8:HIS:HD2	1:B:190:GLN:NE2	1.97	0.63	
1:A:8:HIS:HD2	1:B:190:GLN:HE22	1.46	0.63	
1:B:118:GLN:HE21	1:B:122:VAL:HG23	1.63	0.62	
1:B:76:SER:HB3	1:B:79:ASP:HB2	1.82	0.62	
1:A:10:SER:O	1:B:47:HIS:HE1	1.84	0.61	
1:B:147:GLN:O	1:B:148:ARG:HG2	2.01	0.61	
1:A:108:PRO:HG2	1:A:109:TYR:CE1	2.36	0.60	
1:B:187:LEU:HD12	1:B:188:PRO:CD	2.32	0.60	
1:A:8:HIS:HA	1:B:44:GLU:OE2	2.00	0.60	
1:A:66:ARG:HD3	3:A:297:HOH:O	2.00	0.60	
1:B:75:PHE:N	3:B:232:HOH:O	2.34	0.59	
1:A:14:ARG:NH1	1:A:129:THR:CG2	2.66	0.59	
1:B:142:VAL:O	1:B:146:GLU:HG3	2.03	0.58	
1:B:76:SER:HB3	1:B:79:ASP:H	1.68	0.57	
1:B:109:TYR:OH	1:B:170:ALA:HA	2.06	0.56	
1:B:198:ASP:O	1:B:202:THR:HG23	2.06	0.55	
1:A:124:THR:C	1:A:126:GLY:H	2.08	0.55	
1:A:14:ARG:NH1	1:A:129:THR:HG23	2.12	0.54	
1:A:142:VAL:HG23	1:A:142:VAL:O	2.09	0.52	
1:A:87:ILE:HD13	1:A:105:PHE:CG	2.44	0.52	
1:B:66:ARG:HG2	1:B:67:PHE:CD1	2.46	0.51	
1:B:206:LEU:HB3	1:B:207:PRO:HD2	1.92	0.51	
1:A:124:THR:C	1:A:126:GLY:N	2.66	0.49	
1:B:147:GLN:C	1:B:148:ARG:CG	2.81	0.49	
1:B:76:SER:HB3	1:B:79:ASP:CB	2.43	0.49	
1:B:14:ARG:HH22	1:B:124:THR:CB	2.27	0.48	
1:B:51:ARG:HD2	3:B:271:HOH:O	2.13	0.48	
1:A:48:GLN:HG2	1:A:52:ASP:OD2	2.15	0.47	
1:B:143:GLU:H	1:B:143:GLU:CD	2.18	0.46	
1:B:44:GLU:CG	1:B:193:VAL:HG21	2.45	0.46	
1:A:13:THR:HG23	1:A:16:GLU:OE1	2.15	0.46	
1:B:17:ARG:HD2	1:B:98:ASN:OD1	2.16	0.45	
1:B:44:GLU:HG3	1:B:193:VAL:HG21	1.99	0.44	
1:B:146:GLU:C	1:B:148:ARG:H	2.21	0.44	

Continued from previous page...

Continued on next page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:GLU:HG2	1:A:189:VAL:HG12	1.99	0.44
1:B:70:ASN:OD1	1:B:86:ARG:HD2	2.18	0.43
1:A:122:VAL:HG12	1:A:122:VAL:O	2.19	0.43
1:B:185:TYR:O	1:B:186:GLU:C	2.58	0.42
1:B:66:ARG:HG2	1:B:67:PHE:CE1	2.54	0.42
1:B:147:GLN:O	1:B:148:ARG:CG	2.67	0.41
1:A:198:ASP:O	1:A:202:THR:HG23	2.20	0.41
1:A:20:LEU:HD22	1:B:51:ARG:HA	2.02	0.40
1:A:124:THR:HG21	3:A:325:HOH:O	2.20	0.40
1:A:124:THR:O	1:A:126:GLY:N	2.54	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles		
1	А	173/211 (82%)	164~(95%)	8 (5%)	1 (1%)	25 19		
1	В	178/211 (84%)	164 (92%)	13~(7%)	1 (1%)	25 19		
All	All	351/422~(83%)	328~(93%)	21~(6%)	2(1%)	25 19		

All (2) Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	127	GLU
1	В	146	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	144/180~(80%)	140 (97%)	4 (3%)	43 44		
1	В	148/180~(82%)	143~(97%)	5(3%)	37 36		
All	All	292/360~(81%)	283~(97%)	9(3%)	40 40		

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	26	LEU
1	А	111	LYS
1	А	113	ARG
1	А	122	VAL
1	В	10	SER
1	В	26	LEU
1	В	47	HIS
1	В	113	ARG
1	B	209	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	8	HIS
1	А	22	ASN
1	А	70	ASN
1	В	22	ASN
1	В	47	HIS
1	В	118	GLN
1	В	177	ASN
1	В	190	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Tune	Chain	Dog	Tink	B	ond leng	gths	B	ond ang	gles
		туре		nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	2	TLA	А	212	-	3, 9, 9	1.64	1 (33%)	$6,\!12,\!12$	1.88	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TLA	А	212	-	-	0/4/12/12	-

All (1) bond length outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
2	А	212	TLA	O3-C3	2.85	1.48	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	А	212	TLA	C4-C3-C2	-3.26	106.09	113.11
2	А	212	TLA	C1-C2-C3	-2.71	107.28	113.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

