



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 06:20 am BST

PDB ID : 1DOE  
Title : THE MOBIL FLAVIN OF 4-OH BENZOATE HYDROXYLASE: MOTION OF A PROSTHETIC GROUP REGULATES CATALYSIS  
Authors : Gatti, D.L.; Palfey, B.A.; Lah, M.S.; Entsch, B.; Massey, V.; Ballou, D.P.; Ludwig, M.L.  
Deposited on : 1994-09-06  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.


Mol	Chain	Length	Quality of chain
1	A	394	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BR	A	398	-	-	X	-



- Molecule 4 is 2,4-DIHYDROXYBENZOIC ACID (three-letter code: DOB) (formula: C<sub>7</sub>H<sub>6</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 11 7 4	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	168	Total O 168 168	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: P-HYDROXYBENZOATE HYDROXYLASE



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.90Å 146.44Å 88.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (15.00-2.30)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.164 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3382	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DOB, FAD, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/3190	0.76	1/4316 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	LEU	CA-CB-CG	5.66	128.33	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3125	0	3120	31	0
2	A	2	0	0	2	0
3	A	76	0	34	2	0
4	A	11	0	4	0	0
5	A	168	0	0	1	0
All	All	3382	0	3158	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:ARG:HG3	1:A:179:ARG:HH11	1.49	0.78
1:A:126:GLU:HB3	1:A:147:ARG:HH12	1.53	0.72
1:A:293:PRO:HA	2:A:398:BR:BR	2.51	0.65
1:A:166:ARG:HD2	1:A:273:VAL:HG11	1.81	0.62
1:A:45:ALA:HB2	3:A:395[B]:FAD:C10	2.30	0.62
1:A:142:GLU:HG3	1:A:147:ARG:HH11	1.67	0.58
1:A:183:PHE:HE1	1:A:268:LEU:HD22	1.70	0.56
2:A:398:BR:BR	3:A:395[B]:FAD:C10	3.11	0.54
1:A:64:ARG:HE	1:A:67:ARG:HH22	1.57	0.52
1:A:362:ARG:HG2	1:A:362:ARG:HH11	1.75	0.51
1:A:179:ARG:CG	1:A:179:ARG:HH11	2.21	0.51
1:A:143:ARG:HH11	1:A:143:ARG:HG3	1.76	0.50
1:A:327:ARG:HH11	1:A:327:ARG:HG2	1.77	0.49
1:A:307:SER:O	1:A:311:ARG:HG2	2.12	0.49
1:A:189:LEU:HD12	1:A:219:SER:O	2.13	0.48
1:A:243:LEU:HD22	1:A:247:LEU:HD11	1.97	0.47
1:A:56:LEU:HD12	1:A:56:LEU:HA	1.74	0.47
1:A:182:PRO:HA	1:A:227:LEU:HD21	1.96	0.47
1:A:183:PHE:CE1	1:A:268:LEU:HD22	2.50	0.46
1:A:42:ARG:HD2	1:A:44:ARG:HH21	1.82	0.44
1:A:179:ARG:NH1	1:A:179:ARG:CG	2.80	0.44
1:A:273:VAL:HG22	1:A:288:ALA:O	2.17	0.44
1:A:142:GLU:HG3	1:A:147:ARG:NH1	2.32	0.43
1:A:269:ARG:HG2	1:A:270:SER:N	2.33	0.43
1:A:250:GLU:O	1:A:254:LYS:HE2	2.20	0.42
1:A:307:SER:O	1:A:311:ARG:CG	2.67	0.42
1:A:208:PHE:HB3	1:A:224:GLN:HB2	2.01	0.41
1:A:391:GLU:HB2	5:A:530:HOH:O	2.20	0.41
1:A:153:ASP:O	1:A:280:ARG:HD3	2.20	0.41
1:A:29:VAL:HG22	1:A:119:THR:HB	2.03	0.40
1:A:132:LEU:HA	1:A:132:LEU:HD23	1.88	0.40
1:A:393:ILE:HG22	1:A:394:GLU:N	2.37	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/394 (100%)	382 (97%)	9 (2%)	1 (0%)	41	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	393	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/325 (100%)	301 (93%)	24 (7%)	13	17

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	LEU
1	A	52	MET
1	A	56	LEU
1	A	64	ARG
1	A	70	LEU
1	A	91	LEU
1	A	126	GLU
1	A	131	ASP
1	A	173	ARG
1	A	175	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	191	ASP
1	A	220	ARG
1	A	225	VAL
1	A	227	LEU
1	A	237	GLU
1	A	243	LEU
1	A	255	LEU
1	A	269	ARG
1	A	311	ARG
1	A	314	LEU
1	A	352	ARG
1	A	355	ASP
1	A	372	LEU
1	A	388	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FAD	A	395[A]	-	51,58,58	1.52	9 (17%)	60,89,89	3.05	17 (28%)
3	FAD	A	395[B]	-	51,58,58	1.54	9 (17%)	60,89,89	2.90	14 (23%)
4	DOB	A	396	-	9,11,11	2.93	1 (11%)	11,15,15	1.15	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	395[A]	-	-	3/30/50/50	0/6/6/6
3	FAD	A	395[B]	-	-	2/30/50/50	0/6/6/6
4	DOB	A	396	-	-	0/0/4/4	0/1/1/1

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	396	DOB	C1-C1'	-8.41	1.39	1.47
3	A	395[A]	FAD	C2B-C1B	-4.90	1.46	1.53
3	A	395[B]	FAD	C2B-C1B	-4.90	1.46	1.53
3	A	395[B]	FAD	C4-N3	3.84	1.39	1.33
3	A	395[A]	FAD	C5'-C4'	3.35	1.56	1.51
3	A	395[B]	FAD	C5'-C4'	3.35	1.56	1.51
3	A	395[A]	FAD	C2-N3	-2.94	1.32	1.38
3	A	395[A]	FAD	C2A-N3A	2.72	1.36	1.32
3	A	395[B]	FAD	C2A-N3A	2.72	1.36	1.32
3	A	395[B]	FAD	C10-N1	2.59	1.36	1.33
3	A	395[A]	FAD	C1'-N10	-2.50	1.45	1.48
3	A	395[A]	FAD	C2-N1	-2.47	1.33	1.38
3	A	395[A]	FAD	O2B-C2B	2.30	1.48	1.43
3	A	395[B]	FAD	O2B-C2B	2.30	1.48	1.43
3	A	395[B]	FAD	C9A-N10	2.23	1.41	1.38
3	A	395[A]	FAD	C9A-N10	2.19	1.41	1.38
3	A	395[B]	FAD	C4-C4X	2.15	1.45	1.41
3	A	395[A]	FAD	P-O2P	-2.05	1.45	1.55
3	A	395[B]	FAD	P-O2P	-2.05	1.45	1.55

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	395[A]	FAD	C4-N3-C2	15.69	128.39	115.14
3	A	395[B]	FAD	C4-N3-C2	14.46	127.35	115.14
3	A	395[A]	FAD	C10-C4X-N5	9.60	127.90	121.26
3	A	395[B]	FAD	C10-C4X-N5	9.02	127.50	121.26
3	A	395[A]	FAD	C4X-C4-N3	-7.21	113.56	123.43
3	A	395[B]	FAD	C4X-C4-N3	-6.60	114.41	123.43
3	A	395[A]	FAD	C6-C5X-N5	-4.87	113.68	119.05
3	A	395[B]	FAD	C4-C4X-C10	-4.78	116.78	119.95
3	A	395[B]	FAD	C4X-N5-C5X	-4.42	112.36	116.77
3	A	395[B]	FAD	C6-C5X-N5	-4.12	114.51	119.05
3	A	395[A]	FAD	C4-C4X-C10	-4.10	117.24	119.95
3	A	395[B]	FAD	C4X-C10-N10	-3.94	116.25	120.30
3	A	395[A]	FAD	C4X-N5-C5X	-3.75	113.02	116.77
3	A	395[B]	FAD	C9A-C5X-N5	3.46	127.77	122.36
3	A	395[A]	FAD	C4X-C10-N10	-3.38	116.83	120.30
3	A	395[A]	FAD	C4-C4X-N5	-3.32	114.80	118.60
3	A	395[B]	FAD	C9A-N10-C10	-3.19	117.73	121.91
3	A	395[A]	FAD	C4'-C3'-C2'	3.19	120.00	113.36
3	A	395[A]	FAD	C9A-C5X-N5	3.06	127.15	122.36
3	A	395[A]	FAD	C5X-C9A-N10	-2.78	115.70	117.72
3	A	395[A]	FAD	C2B-C3B-C4B	2.69	107.86	102.64
3	A	395[B]	FAD	C2B-C3B-C4B	2.69	107.86	102.64
3	A	395[A]	FAD	O3B-C3B-C4B	-2.61	103.49	111.05
3	A	395[B]	FAD	O3B-C3B-C4B	-2.61	103.49	111.05
3	A	395[B]	FAD	C4-C4X-N5	-2.52	115.71	118.60
3	A	395[A]	FAD	N6A-C6A-N1A	2.43	123.62	118.57
3	A	395[B]	FAD	N6A-C6A-N1A	2.43	123.62	118.57
4	A	396	DOB	C6-C5-C4	-2.35	117.29	119.88
3	A	395[A]	FAD	C1'-N10-C10	2.19	120.37	118.41
3	A	395[A]	FAD	C1'-N10-C9A	2.17	120.00	118.29
3	A	395[A]	FAD	O3'-C3'-C4'	2.17	114.05	108.81
3	A	395[B]	FAD	O4'-C4'-C3'	2.04	114.06	109.10

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	395[A]	FAD	N10-C1'-C2'-O2'
3	A	395[A]	FAD	N10-C1'-C2'-C3'
3	A	395[B]	FAD	O3'-C3'-C4'-C5'
3	A	395[A]	FAD	O4B-C4B-C5B-O5B

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	A	395[B]	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	395[B]	FAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.