



wwPDB NMR Structure Validation Summary Report ⓘ

Feb 10, 2022 – 09:26 AM EST

PDB ID : 1EKZ
Title : NMR STRUCTURE OF THE COMPLEX BETWEEN THE THIRD DSRBD
FROM DROSOPHILA STAUFEN AND A RNA HAIRPIN
Authors : Ramos, A.; Grunert, S.; Bycroft, M.; St Johnston, D.; Varani, G.
Deposited on : 2000-03-11

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.26
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428
RNA backbone	4643	676

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	30	
2	A	76	

2 Ensemble composition and analysis i

This entry contains 36 models. Model 27 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:9-A:30, A:40-A:55, A:62-A:76 (53)	0.60	27

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 13 single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 7, 8, 10, 11, 13, 17, 20, 23, 24, 25, 27, 28, 30, 32, 33
2	1, 5, 22
3	6, 21
Single-model clusters	9; 12; 14; 15; 16; 18; 19; 26; 29; 31; 34; 35; 36

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2173 atoms, of which 949 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called STAUFEN DOUBLE-STRANDED RNA BINDING DOMAIN.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	B	30	961	284	326	112	210	29	0

- Molecule 2 is a protein called MATERNAL EFFECT PROTEIN (STAUFEN).

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	A	76	1212	368	623	107	109	5	0

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble


These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: STAUFEN DOUBLE-STRANDED RNA BINDING DOMAIN

Chain B: 



- Molecule 2: MATERNAL EFFECT PROTEIN (STAUFEN)

Chain A: 



4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 27. Colouring as in section 4.1 above.

- Molecule 1: STAUFEN DOUBLE-STRANDED RNA BINDING DOMAIN

Chain B: 



- Molecule 2: MATERNAL EFFECT PROTEIN (STAUFEN)

Chain A: 



M69
L70
V71
E72
L73
L76

5 Refinement protocol and experimental data overview

Of the 50 calculated structures, 36 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy.*

The authors did not provide any information on software used for structure solution, optimization or refinement.

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	B	1.10±0.01	0±0/708 (0.0± 0.0%)	1.89±0.01	38±0/1102 (3.5± 0.0%)
2	A	1.13±0.01	0±0/420 (0.0± 0.0%)	0.84±0.01	0±0/558 (0.0± 0.0%)
All	All	1.11	0/40608 (0.0%)	1.62	1369/59760 (2.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	B	0.0±0.0	0.1±0.3
2	A	0.0±0.0	3.0±0.0
All	All	0	110

There are no bond-length outliers.

5 of 42 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	17	G	N7-C8-N9	9.47	117.83	113.10	5	36
1	B	1	G	N7-C8-N9	9.45	117.82	113.10	36	36
1	B	9	G	N7-C8-N9	9.37	117.79	113.10	32	36
1	B	18	G	N7-C8-N9	9.31	117.75	113.10	7	36
1	B	20	G	N7-C8-N9	9.29	117.75	113.10	32	36

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	A	20	ARG	Sidechain	36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
2	A	30	ARG	Sidechain	36
2	A	64	ARG	Sidechain	36
1	B	9	G	Sidechain	1
1	B	22	C	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	635	326	326	10±6
2	A	416	453	453	24±11
All	All	37836	28044	28044	1193

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

5 of 625 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:A:41:ILE:HG23	2:A:65:ALA:HB2	1.01	1.26	31	1
2:A:13:VAL:HG22	2:A:70:LEU:HD21	0.96	1.36	25	20
2:A:50:VAL:HG11	2:A:69:MET:CE	0.95	1.92	36	1
2:A:50:VAL:HG11	2:A:69:MET:HE1	0.94	1.35	36	1
2:A:44:CYS:SG	2:A:65:ALA:HB1	0.93	2.02	2	5

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	52/76 (68%)	42±3 (81±5%)	7±2 (14±4%)	3±2 (6±3%)	3	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1872/2736 (68%)	1507 (81%)	261 (14%)	104 (6%)	3 22

5 of 25 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	A	21	ASN	14
2	A	47	GLY	11
2	A	9	PRO	11
2	A	54	GLU	7
2	A	48	SER	6

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	46/65 (71%)	35±3 (77±6%)	11±3 (23±6%)	3 28
All	All	1656/2340 (71%)	1270 (77%)	386 (23%)	3 28

5 of 41 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	A	23	THR	27
2	A	11	SER	26
2	A	75	LYS	20
2	A	19	LYS	20
2	A	20	ARG	18

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	B	29/30 (97%)	13±3 (46±10%)	1±1 (4±5%)	0.13±0.03
All	All	1052/1080 (97%)	479 (46%)	45 (4%)	0.12

The overall RNA backbone suiteness is 0.13.

5 of 29 unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	B	18	G	30
1	B	17	G	28
1	B	27	G	26
1	B	30	C	23
1	B	2	G	21

5 of 14 unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	B	19	G	11
1	B	1	G	8
1	B	3	A	5
1	B	12	C	3
1	B	22	C	3

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided