

Full wwPDB X-ray Structure Validation Report (i)

Jan 29, 2024 – 05:17 PM EST

PDB ID : 1EOB

Title : CRYSTAL STRUCTURE OF ACINETOBACTER SP. ADP1 PROTOCAT-

ECHUATE 3,4-DIOXYGENASE IN COMPLEX WITH 3,4-DIHYDROXYB

ENZOATE

Authors: Vetting, M.W.; D'Argenio, D.A.; Ornston, L.N.; Ohlendorf, D.H.

Deposited on : 2000-03-22

Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	209	77%	18%	
2	В	241	80%	18%	



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 3732 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called PROTOCATECHUATE 3,4-DIOXYGENASE ALPHA CHAIN.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
1	A	202	Total 1602	C 1010	N 285	O 307	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	60	GLY	SER	SEE REMARK 999	UNP P20371

• Molecule 2 is a protein called PROTOCATECHUATE 3,4-DIOXYGENASE BETA CHAIN.


\mathbf{N}	[ol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
	2	В	238	Total 1926	C 1224	N 345	O 352	S 5	0	0	0

• Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total Fe 1 1	0	0

• Molecule 4 is 3,4-DIHYDROXYBENZOIC ACID (three-letter code: DHB) (formula: $C_7H_6O_4$).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	В	1	Total C 11 7	O 4	0	0

• Molecule 5 is water.


\mathbf{Mol}	Chain	Residues	Atoms	$\mathbf{ZeroOcc}$	AltConf
5	A	78	Total O 78 78	0	0
5	В	114	Total O 114 114	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTOCATECHUATE 3,4-DIOXYGENASE ALPHA CHAIN







4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants	144.40Å 144.40Å 144.40Å	Donogiton
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 - 2.20	Depositor
Resolution (A)	21.29 - 2.20	EDS
% Data completeness	86.8 (20.00-2.20)	Depositor
(in resolution range)	86.9 (21.29-2.20)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.44 (at 2.19Å)	Xtriage
Refinement program	CNS	Depositor
D D.	0.182 , 0.222	Depositor
R, R_{free}	0.176 , 0.214	DCC
R_{free} test set	1091 reflections (4.32%)	wwPDB-VP
Wilson B-factor (Å ²)	11.8	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 37.8	EDS
L-test for twinning ²	$< L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.046 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3732	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, DHB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.32	0/1638	0.60	0/2223	
2	В	0.33	0/1981	0.65	2/2691 (0.1%)	
All	All	0.33	0/3619	0.63	2/4914 (0.0%)	

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	451	ASN	N-CA-C	-6.71	92.90	111.00
2	В	452	ARG	N-CA-C	-5.83	95.26	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1602	0	1548	38	0
2	В	1926	0	1858	38	0
3	В	1	0	0	0	0
4	В	11	0	3	1	0
5	A	78	0	0	1	0
5	В	114	0	0	1	1
All	All	3732	0	3409	70	1



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 10.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
1:A:38:ASN:HD21	1. 4. 40. 7/41 . 11/200	distance (Å)	overlap (Å)
1:A:38:A5N:HD21 1:A:28:ILE:HG22	1:A:40:VAL:HG22 1:A:30:VAL:HG23	1.44	0.82
		1.66	0.78
1:A:68:ILE:HD12	1:A:130:ILE:HD13	1.68	0.73
2:B:369:LYS:H	2:B:422:ASN:HD22	1.40	0.68
1:A:53:GLY:HA2	1:A:184:ARG:HE	1.59	0.67
2:B:403:ASN:C	2:B:403:ASN:HD22	2.00	0.64
2:B:315:ASP:HB3	2:B:500:GLN:NE2	2.13	0.64
1:A:38:ASN:ND2	1:A:40:VAL:HG22	2.13	0.64
1:A:68:ILE:HD12	1:A:130:ILE:CD1	2.28	0.63
2:B:403:ASN:ND2	2:B:405:SER:H	1.97	0.62
2:B:315:ASP:HB3	2:B:500:GLN:HE21	1.66	0.61
1:A:39:LEU:HD11	1:A:93:GLY:HA3	1.81	0.61
1:A:177:ARG:O	1:A:178:ASP:HB3	2.00	0.61
1:A:5:LEU:N	1:A:5:LEU:HD12	2.16	0.60
1:A:37:ASN:HB3	1:A:95:THR:HB	1.83	0.60
2:B:369:LYS:H	2:B:422:ASN:ND2	2.02	0.58
2:B:415:TYR:CE1	2:B:448:PRO:HG3	2.39	0.57
1:A:20:GLY:HA3	2:B:400:TRP:HB3	1.85	0.57
2:B:467:ALA:HB3	2:B:472:GLN:NE2	2.20	0.57
1:A:33:HIS:NE2	2:B:355:LYS:HD3	2.21	0.55
1:A:177:ARG:HD3	1:A:182:VAL:HG21	1.88	0.55
2:B:507:LEU:N	2:B:507:LEU:HD22	2.23	0.54
1:A:95:THR:HG22	1:A:106:PHE:CZ	2.43	0.54
1:A:40:VAL:HG12	1:A:109:ILE:HD13	1.89	0.53
2:B:463:PHE:CZ	2:B:476:SER:HB3	2.43	0.53
1:A:30:VAL:HG22	2:B:367:TYR:HD1	1.74	0.53
1:A:38:ASN:HD21	1:A:40:VAL:CG2	2.17	0.51
1:A:128:LEU:HD23	1:A:129:ILE:N	2.25	0.51
2:B:428:ARG:C	2:B:429:MET:HG3	2.32	0.50
2:B:406:GLY:O	2:B:447:TYR:HD1	1.95	0.49
2:B:403:ASN:ND2	2:B:407:ARG:H	2.10	0.49
1:A:20:GLY:CA	2:B:400:TRP:HB3	2.42	0.49
2:B:428:ARG:O	2:B:429:MET:HG3	2.12	0.49
1:A:51:LEU:O	1:A:105:SER:HA	2.12	0.49
2:B:486:ILE:HG23	2:B:492:LEU:CD2	2.43	0.49
1:A:33:HIS:CD2	2:B:355:LYS:HD3	2.48	0.49
1:A:118:LYS:HE3	5:A:867:HOH:O	2.13	0.48

Continued on next page...



 $Continued\ from\ previous\ page...$

Constitued from presso		Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
2:B:400:TRP:HA	2:B:425:GLY:O	2.14	0.48
2:B:462:HIS:CE1	4:B:999:DHB:O3	2.67	0.47
2:B:403:ASN:HD21	2:B:405:SER:HB2	1.79	0.47
2:B:468:ASP:H	2:B:472:GLN:HE22	1.61	0.47
1:A:35:LEU:N	1:A:35:LEU:CD1	2.78	0.46
1:A:55:VAL:O	1:A:62:PRO:HA	2.15	0.46
2:B:469:GLY:H	2:B:472:GLN:NE2	2.13	0.46
1:A:108:THR:OG1	1:A:109:ILE:N	2.49	0.46
2:B:373:PRO:HG3	2:B:422:ASN:O	2.16	0.46
2:B:353:ALA:HA	2:B:356:PHE:CD2	2.51	0.45
1:A:88(B):VAL:HG22	1:A:88(C):ASP:N	2.32	0.45
2:B:458:PRO:HG2	2:B:479:TYR:CD2	2.51	0.45
1:A:131:PHE:O	1:A:132:ALA:HB2	2.16	0.44
1:A:5:LEU:N	1:A:5:LEU:CD1	2.80	0.44
1:A:28:ILE:HG22	1:A:30:VAL:CG2	2.43	0.44
2:B:513:PHE:CE2	2:B:519:ARG:HG2	2.54	0.43
2:B:403:ASN:C	2:B:403:ASN:ND2	2.67	0.43
2:B:464:SER:C	2:B:465:LEU:HD12	2.38	0.43
2:B:510:LYS:HA	2:B:513:PHE:CE1	2.54	0.43
1:A:50:ARG:HB3	1:A:182:VAL:HG22	2.00	0.43
2:B:364:ILE:CD1	2:B:440:ARG:HG2	2.50	0.42
1:A:58:GLY:HA2	1:A:190:GLN:HB3	2.02	0.42
2:B:353:ALA:HA	2:B:356:PHE:HD2	1.84	0.42
1:A:129:ILE:HD12	1:A:140:HIS:CE1	2.55	0.42
1:A:35:LEU:N	1:A:35:LEU:HD12	2.35	0.41
1:A:155:ASP:O	1:A:159:ASN:ND2	2.53	0.41
1:A:39:LEU:HD12	1:A:39:LEU:N	2.35	0.41
1:A:30:VAL:HG22	2:B:367:TYR:CD1	2.53	0.41
2:B:356:PHE:HE2	2:B:430:LEU:HD11	1.86	0.41
1:A:89:PHE:CE2	1:A:91:GLY:HA2	2.55	0.41
1:A:50:ARG:HH11	1:A:50:ARG:HG3	1.86	0.40
2:B:359:LYS:O	2:B:366:ASN:HB3	2.22	0.40
2:B:507:LEU:HD21	5:B:769:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
5:B:892:HOH:O	5:B:892:HOH:O[2_555]	2.12	0.08



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percenti	les
1	A	200/209~(96%)	190 (95%)	9 (4%)	1 (0%)	29 31	-
2	В	$236/241 \ (98\%)$	224 (95%)	12 (5%)	0	100 10	00
All	All	436/450 (97%)	414 (95%)	21 (5%)	1 (0%)	47 55	5

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	178	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Chain Analysed Rotameric Outliers		Percentiles		
1	A	169/175 (97%)	166 (98%)	3 (2%)	59 72	
2	В	201/204 (98%)	197 (98%)	4 (2%)	55 69	
All	All	370/379 (98%)	363 (98%)	7 (2%)	57 71	

All (7) residues with a non-rotameric sidechain are listed below:

\mathbf{Mol}	Chain	Res	\mathbf{Type}
1	A	38	ASN
1	A 109		ILE
1	A	177	ARG
2	В	333	LYS
2	В	381	HIS

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
2	В	403	ASN
2	В	429	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	25	GLN
1	A	38	ASN
1	A	47	GLN
1	A	72	GLN
1	A	159	ASN
2	В	403	ASN
2	В	412	ASN
2	В	422	ASN
2	В	472	GLN
2	В	499	GLN
2	В	512	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	ond leng	$ ag{ths}$	В	ond ang	eles
IVIOI		Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DHB	В	999	3	11,11,11	0.99	0	15,15,15	0.81	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DHB	В	999	3	-	0/4/4/4	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
4	В	999	DHB	O2-C-C1	2.14	120.40	114.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	999	DHB	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$egin{array}{c c} Analysed & <& RSRZ> & \#RSRZ>2 \end{array}$		$OWAB(Å^2)$	Q < 0.9
1	A	202/209 (96%)	-0.19	6 (2%) 50 48	4, 15, 31, 57	0
2	В	238/241 (98%)	-0.53	2 (0%) 86 85	4, 10, 28, 35	0
All	All	440/450 (97%)	-0.37	8 (1%) 68 66	4, 12, 31, 57	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	178	ASP	5.6
1	A	177	ARG	3.9
1	A	179	GLY	3.8
1	A	42	ASP	3.1
2	В	370	ASP	3.0
2	В	303	ILE	2.4
1	A	192	GLU	2.3
1	A	43	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	DHB	В	999	11/11	0.94	0.14	22,23,24,24	0
3	FE	В	600	1/1	0.99	0.04	11,11,11,11	0

6.5 Other polymers (i)

There are no such residues in this entry.

