



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 01:04 am BST

PDB ID : 1F74  
Title : CRYSTAL STRUCTURE ANALYSIS OF N-ACETYLNEURAMINATE  
LYASE FROM HAEMOPHILUS INFLUENZAE: CRYSTAL FORM II COM-  
PLEXED WITH 4-DEOXY-SIALIC ACID  
Authors : Barbosa, J.A.R.G.; Smith, B.J.; DeGori, R.; Lawrence, M.C.  
Deposited on : 2000-06-26  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	 5% 89% 11%
1	C	293	 4% 88% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	934	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called N-ACETYL-NEURAMINATE LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	293	2339	1511	376	440	12	0	12	0
1	C	293	2327	1500	375	439	13	0	11	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	SER	ASN	SEE REMARK 999	UNP P44539
A	229	LYS	ALA	SEE REMARK 999	UNP P44539
A	278	ALA	GLU	SEE REMARK 999	UNP P44539
A	281	VAL	LEU	SEE REMARK 999	UNP P44539
C	131	SER	ASN	SEE REMARK 999	UNP P44539
C	229	LYS	ALA	SEE REMARK 999	UNP P44539
C	278	ALA	GLU	SEE REMARK 999	UNP P44539
C	281	VAL	LEU	SEE REMARK 999	UNP P44539

- Molecule 2 is 6,7,8,9-TETRAHYDROXY-5-METHYLCARBOXAMIDO-2-OXONONANOIC ACID (three-letter code: NAY) (formula: C<sub>11</sub>H<sub>19</sub>NO<sub>8</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			20	11	1	8		
2	C	1	Total	C	N	O	0	0
			20	11	1	8		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	450	Total	O	0	0
			450	450		
5	C	395	Total	O	0	0
			395	395		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-ACETYL-NEURAMINATE LYASE



- Molecule 1: N-ACETYL-NEURAMINATE LYASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.21Å 118.01Å 80.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.60 24.83 – 1.53	Depositor EDS
% Data completeness (in resolution range)	94.9 (15.00-1.60) 85.4 (24.83-1.53)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.17 (at 1.53Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.169 , 0.202 0.166 , 0.193	Depositor DCC
$R_{free}$ test set	4076 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.0	Xtrriage
Anisotropy	0.178	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 59.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5576	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAY, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/2437	0.92	8/3280 (0.2%)
1	C	0.41	0/2424	1.02	16/3260 (0.5%)
All	All	0.40	0/4861	0.97	24/6540 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	C	0	1
All	All	0	3

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	155	TYR	N-CA-CB	9.48	127.67	110.60
1	A	271	PRO	N-CA-CB	7.99	112.89	103.30
1	A	270	GLU	CA-C-O	-7.81	103.69	120.10
1	C	270[A]	GLU	CA-C-O	-7.79	103.74	120.10
1	C	270[B]	GLU	CA-C-O	-7.79	103.74	120.10
1	A	271	PRO	CA-N-CD	-7.72	100.69	111.50
1	C	269	ARG	NE-CZ-NH2	-7.42	116.59	120.30
1	C	271	PRO	N-CA-CB	7.21	111.95	103.30
1	C	102[A]	CYS	CA-CB-SG	6.66	126.00	114.00
1	C	102[B]	CYS	CA-CB-SG	6.66	126.00	114.00
1	C	271	PRO	CA-N-CD	-6.57	102.30	111.50
1	C	2	ARG	NE-CZ-NH2	-6.31	117.14	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	271	PRO	N-CD-CG	6.17	112.45	103.20
1	A	121	TYR	CA-CB-CG	6.11	125.01	113.40
1	C	110	TYR	N-CA-CB	6.11	121.60	110.60
1	C	64	ARG	NE-CZ-NH1	5.83	123.22	120.30
1	C	121	TYR	CA-CB-CG	5.69	124.21	113.40
1	C	2	ARG	N-CA-CB	-5.57	100.57	110.60
1	A	269	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	C	110	TYR	CA-C-O	-5.38	108.81	120.10
1	A	155	TYR	CB-CG-CD1	5.28	124.17	121.00
1	A	31	ARG	NE-CZ-NH1	5.12	122.86	120.30
1	C	64	ARG	CD-NE-CZ	5.09	130.72	123.60
1	C	110	TYR	CA-C-N	5.05	128.32	117.20

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	270	GLU	Mainchain,Peptide
1	C	155	TYR	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2339	0	2370	27	0
1	C	2327	0	2362	31	0
2	A	20	0	18	1	0
2	C	20	0	18	1	0
3	A	12	0	16	2	0
3	C	12	0	16	1	0
4	C	1	0	0	0	0
5	A	450	0	0	4	0
5	C	395	0	0	3	0
All	All	5576	0	4800	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1[A]:MET:CE	1:C:132:ASN:HD22	1.54	1.20
1:C:136:TYR:CE2	1:C:164:LYS:NZ	2.19	1.11
1:C:1[A]:MET:HE3	1:C:132:ASN:HD22	0.91	1.08
1:C:1[A]:MET:HE3	1:C:132:ASN:ND2	1.71	1.06
1:C:136:TYR:HE2	1:C:164:LYS:NZ	1.64	0.93
1:C:177[A]:LYS:HE2	1:C:200:GLY:O	1.71	0.90
1:A:144:VAL:HG11	1:A:175:ARG:HG3	1.64	0.80
1:C:1[A]:MET:CE	1:C:132:ASN:ND2	2.36	0.75
1:A:229:LYS:H	3:A:934:GOL:H12	1.53	0.72
1:C:142:THR:H	1:C:146:MET:HE2	1.56	0.70
1:A:136:TYR:CE2	1:A:164:LYS:NZ	2.60	0.69
1:C:142:THR:H	1:C:146:MET:CE	2.07	0.67
1:C:144:VAL:HG11	1:C:175:ARG:HG3	1.75	0.67
1:C:248:ASN:HD21	1:C:280:GLN:HA	1.60	0.66
1:C:164:LYS:HD2	2:C:951:NAY:O2	1.99	0.62
1:C:136:TYR:HE2	1:C:164:LYS:HZ1	1.45	0.62
1:C:173[A]:LEU:HD12	1:C:185:ILE:HD13	1.82	0.60
1:C:176:LEU:HD12	1:C:180[A]:TYR:HD2	1.67	0.59
1:A:229:LYS:H	3:A:934:GOL:C1	2.17	0.55
1:C:164:LYS:HG3	1:C:205:ILE:HD12	1.89	0.54
1:A:283[B]:LYS:HE2	1:A:287:LEU:HG	1.90	0.54
1:A:248:ASN:HD21	1:A:280:GLN:HA	1.72	0.54
1:C:173[A]:LEU:HD11	1:C:185:ILE:HG21	1.90	0.54
1:C:177[A]:LYS:HE2	1:C:200:GLY:C	2.29	0.53
1:C:177[A]:LYS:CE	1:C:200:GLY:O	2.50	0.53
1:C:177[A]:LYS:HE3	1:C:200:GLY:HA3	1.91	0.53
1:A:136:TYR:CE2	1:A:164:LYS:HE2	2.44	0.52
1:C:1[A]:MET:HE1	1:C:132:ASN:HB2	1.91	0.52
1:A:164:LYS:HD2	2:A:950:NAY:O2	2.08	0.52
1:C:173[A]:LEU:CD1	1:C:185:ILE:HD13	2.40	0.51
1:C:136:TYR:CE2	1:C:164:LYS:CE	2.92	0.51
1:A:106[A]:VAL:HG22	5:A:1297:HOH:O	2.11	0.50
1:A:142:THR:H	1:A:146:MET:CE	2.25	0.49
1:A:224:LYS:HD3	5:A:1137:HOH:O	2.13	0.48
1:A:142:THR:H	1:A:146:MET:HE2	1.77	0.48
1:A:144:VAL:CG1	1:A:175:ARG:HG3	2.42	0.48
1:A:145:ASN:HB2	5:C:1365:HOH:O	2.13	0.48
1:C:177[A]:LYS:HD2	1:C:199:LEU:O	2.14	0.47
1:A:136:TYR:CE2	1:A:164:LYS:CE	2.97	0.47
1:A:94:TYR:CE2	1:A:98[B]:LEU:HD21	2.50	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:173[A]:LEU:CD1	1:C:185:ILE:HG21	2.46	0.46
1:A:41:GLY:HA2	1:A:74:ILE:HB	1.97	0.45
1:A:224:LYS:HD2	1:A:224:LYS:HA	1.87	0.44
1:C:194:LEU:HD13	1:C:223:THR:OG1	2.16	0.44
1:A:53:MET:SD	1:A:271:PRO:HD2	2.56	0.44
1:A:94:TYR:O	1:A:98[B]:LEU:HD13	2.18	0.44
1:A:229:LYS:HG2	5:A:1371:HOH:O	2.18	0.44
1:A:177:LYS:HD3	1:A:177:LYS:HA	1.91	0.43
1:C:218:GLN:NE2	3:C:932:GOL:H12	2.34	0.43
1:C:176:LEU:HD12	1:C:180[A]:TYR:CD2	2.52	0.43
1:A:106[A]:VAL:HG23	5:A:1305:HOH:O	2.17	0.42
1:C:138:ILE:HD12	5:C:1344:HOH:O	2.19	0.42
1:A:173:LEU:HD11	1:A:185:ILE:HG21	2.03	0.41
1:A:220:PHE:O	1:A:224:LYS:HG2	2.21	0.41
1:A:228:LEU:HD11	1:C:228:LEU:HD21	2.02	0.41
1:C:145:ASN:HB3	5:C:1338:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	302/293 (103%)	297 (98%)	5 (2%)	0	100	100
1	C	301/293 (103%)	295 (98%)	6 (2%)	0	100	100
All	All	603/586 (103%)	592 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/242 (105%)	252 (99%)	2 (1%)	81	70
1	C	253/242 (104%)	250 (99%)	3 (1%)	71	54
All	All	507/484 (105%)	502 (99%)	5 (1%)	76	61

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	GLN
1	A	189	PHE
1	C	2	ARG
1	C	79	GLN
1	C	189	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	248	ASN
1	C	132	ASN
1	C	248	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	C	931	-	5,5,5	0.34	0	5,5,5	0.20	0
3	GOL	C	932	-	5,5,5	0.44	0	5,5,5	0.40	0
2	NAY	A	950	-	16,19,19	1.43	4 (25%)	18,25,25	1.89	5 (27%)
3	GOL	A	933	-	5,5,5	0.37	0	5,5,5	0.43	0
3	GOL	A	934	-	5,5,5	0.28	0	5,5,5	0.39	0
2	NAY	C	951	-	16,19,19	1.47	5 (31%)	18,25,25	1.44	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	931	-	-	4/4/4/4	-
3	GOL	C	932	-	-	4/4/4/4	-
2	NAY	A	950	-	-	5/23/27/27	-
3	GOL	A	933	-	-	2/4/4/4	-
3	GOL	A	934	-	-	4/4/4/4	-
2	NAY	C	951	-	-	5/23/27/27	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	951	NAY	C3-C2	2.35	1.54	1.51
2	A	950	NAY	C6-C5	2.33	1.57	1.53
2	C	951	NAY	C6-C5	2.33	1.57	1.53
2	A	950	NAY	C3-C2	2.22	1.54	1.51
2	C	951	NAY	C5-N	2.21	1.50	1.46
2	C	951	NAY	C7-C6	2.07	1.57	1.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	951	NAY	C4-C5	2.07	1.57	1.52
2	A	950	NAY	C4-C3	2.04	1.59	1.52
2	A	950	NAY	C4-C5	2.03	1.57	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	950	NAY	O2-C2-C3	4.49	128.04	120.38
2	A	950	NAY	C3-C4-C5	-3.42	108.71	114.04
2	A	950	NAY	O10-C10-N	3.32	128.05	121.95
2	C	951	NAY	C11-C10-N	-3.03	110.97	116.10
2	C	951	NAY	O2-C2-C3	2.96	125.43	120.38
2	A	950	NAY	C11-C10-N	-2.46	111.94	116.10
2	A	950	NAY	C3-C2-C1	-2.35	115.47	121.32

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	931	GOL	O1-C1-C2-C3
3	C	931	GOL	C1-C2-C3-O3
3	C	932	GOL	O1-C1-C2-C3
3	C	932	GOL	C1-C2-C3-O3
2	A	950	NAY	C3-C4-C5-N
2	A	950	NAY	C3-C4-C5-C6
3	A	933	GOL	O1-C1-C2-C3
3	A	934	GOL	C1-C2-C3-O3
2	C	951	NAY	C3-C4-C5-N
2	C	951	NAY	C3-C4-C5-C6
2	A	950	NAY	C2-C3-C4-C5
2	C	951	NAY	C2-C3-C4-C5
2	A	950	NAY	C1-C2-C3-C4
2	A	950	NAY	O2-C2-C3-C4
3	C	931	GOL	O1-C1-C2-O2
3	C	931	GOL	O2-C2-C3-O3
3	C	932	GOL	O2-C2-C3-O3
3	A	933	GOL	O1-C1-C2-O2
3	A	934	GOL	O2-C2-C3-O3
2	C	951	NAY	O2-C2-C3-C4
2	C	951	NAY	C1-C2-C3-C4
3	C	932	GOL	O1-C1-C2-O2
3	A	934	GOL	O1-C1-C2-O2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	934	GOL	O1-C1-C2-C3


There are no ring outliers.


4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	932	GOL	1	0
2	A	950	NAY	1	0
3	A	934	GOL	2	0
2	C	951	NAY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	293/293 (100%)	-0.04	14 (4%) 30 28	7, 11, 22, 39	0
1	C	293/293 (100%)	-0.11	12 (4%) 37 34	7, 11, 21, 32	0
All	All	586/586 (100%)	-0.08	26 (4%) 34 31	7, 11, 22, 39	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	140	PHE	11.0
1	A	142	THR	7.7
1	A	168	GLY	6.3
1	C	168	GLY	6.2
1	A	1[A]	MET	6.2
1	C	140	PHE	6.2
1	C	142	THR	5.3
1	A	141	LEU	5.2
1	A	144	VAL	3.9
1	C	167	ALA	3.8
1	A	143	GLY	3.5
1	C	145	ASN	3.0
1	A	145	ASN	2.9
1	C	139	PRO	2.7
1	A	167	ALA	2.6
1	A	149	GLU	2.6
1	C	1[A]	MET	2.6
1	C	155	TYR	2.5
1	A	139	PRO	2.4
1	C	143	GLY	2.4
1	C	141	LEU	2.3
1	A	182	ASN	2.2
1	A	138[A]	ILE	2.2
1	C	293	SER	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	112	LYS	2.0
1	A	229	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	934	6/6	0.35	0.41	40,41,41,42	0
3	GOL	C	932	6/6	0.58	0.28	33,36,37,38	0
3	GOL	C	931	6/6	0.68	0.20	49,49,50,50	0
2	NAY	C	951	20/20	0.81	0.15	14,19,26,27	0
3	GOL	A	933	6/6	0.83	0.15	21,23,24,26	0
2	NAY	A	950	20/20	0.85	0.14	12,19,23,25	0
4	CL	C	999	1/1	1.00	0.04	11,11,11,11	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers

There are no such residues in this entry.