



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 25, 2023 – 12:41 AM EDT

PDB ID : 1FIR
Title : CRYSTAL STRUCTURE OF HIV-1 REVERSE TRANSCRIPTION PRIMER TRNA(LYS3)
Authors : Benas, P.; Dumas, P.
Deposited on : 2000-08-06
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.34
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.34


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)
RNA backbone	3102	1117 (3.70-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	76	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1706 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called HIV-1 REVERSE TRANSCRIPTION PRIMER TRNA(LYS3).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	76	1688	758	294	556	78	2	0	2	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	16	Total	O	0	0
			16	16		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HIV-1 REVERSE TRANSCRIPTION PRIMER TRNA(LYS3)



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 1 2	Depositor
Cell constants a, b, c, α , β , γ	100.72Å 100.72Å 82.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 3.30 20.01 – 3.30	Depositor EDS
% Data completeness (in resolution range)	80.5 (40.00-3.30) 87.7 (20.01-3.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.89 (at 3.29Å)	Xtrriage
Refinement program	CNS 0.4	Depositor
R, R_{free}	0.181 , 0.227 0.187 , 0.229	Depositor DCC
R_{free} test set	588 reflections (9.07%)	wwPDB-VP
Wilson B-factor (Å ²)	64.8	Xtrriage
Anisotropy	0.105	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.09 , 11.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.050 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1706	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 2MG, 12A, PSU, MG, 1MA, 70U, H2U, 7MG, 5MC, NA, 2MU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	2/1485 (0.1%)	0.93	4/2308 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	G	OP3-P	-6.73	1.53	1.61
1	A	56	C	N1-C2	-5.01	1.35	1.40

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	60	U	C2'-C3'-O3'	9.95	131.39	109.50
1	A	17	C	N1-C1'-C2'	7.38	123.59	114.00
1	A	9	A	N9-C1'-C2'	6.23	122.11	114.00
1	A	60	U	C4'-C3'-C2'	5.84	108.44	102.60

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	60	U	C3'

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	42	G	Sidechain
1	A	5	G	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1688	0	871	70	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	16	0	0	0	0
All	All	1706	0	871	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:G:C2'	1:A:60:U:H5'	1.84	1.06
1:A:59:G:H2'	1:A:60:U:H5'	1.46	0.98
1:A:6:2MG:HM22	1:A:68:C:H1'	1.52	0.91
1:A:59:G:O2'	1:A:60:U:H5'	1.77	0.85
1:A:50:A:H2'	1:A:51:G:H8	1.48	0.77
1:A:66:U:H2'	1:A:67:U:H6	1.50	0.74
1:A:29:A:H2'	1:A:30:G:H8	1.52	0.74
1:A:65:G:O2'	1:A:66:U:H5'	1.88	0.74
1:A:19[A]:G:H4'	1:A:20[A]:H2U:OP1	1.88	0.73
1:A:20[B]:H2U:H5''	1:A:21:A:OP2	1.89	0.72
1:A:37:12A:OO	1:A:37:12A:C	2.39	0.71
1:A:20[A]:H2U:H5''	1:A:21:A:OP2	1.90	0.71
1:A:50:A:H2'	1:A:51:G:C8	2.25	0.71
1:A:10:2MG:HN2	1:A:26:A:H1'	1.54	0.70
1:A:19[B]:G:H4'	1:A:20[B]:H2U:OP1	1.91	0.69
1:A:29:A:H2'	1:A:30:G:C8	2.30	0.67
1:A:8:U:O4'	1:A:48:5MC:O2'	2.17	0.63
1:A:66:U:C2	1:A:67:U:C6	2.88	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:2MG:HN2	1:A:26:A:C1'	2.13	0.61
1:A:28:C:N3	1:A:29:A:N7	2.48	0.61
1:A:67:U:H2'	1:A:68:C:C6	2.35	0.60
1:A:10:2MG:HM23	1:A:11:C:C2	2.36	0.60
1:A:43:A:H2'	1:A:44:G:C8	2.37	0.59
1:A:66:U:H2'	1:A:67:U:C6	2.36	0.59
1:A:9:A:O2'	1:A:10:2MG:N7	2.37	0.57
1:A:10:2MG:H2'	1:A:11:C:H6	1.69	0.57
1:A:10:2MG:N2	1:A:26:A:H1'	2.18	0.57
1:A:66:U:O2'	1:A:67:U:H5'	2.05	0.57
1:A:59:G:H2'	1:A:60:U:C5'	2.29	0.55
1:A:28:C:C2	1:A:29:A:C8	2.95	0.55
1:A:36:U:H2'	1:A:37:12A:O5'	2.08	0.54
1:A:66:U:H2'	1:A:67:U:O4'	2.07	0.53
1:A:49:5MC:H2'	1:A:49:5MC:O2	2.09	0.53
1:A:37:12A:H5''	1:A:37:12A:H8	1.91	0.53
1:A:10:2MG:O5'	1:A:10:2MG:H8	1.92	0.53
1:A:67:U:H2'	1:A:68:C:H6	1.74	0.53
1:A:44:G:H2'	1:A:44:G:N3	2.24	0.52
1:A:16:H2U:H5''	1:A:16:H2U:H62	1.92	0.52
1:A:23:A:H2'	1:A:24:G:C8	2.45	0.51
1:A:51:G:H2'	1:A:52:G:H8	1.75	0.51
1:A:10:2MG:H2'	1:A:11:C:C6	2.46	0.51
1:A:4:C:H2'	1:A:5:G:H8	1.77	0.50
1:A:7:A:C5	1:A:49:5MC:HM52	2.47	0.49
1:A:37:12A:OO	1:A:37:12A:N1	2.38	0.49
1:A:66:U:C4	1:A:67:U:C5	3.01	0.49
1:A:28:C:C2	1:A:29:A:N7	2.82	0.48
1:A:50:A:O2'	1:A:51:G:H5'	2.14	0.48
1:A:64:U:O2'	1:A:65:G:H5'	2.14	0.47
1:A:5:G:C2	1:A:6:2MG:C8	3.03	0.47
1:A:65:G:H2'	1:A:66:U:H6	1.81	0.46
1:A:62:C:H2'	1:A:63:C:C6	2.50	0.46
1:A:32:C:H2'	1:A:33:U:H5'	1.98	0.46
1:A:72:C:H2'	1:A:73:G:O4'	2.16	0.45
1:A:5:G:H2'	1:A:6:2MG:H8	1.82	0.45
1:A:51:G:H2'	1:A:52:G:C8	2.51	0.45
1:A:17:C:H4'	1:A:18:G:OP2	2.07	0.45
1:A:34:70U:S2	1:A:34:70U:C2'	3.03	0.45
1:A:51:G:H2'	1:A:52:G:O4'	2.16	0.44
1:A:28:C:H42	1:A:29:A:N6	2.15	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34:70U:S2	1:A:34:70U:H2'	2.58	0.44
1:A:54:2MU:H6	1:A:54:2MU:O5'	2.19	0.43
1:A:71:G:H2'	1:A:72:C:O4'	2.18	0.43
1:A:36:U:C2'	1:A:37:12A:O5'	2.66	0.43
1:A:54:2MU:H6'3	1:A:54:2MU:H1'	1.51	0.43
1:A:28:C:N4	1:A:29:A:H62	2.16	0.42
1:A:68:C:C2	1:A:69:G:C8	3.07	0.42
1:A:27:PSU:C4	1:A:28:C:C5	3.08	0.42
1:A:29:A:C4	1:A:30:G:C8	3.08	0.42
1:A:14:A:H2'	1:A:15:G:O4'	2.21	0.41
1:A:19[B]:G:C8	1:A:57:A:C2	3.09	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	70/76 (92%)	11 (15%)	4 (5%)

All (11) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	11	C
1	A	16	H2U
1	A	17	C
1	A	18	G
1	A	22	G
1	A	46	7MG
1	A	47	H2U

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	48	5MC
1	A	49	5MC
1	A	61	C
1	A	75	C

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	16	H2U
1	A	17	C
1	A	46	7MG
1	A	60	U

5.4 Non-standard residues in protein, DNA, RNA chains

16 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	5MC	A	48	1	18,22,23	0.51	0	26,32,35	0.91	1 (3%)
1	H2U	A	47	1	18,21,22	0.75	1 (5%)	21,30,33	0.84	1 (4%)
1	PSU	A	55	1	18,21,22	1.89	1 (5%)	22,30,33	1.01	2 (9%)
1	PSU	A	27	1	18,21,22	1.81	1 (5%)	22,30,33	1.03	1 (4%)
1	H2U	A	20[B]	1	18,21,22	0.95	1 (5%)	21,30,33	0.90	1 (4%)
1	12A	A	37	1	29,36,37	9.18	4 (13%)	34,52,55	1.30	3 (8%)
1	H2U	A	20[A]	1	18,21,22	0.74	1 (5%)	21,30,33	0.97	2 (9%)
1	PSU	A	39	1	18,21,22	1.89	1 (5%)	22,30,33	0.90	1 (4%)
1	7MG	A	46	1	22,26,27	3.70	3 (13%)	29,39,42	1.57	2 (6%)
1	5MC	A	49	1	18,22,23	0.60	0	26,32,35	0.71	1 (3%)
1	70U	A	34	1	22,26,27	5.82	2 (9%)	28,37,40	1.26	2 (7%)
1	2MG	A	10	1	18,26,27	1.21	3 (16%)	16,38,41	0.86	1 (6%)
1	1MA	A	58	1	16,25,26	1.41	4 (25%)	18,37,40	1.15	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	2MG	A	6	1	18,26,27	1.18	2 (11%)	16,38,41	0.96	1 (6%)
1	H2U	A	16	1	18,21,22	0.76	1 (5%)	21,30,33	1.44	4 (19%)
1	2MU	A	54	1	20,23,24	0.56	0	28,33,36	1.24	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5MC	A	48	1	-	1/7/25/26	0/2/2/2
1	H2U	A	47	1	-	3/7/38/39	0/2/2/2
1	PSU	A	55	1	-	1/7/25/26	0/2/2/2
1	PSU	A	27	1	-	0/7/25/26	0/2/2/2
1	H2U	A	20[B]	1	-	2/7/38/39	0/2/2/2
1	12A	A	37	1	-	9/21/43/44	0/3/3/3
1	H2U	A	20[A]	1	-	2/7/38/39	0/2/2/2
1	PSU	A	39	1	-	0/7/25/26	0/2/2/2
1	7MG	A	46	1	-	0/7/37/38	0/3/3/3
1	5MC	A	49	1	-	2/7/25/26	0/2/2/2
1	70U	A	34	1	-	3/13/31/32	0/2/2/2
1	2MG	A	10	1	-	2/5/27/28	0/3/3/3
1	1MA	A	58	1	-	0/3/25/26	0/3/3/3
1	2MG	A	6	1	-	2/5/27/28	0/3/3/3
1	H2U	A	16	1	-	1/7/38/39	0/2/2/2
1	2MU	A	54	1	-	1/9/27/28	0/2/2/2

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	37	12A	C2-S2	-49.04	1.34	1.75
1	A	34	70U	C2-S2	-27.03	1.24	1.67
1	A	46	7MG	C8-N9	-16.77	1.36	1.46
1	A	39	PSU	C6-N1	-7.64	1.23	1.36
1	A	55	PSU	C6-N1	-7.58	1.23	1.36
1	A	27	PSU	C6-N1	-7.40	1.23	1.36
1	A	58	1MA	C2-N3	3.28	1.33	1.29
1	A	20[B]	H2U	C2-N1	3.11	1.40	1.35
1	A	10	2MG	C5-C6	-3.08	1.41	1.47
1	A	34	70U	O9-C8	2.91	1.42	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6	2MG	C5-C6	-2.82	1.41	1.47
1	A	10	2MG	C8-N7	-2.71	1.30	1.35
1	A	37	12A	C6-N6	-2.61	1.31	1.36
1	A	6	2MG	C8-N7	-2.60	1.30	1.35
1	A	37	12A	CC-N6	-2.59	1.32	1.37
1	A	16	H2U	C2-N1	2.59	1.39	1.35
1	A	46	7MG	C5-N7	2.58	1.38	1.35
1	A	46	7MG	CM7-N7	-2.48	1.37	1.45
1	A	10	2MG	C5-C4	-2.25	1.37	1.43
1	A	37	12A	C8-N7	-2.16	1.30	1.34
1	A	20[A]	H2U	C2-N1	2.14	1.38	1.35
1	A	58	1MA	C8-N7	-2.08	1.31	1.35
1	A	58	1MA	C5-C4	-2.03	1.37	1.43
1	A	58	1MA	C2-N1	-2.02	1.32	1.35
1	A	47	H2U	C2-N1	2.01	1.38	1.35

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	46	7MG	N9-C8-N7	6.45	112.60	103.38
1	A	54	2MU	C6'-O2'-C2'	-5.16	100.98	114.52
1	A	34	70U	C9-O9-C8	-4.45	101.45	116.11
1	A	16	H2U	O3'-C3'-C2'	3.83	124.22	111.82
1	A	37	12A	C2-N3-C4	-3.75	110.16	115.32
1	A	46	7MG	C4-C5-N7	3.68	110.64	105.53
1	A	37	12A	C5-C6-N1	-3.54	117.87	120.81
1	A	34	70U	O9-C8-C5M	3.34	117.46	111.27
1	A	55	PSU	C6-N1-C2	3.19	125.95	122.68
1	A	27	PSU	C6-N1-C2	3.17	125.92	122.68
1	A	37	12A	N6-CC-N	3.17	118.19	113.76
1	A	54	2MU	C2'-C1'-N1	-2.89	108.61	114.22
1	A	39	PSU	C6-N1-C2	2.73	125.48	122.68
1	A	16	H2U	O3'-C3'-C4'	2.68	118.79	111.05
1	A	58	1MA	N1-C2-N3	2.64	129.09	126.02
1	A	20[B]	H2U	C4-N3-C2	2.55	127.91	125.79
1	A	20[A]	H2U	C4-N3-C2	2.46	127.83	125.79
1	A	16	H2U	C2'-C3'-C4'	2.42	107.35	102.64
1	A	16	H2U	C4-N3-C2	2.41	127.79	125.79
1	A	20[A]	H2U	O3'-C3'-C2'	-2.33	104.27	111.82
1	A	49	5MC	C5-C6-N1	-2.33	120.94	123.34
1	A	48	5MC	C5-C6-N1	-2.29	120.99	123.34
1	A	58	1MA	C5-C6-N1	2.26	117.27	113.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	2MU	O2'-C2'-C1'	2.21	113.39	109.08
1	A	55	PSU	N1-C2-N3	-2.16	112.68	115.13
1	A	6	2MG	O6-C6-C5	2.12	128.51	124.37
1	A	47	H2U	N3-C2-N1	-2.03	114.50	116.65
1	A	10	2MG	O6-C6-C5	2.02	128.32	124.37

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	6	2MG	N1-C2-N2-CM2
1	A	6	2MG	N3-C2-N2-CM2
1	A	10	2MG	N1-C2-N2-CM2
1	A	10	2MG	N3-C2-N2-CM2
1	A	37	12A	C-CA-N-CC
1	A	37	12A	C-CA-CB-OG1
1	A	37	12A	C-CA-CB-CG2
1	A	47	H2U	C3'-C4'-C5'-O5'
1	A	54	2MU	C1'-C2'-O2'-C6'
1	A	34	70U	O8-C8-O9-C9
1	A	34	70U	C5M-C8-O9-C9
1	A	47	H2U	O4'-C4'-C5'-O5'
1	A	16	H2U	C4'-C5'-O5'-P
1	A	37	12A	C5-C6-N6-CC
1	A	37	12A	OXT-C-CA-N
1	A	47	H2U	C4'-C5'-O5'-P
1	A	48	5MC	C4'-C5'-O5'-P
1	A	20[B]	H2U	O4'-C4'-C5'-O5'
1	A	20[A]	H2U	O4'-C4'-C5'-O5'
1	A	37	12A	O-C-CA-N
1	A	49	5MC	C2'-C1'-N1-C6
1	A	55	PSU	O4'-C1'-C5-C4
1	A	49	5MC	C2'-C1'-N1-C2
1	A	20[A]	H2U	C3'-C4'-C5'-O5'
1	A	20[B]	H2U	C3'-C4'-C5'-O5'
1	A	34	70U	C5-C5M-C8-O9
1	A	37	12A	N-CA-CB-CG2
1	A	37	12A	N1-C2-S2-C2M
1	A	37	12A	N3-C2-S2-C2M

There are no ring outliers.

11 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	48	5MC	1	0
1	A	27	PSU	1	0
1	A	20[B]	H2U	2	0
1	A	37	12A	5	0
1	A	20[A]	H2U	2	0
1	A	49	5MC	2	0
1	A	34	70U	2	0
1	A	10	2MG	8	0
1	A	6	2MG	3	0
1	A	16	H2U	1	0
1	A	54	2MU	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	61/76 (80%)	-0.86	0 100 100	11, 29, 46, 57	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	12A	A	37	34/35	0.92	0.18	16,48,103,109	0
1	H2U	A	16	20/21	0.93	0.27	17,77,120,121	0
1	H2U	A	20[B]	20/21	0.95	0.17	38,67,81,82	20
1	70U	A	34	25/26	0.95	0.14	19,32,68,75	0
1	H2U	A	20[A]	20/21	0.95	0.17	37,66,80,82	20
1	PSU	A	39	20/21	0.95	0.14	3,23,44,49	0
1	H2U	A	47	20/21	0.95	0.16	20,46,89,96	0
1	7MG	A	46	24/25	0.96	0.13	10,33,50,67	0
1	PSU	A	27	20/21	0.97	0.10	16,36,49,65	0
1	2MG	A	10	24/25	0.97	0.12	3,28,44,49	0
1	5MC	A	48	21/22	0.97	0.14	10,33,62,77	0
1	5MC	A	49	21/22	0.97	0.12	9,32,60,65	0
1	2MG	A	6	24/25	0.98	0.09	3,23,40,49	0
1	2MU	A	54	22/23	0.98	0.13	3,17,29,59	0
1	PSU	A	55	20/21	0.98	0.08	3,8,41,45	0
1	1MA	A	58	23/24	0.98	0.11	3,8,28,37	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	A	201	1/1	0.80	0.33	8,8,8,8	0
3	NA	A	202	1/1	0.93	0.21	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.