

# Full wwPDB X-ray Structure Validation Report (i)

Jan 27, 2024 – 09:57 PM EST

PDB ID : 1GDD

Title : TERTIARY AND QUATERNARY STRUCTURAL CHANGES IN GIA1 IN-

DUCED BY GTP HYDROLYSIS

Authors: Mixon, M.B.; Sprang, S.R.

Deposited on : 1995-07-25

Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	A	353	71%	20%	• 8%



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2699 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GI ALPHA 1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace			
1	A	324	Total 2596	C 1647	N 439	O 494	S 16	139	0	0	


• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 5	O 4	S 1	0	0

• Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:  $C_{10}H_{15}N_5O_{11}P_2$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	Р	0	0
3	A	1	28	10	5	11	2	U	0

#### • Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	70	Total O 70 70	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: GI ALPHA 1





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 4	Depositor
Cell constants	121.84Å 121.84Å 68.39Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 - 2.20	Depositor
% Data completeness	95.0 (6.00-2.20)	Depositor
(in resolution range)	35.0 (0.00 2.20)	Беровног
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
$R, R_{free}$	0.214 , $0.291$	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2699	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP



## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bo	nd angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.80	0/2637	0.92	$6/3546 \ (0.2\%)$

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
1	A	15	ARG	NE-CZ-NH2	-6.37	117.11	120.30
1	A	201	VAL	N-CA-C	-6.05	94.65	111.00
1	A	347	ASN	N-CA-C	5.31	125.35	111.00
1	A	36	LEU	CA-CB-CG	5.30	127.50	115.30
1	A	325	CYS	N-CA-C	-5.13	97.16	111.00
1	A	197	LYS	N-CA-C	-5.05	97.38	111.00

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2596	0	2589	45	0
2	A	5	0	0	0	0
3	A	28	0	11	1	0
4	A	70	0	0	0	0
All	All	2699	0	2600	45	0



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 9.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance} ({ m \AA})$	overlap (Å)
1:A:62:SER:OG	1:A:65:GLU:HG3	1.85	0.77
1:A:52:GLN:HE21	1:A:52:GLN:HA	1.53	0.73
1:A:244:HIS:HE1	1:A:286:CYS:O	1.73	0.70
1:A:15:ARG:NH2	1:A:192:LYS:O	2.27	0.68
1:A:51:LYS:HG2	1:A:61:TYR:OH	1.95	0.66
1:A:118:PHE:CZ	1:A:123:LEU:HD22	2.30	0.66
1:A:63:GLU:HG3	1:A:168:ILE:HG21	1.81	0.62
1:A:192:LYS:HD2	1:A:340:THR:HG21	1.83	0.58
1:A:35:LYS:HG3	1:A:199:PHE:HE1	1.71	0.56
1:A:118:PHE:CE2	1:A:123:LEU:HD22	2.42	0.55
1:A:240:MET:HG2	1:A:241:ASN:N	2.24	0.53
1:A:160:ASP:O	1:A:164:GLN:HG2	2.10	0.51
1:A:144:ARG:NE	1:A:231:ASP:O	2.43	0.51
1:A:184:ILE:HD13	1:A:201:VAL:HA	1.93	0.51
1:A:190:THR:HA	1:A:194:LEU:O	2.13	0.48
1:A:62:SER:O	1:A:65:GLU:N	2.47	0.48
1:A:43:GLU:N	3:A:355:GDP:O1B	2.45	0.48
1:A:188:HIS:HA	1:A:196:PHE:O	2.13	0.48
1:A:56:ILE:HG22	1:A:57:HIS:CD2	2.50	0.47
1:A:284:THR:HA	1:A:287:TYR:O	2.15	0.47
1:A:100:ARG:HA	1:A:100:ARG:HH11	1.79	0.47
1:A:226:ALA:HB1	1:A:270:LYS:HD2	1.97	0.47
1:A:282:PRO:O	1:A:285:ILE:HG13	2.14	0.47
1:A:91:LEU:O	1:A:92:LYS:HB2	2.15	0.47
1:A:14:GLU:O	1:A:17:LYS:HG2	2.14	0.46
1:A:116:GLU:HG2	1:A:117:GLY:N	2.30	0.46
1:A:240:MET:HG2	1:A:241:ASN:H	1.80	0.45
1:A:269:ASN:O	1:A:270:LYS:HB2	2.17	0.44
1:A:48:THR:O	1:A:52:GLN:HG2	2.17	0.44
1:A:243:MET:HB3	1:A:286:CYS:SG	2.58	0.44
1:A:275:GLU:O	1:A:278:ILE:HG22	2.18	0.44
1:A:52:GLN:HE21	1:A:52:GLN:CA	2.20	0.43
1:A:62:SER:O	1:A:65:GLU:HB2	2.19	0.43
1:A:184:ILE:CD1	1:A:201:VAL:HA	2.48	0.43
1:A:243:MET:HE3	1:A:303:ILE:HD13	2.01	0.42
1:A:106:GLN:O	1:A:110:LEU:HG	2.19	0.42
1:A:21:ARG:O	1:A:27:GLY:HA3	2.19	0.42

Continued on next page...



$\alpha \cdots$	, r	•	
Continued	trom	mromonie	maaa
-	110116	DICULUUS	Duuc
	J	1	1

Atom-1	Atom-2	$egin{array}{ll}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{array}$	$egin{aligned} \operatorname{Clash} \ \operatorname{overlap}\ (\begin{subarray}{c} \begin{subarray}{c} \begi$
1:A:302:TYR:O	1:A:306:GLN:HG2	2.20	0.42
1:A:36:LEU:O	1:A:198:MET:HA	2.19	0.42
1:A:100:ARG:HA	1:A:100:ARG:HD3	1.68	0.42
1:A:240:MET:SD	1:A:245:GLU:HG2	2.59	0.42
1:A:256:ASN:OD1	1:A:258:TRP:N	2.44	0.41
1:A:143:SER:HA	1:A:146:TYR:CE1	2.55	0.41
1:A:35:LYS:HG3	1:A:199:PHE:CE1	2.53	0.41
1:A:156:LEU:HD23	1:A:156:LEU:HA	1.85	0.41

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Favoured Allowed		Outliers	Percentiles
1	A	318/353 (90%)	310 (98%)	8 (2%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

N	/Iol	Chain	Analysed	Analysed Rotameric Outliers		Percentiles		
	1	A	280/303 (92%)	268 (96%)	12 (4%)	29 36		

All (12) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	9	ASP
1	A	15	ARG
1	A	26	ASP
1	A	52	GLN
1	A	93	ILE
1	A	100	ARG
1	A	143	SER
1	A	158	ASP
1	A	159	LEU
1	A	178	ARG
1	A	181	THR
1	A	257	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	GLN
1	A	106	GLN
1	A	141	ASN
1	A	166	ASN
1	A	195	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trme	Chain	Dag	Link	Bo	ond leng	ths	В	ond ang	les
IVIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	356	-	4,4,4	0.81	0	6,6,6	1.53	1 (16%)
3	GDP	A	355	-	24,30,30	1.78	6 (25%)	30,47,47	1.54	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	355	-	-	2/12/32/32	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
3	A	355	GDP	O4'-C1'	4.71	1.47	1.41
3	A	355	GDP	PB-O3B	-3.17	1.42	1.54
3	A	355	GDP	O3'-C3'	-2.87	1.36	1.43
3	A	355	GDP	PA-O1A	-2.37	1.42	1.50
3	A	355	GDP	C2'-C1'	-2.18	1.50	1.53
3	A	355	GDP	C5'-C4'	-2.11	1.45	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	355	GDP	O6-C6-C5	3.55	131.30	124.37
3	A	355	GDP	O6-C6-N1	-3.52	116.49	120.65
2	A	356	SO4	O4-S-O2	-3.26	92.27	109.31
3	A	355	GDP	O4'-C1'-C2'	-2.65	103.05	106.93
3	A	355	GDP	O5'-C5'-C4'	2.29	116.86	108.99
3	A	355	GDP	C3'-C2'-C1'	2.27	104.40	100.98
3	A	355	GDP	O2'-C2'-C1'	2.13	118.71	110.85

There are no chirality outliers.

All (2) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
3	A	355	GDP	PA-O3A-PB-O2B
3	A	355	GDP	PA-O3A-PB-O3B

There are no ring outliers.

1 monomer is involved in 1 short contact:

$\mathbf{Mol}$	Chain	Res	Type	Clashes	Symm-Clashes
3	A	355	GDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers (i)

There are no such residues in this entry.



# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

#### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

## 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

