



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 9, 2023 – 05:46 pm GMT

PDB ID : 1H83
Title : STRUCTURE OF POLYAMINE OXIDASE IN COMPLEX WITH 1,8-DIAMINO-OCTANE
Authors : Binda, C.; Coda, A.; Angelini, R.; Federico, R.; Ascenzi, P.; Mattevi, A.
Deposited on : 2001-01-24
Resolution : 1.90 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	472	
1	B	472	
1	C	472	
2	D	2	
2	E	2	
3	F	5	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12183 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POLYAMINE OXIDASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	459	Total 3684	C 2353	N 621	O 696	S 14	65	0	0
1	B	462	Total 3715	C 2374	N 627	O 700	S 14	66	0	0
1	C	462	Total 3715	C 2374	N 627	O 700	S 14	58	0	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	2	Total 28	C 16	N 2	O 10	0	0	0
2	E	2	Total 28	C 16	N 2	O 10	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-D-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	5	Total 60	C 34	N 2	O 24	0	0	0

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			
4	A	1	Total	53	27	9	15	2	0	0
4	B	1	Total	53	27	9	15	2	0	0
4	C	1	Total	53	27	9	15	2	0	0

- Molecule 5 is OCTANE 1,8-DIAMINE (three-letter code: DIA) (formula: $C_8H_{20}N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	1
			20	16	4		
5	B	1	Total	C	N	0	1
			20	16	4		
5	C	1	Total	C	N	0	1
			20	16	4		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	223	Total	O	0	0
			223	223		
6	B	242	Total	O	0	0
			242	242		
6	C	269	Total	O	0	0
			269	269		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: POLYAMINE OXIDASE



- Molecule 1: POLYAMINE OXIDASE



- Molecule 1: POLYAMINE OXIDASE



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2

- Molecule 3: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-D-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  40% 60%

MAG1
MAG2
MAN3
MAN4
FCA5

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	184.69Å 184.69Å 282.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.90	Depositor
% Data completeness (in resolution range)	97.9 (20.00-1.90)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT 5D	Depositor
R, R_{free}	0.196 , 0.237	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	12183	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FCA, DIA, FAD, MAN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/3775	1.21	15/5116 (0.3%)
1	B	0.55	0/3808	1.21	15/5160 (0.3%)
1	C	0.56	0/3808	1.22	17/5160 (0.3%)
All	All	0.56	0/11391	1.21	47/15436 (0.3%)

There are no bond length outliers.

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	355	ARG	NE-CZ-NH2	-10.02	115.29	120.30
1	A	355	ARG	NE-CZ-NH2	-8.79	115.91	120.30
1	C	176	ARG	NE-CZ-NH2	-8.47	116.06	120.30
1	A	424	ARG	NE-CZ-NH1	8.23	124.42	120.30
1	C	176	ARG	NE-CZ-NH1	8.12	124.36	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3684	0	3585	56	0
1	B	3715	0	3614	47	0
1	C	3715	0	3614	41	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	28	0	26	5	0
2	E	28	0	26	4	0
3	F	60	0	53	6	0
4	A	53	0	31	2	0
4	B	53	0	31	1	0
4	C	53	0	31	4	0
5	A	20	0	40	5	0
5	B	20	0	40	4	0
5	C	20	0	40	5	0
6	A	223	0	0	7	1
6	B	242	0	0	4	0
6	C	269	0	0	3	2
All	All	12183	0	11131	158	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 158 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:590[A]:DIA:HC82	6:C:2227:HOH:O	1.55	1.06
1:A:273:SER:O	1:A:274:ASP:HB2	1.72	0.88
3:F:1:NAG:C4	3:F:2:NAG:C1	2.54	0.86
2:E:1:NAG:C4	2:E:2:NAG:C1	2.55	0.84
1:C:91:TYR:OH	1:C:314:LYS:HE2	1.80	0.81

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:C:2094:HOH:O	6:C:2094:HOH:O[10_665]	0.73	1.47
6:A:2131:HOH:O	6:C:2058:HOH:O[6_654]	1.30	0.90

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/472 (97%)	438 (96%)	18 (4%)	1 (0%)	47	38
1	B	460/472 (98%)	444 (96%)	16 (4%)	0	100	100
1	C	460/472 (98%)	442 (96%)	18 (4%)	0	100	100
All	All	1377/1416 (97%)	1324 (96%)	52 (4%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	101	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	394/404 (98%)	387 (98%)	7 (2%)	59	55
1	B	397/404 (98%)	390 (98%)	7 (2%)	59	55
1	C	397/404 (98%)	392 (99%)	5 (1%)	69	68
All	All	1188/1212 (98%)	1169 (98%)	19 (2%)	62	60

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	59	ASN
1	C	149	HIS
1	C	175	PRO
1	C	127	SER
1	B	127	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	466	HIS
1	C	360	GLN
1	C	431	HIS
1	C	152	ASN
1	B	152	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	D	1	1,2	14,14,15	0.94	1 (7%)	17,19,21	1.44	4 (23%)
2	NAG	D	2	2	14,14,15	0.88	1 (7%)	17,19,21	1.77	4 (23%)
2	NAG	E	1	1,2	14,14,15	1.21	2 (14%)	17,19,21	1.58	3 (17%)
2	NAG	E	2	2	14,14,15	0.95	1 (7%)	17,19,21	1.33	2 (11%)
3	NAG	F	1	3,1	14,14,15	1.06	2 (14%)	17,19,21	2.20	7 (41%)
3	NAG	F	2	3	14,14,15	0.96	1 (7%)	17,19,21	1.53	4 (23%)
3	MAN	F	3	3	11,11,12	0.72	0	15,15,17	3.03	5 (33%)
3	MAN	F	4	3	11,11,12	0.55	0	15,15,17	1.59	3 (20%)
3	FCA	F	5	3	10,10,11	1.21	1 (10%)	14,14,16	2.12	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	NAG	E	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1
3	NAG	F	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
3	MAN	F	3	3	-	0/2/19/22	0/1/1/1
3	MAN	F	4	3	-	2/2/19/22	0/1/1/1
3	FCA	F	5	3	-	-	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	2	NAG	O5-C1	-2.93	1.39	1.43
2	E	1	NAG	C1-C2	-2.86	1.48	1.52
3	F	5	FCA	C2-C3	-2.69	1.48	1.52
2	E	1	NAG	O5-C1	-2.57	1.39	1.43
2	E	2	NAG	O5-C1	-2.48	1.39	1.43

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	3	MAN	O2-C2-C3	10.34	130.85	110.14
2	D	2	NAG	C2-N2-C7	-5.10	115.64	122.90
3	F	1	NAG	C1-C2-N2	-4.66	102.53	110.49
3	F	5	FCA	O5-C1-C2	-4.40	103.98	110.77
3	F	4	MAN	C1-C2-C3	-4.32	104.35	109.67

There are no chirality outliers.

5 of 14 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	2	NAG	C4-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	E	1	NAG	C4-C5-C6-O6
3	F	2	NAG	C4-C5-C6-O6


There are no ring outliers.


7 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	2	NAG	4	0
3	F	5	FCA	3	0
2	D	1	NAG	5	0
2	E	1	NAG	4	0
3	F	2	NAG	4	0
2	D	2	NAG	5	0
3	F	1	NAG	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	FAD	A	579	-	53,58,58	0.87	2 (3%)	68,89,89	1.47	11 (16%)
5	DIA	A	590[B]	-	9,9,9	0.51	0	8,8,8	0.30	0
4	FAD	C	579	-	53,58,58	1.05	1 (1%)	68,89,89	1.29	8 (11%)
5	DIA	C	590[B]	-	9,9,9	0.50	0	8,8,8	0.35	0
4	FAD	B	579	-	53,58,58	0.80	1 (1%)	68,89,89	1.26	7 (10%)
5	DIA	B	590[A]	-	9,9,9	0.48	0	8,8,8	0.44	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	DIA	A	590[A]	-	9,9,9	0.47	0	8,8,8	0.28	0
5	DIA	C	590[A]	-	9,9,9	0.49	0	8,8,8	0.42	0
5	DIA	B	590[B]	-	9,9,9	0.51	0	8,8,8	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAD	A	579	-	-	2/30/50/50	0/6/6/6
5	DIA	A	590[B]	-	-	3/7/7/7	-
4	FAD	C	579	-	-	1/30/50/50	0/6/6/6
5	DIA	C	590[B]	-	-	4/7/7/7	-
4	FAD	B	579	-	-	5/30/50/50	0/6/6/6
5	DIA	B	590[A]	-	-	5/7/7/7	-
5	DIA	A	590[A]	-	-	4/7/7/7	-
5	DIA	C	590[A]	-	-	6/7/7/7	-
5	DIA	B	590[B]	-	-	3/7/7/7	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	579	FAD	C4X-N5	3.49	1.37	1.30
4	B	579	FAD	C4X-N5	2.55	1.35	1.30
4	A	579	FAD	C4X-N5	2.50	1.35	1.30
4	A	579	FAD	C5X-N5	-2.10	1.35	1.39

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	579	FAD	C9A-C5X-N5	-5.36	116.60	122.43
4	C	579	FAD	O4B-C1B-C2B	-4.21	100.77	106.93
4	B	579	FAD	O4B-C1B-C2B	-3.94	101.17	106.93
4	A	579	FAD	O4'-C4'-C3'	3.12	116.68	109.10
4	A	579	FAD	C6-C5X-C9A	3.07	123.28	118.94

There are no chirality outliers.

5 of 33 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	579	FAD	C5B-O5B-PA-O1A
5	A	590[B]	DIA	C2-C3-C4-C5
5	B	590[A]	DIA	C3-C4-C5-C6
5	B	590[B]	DIA	C2-C3-C4-C5
5	B	590[A]	DIA	C2-C3-C4-C5


There are no ring outliers.


9 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	579	FAD	2	0
5	A	590[B]	DIA	1	0
4	C	579	FAD	4	0
5	C	590[B]	DIA	2	0
4	B	579	FAD	1	0
5	B	590[A]	DIA	3	0
5	A	590[A]	DIA	4	0
5	C	590[A]	DIA	3	0
5	B	590[B]	DIA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.