

# Full wwPDB X-ray Structure Validation Report (i)

May 29, 2020 – 07:44 am BST

PDB ID : 1HOS

Title : INHIBITION OF HUMAN IMMUNODEFICIENCY VIRUS-1 PROTEASE

BY A C2-SYMMETRIC PHOSPHINATE SYNTHESIS AND CRYSTALLO-

GRAPHIC ANALYSIS

Authors: Abdel-Meguid, S.; Zhao, B.

Deposited on : 1993-04-06

Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at  $\frac{\text{https://www.wwpdb.org/validation/2017/XrayValidationReportHelp}}{\text{with specific help available everywhere you see the (i) symbol.}$ 

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

EDS: NOT EXECUT buster-report: 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{resolution range}( ext{Å}))$
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain					
1	A	99	60%	36%				
1	В	99	60%	29%	8% •			



# 2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 1663 atoms, of which 10 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called HIV-1 PROTEASE.

Mol	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
1	A	99	Total 758	C 489	- 1	O 135	S 4	0	0	0
1	В	99	Total 758	C 489	N 130	O 135	S 4	0	0	0

• Molecule 2 is (2-PHENYL-1-CARBOBENZYL-OXYVALYL-AMINO)-ETHYL-PHOSPHI NIC ACID (three-letter code: PHP) (formula: C<sub>42</sub>H<sub>51</sub>N<sub>4</sub>O<sub>8</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
2	A	1	Total 120	C 84	H 10	N 8	O 16	P 2	0	1

• Molecule 3 is water.

$\mathbf{Mol}$	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	14	Total O 14 14	0	0



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	13	Total O 13 13	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: HIV-1 PROTEASE





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 61	Depositor	
Cell constants	63.20Å 63.20Å 83.40Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor	
Resolution (Å)	6.00 - 2.30	Depositor	
% Data completeness	(Not available) (6.00-2.30)	Depositor	
(in resolution range)	, , , , , , , , , , , , , , , , , , , ,	Depositor	
$R_{merge}$	(Not available)	Depositor	
$R_{sym}$	(Not available)	Depositor	
Refinement program	X-PLOR	Depositor	
$R, R_{free}$	0.186 , (Not available)	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	1663	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP	



# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PHP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Bond angles		
Moi Chain		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	1.07	1/771~(0.1%)	1.94	19/1043 (1.8%)	
1	В	1.01	1/771 (0.1%)	1.87	25/1043~(2.4%)	
All	All	1.04	$2/1542 \ (0.1\%)$	1.90	$44/2086 \ (2.1\%)$	

#### All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$Ideal(\AA)$
1	A	37	SER	CB-OG	10.85	1.56	1.42
1	В	37	SER	CB-OG	5.95	1.50	1.42

#### All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	В	6	TRP	CD1-CG-CD2	9.32	113.76	106.30
1	A	57	ARG	NE-CZ-NH1	9.00	124.80	120.30
1	A	6	TRP	CD1-CG-CD2	8.73	113.29	106.30
1	В	6	TRP	CE2-CD2-CG	-8.42	100.56	107.30
1	A	42	TRP	CD1-CG-CD2	8.11	112.79	106.30
1	A	57	ARG	NE-CZ-NH2	-8.04	116.28	120.30
1	В	36	MET	CG-SD-CE	-7.95	87.48	100.20
1	A	6	TRP	CE2-CD2-CG	-7.86	101.01	107.30
1	A	42	TRP	CE2-CD2-CG	-7.84	101.03	107.30
1	A	25	ASP	N-CA-CB	-7.57	96.97	110.60
1	В	59	TYR	CB-CG-CD1	-7.52	116.49	121.00
1	A	70	LYS	CB-CG-CD	-7.23	92.81	111.60
1	В	42	TRP	CE2-CD2-CG	-6.57	102.05	107.30
1	A	59	TYR	CB-CG-CD1	-6.52	117.09	121.00
1	A	29	ASP	CB-CG-OD2	-6.44	112.50	118.30
1	A	25	ASP	N-CA-C	6.26	127.92	111.00
1	В	65	GLU	CA-CB-CG	-6.25	99.64	113.40



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\mathbf{Ideal}(^o)$
1	В	42	TRP	CB-CG-CD1	-6.18	118.97	127.00
1	В	33	LEU	CA-CB-CG	6.07	129.25	115.30
1	A	32	VAL	CA-CB-CG2	-5.92	102.03	110.90
1	В	8	ARG	NE-CZ-NH2	5.74	123.17	120.30
1	В	2	GLN	CA-CB-CG	-5.73	100.79	113.40
1	В	25	ASP	N-CA-CB	-5.69	100.36	110.60
1	В	32	VAL	CG1-CB-CG2	-5.65	101.86	110.90
1	В	50	ILE	CB-CG1-CD1	-5.63	98.12	113.90
1	В	25	ASP	N-CA-C	5.56	126.02	111.00
1	В	42	TRP	CD1-CG-CD2	5.56	110.75	106.30
1	В	23	LEU	CA-CB-CG	5.52	128.00	115.30
1	A	29	ASP	CB-CG-OD1	5.48	123.23	118.30
1	В	6	TRP	CB-CG-CD1	-5.45	119.92	127.00
1	В	42	TRP	CG-CD2-CE3	5.41	138.77	133.90
1	A	58	GLN	CA-CB-CG	5.36	125.20	113.40
1	В	25	ASP	CB-CG-OD1	5.35	123.11	118.30
1	A	52	GLY	CA-C-N	-5.29	105.56	117.20
1	A	37	SER	N-CA-C	-5.26	96.79	111.00
1	В	32	VAL	CA-CB-CG1	5.24	118.75	110.90
1	В	41	ARG	NE-CZ-NH2	-5.23	117.68	120.30
1	В	6	TRP	CG-CD2-CE3	5.21	138.58	133.90
1	A	32	VAL	CA-CB-CG1	5.18	118.66	110.90
1	A	42	TRP	CB-CG-CD1	-5.18	120.27	127.00
1	A	52	GLY	O-C-N	5.13	130.90	122.70
1	В	85	ILE	N-CA-C	-5.06	97.33	111.00
1	В	11	VAL	N-CA-CB	-5.03	100.43	111.50
1	В	97	LEU	N-CA-C	-5.03	97.43	111.00

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	758	0	806	21	15
1	В	758	0	806	21	15



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
2	A	110	10	100	6	0
3	A	14	0	0	0	0
3	В	13	0	0	0	0
All	All	1653	10	1712	35	15

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

	1	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ ({\rm \AA})$	overlap(Å)
1:A:56:VAL:HG21	1:A:76:LEU:HD13	1.69	0.74
1:A:99:PHE:HE1	1:B:3:ILE:HD11	1.56	0.70
1:A:45:LYS:HD2	1:A:58:GLN:HE21	1.64	0.63
1:A:34:GLU:HG3	1:A:83:ASN:OD1	2.00	0.61
1:A:54:ILE:HD12	1:A:79:PRO:HD2	1.85	0.58
1:A:3:ILE:HD11	1:B:99:PHE:HE1	1.71	0.56
1:A:50:ILE:HG12	1:B:49:GLY:HA3	1.88	0.56
2:A:400[A]:PHP:HG54	1:B:47:ILE:HD12	1.86	0.56
1:A:32:VAL:HG23	1:A:76:LEU:HB2	1.89	0.54
1:B:59:TYR:O	1:B:74:THR:HA	2.09	0.53
1:B:20:LYS:NZ	1:B:36:MET:SD	2.82	0.51
1:B:13:ILE:HG22	1:B:66:ILE:HA	1.92	0.51
1:B:34:GLU:O	1:B:36:MET:HG2	2.11	0.51
1:A:87:ARG:HD2	1:B:5:LEU:O	2.11	0.50
1:A:87:ARG:HG2	1:B:5:LEU:HB3	1.95	0.48
2:A:400[B]:PHP:HG24	1:B:47:ILE:HD12	1.97	0.47
1:A:72:ILE:H	1:A:92:GLN:HE22	1.61	0.47
1:A:59:TYR:HB3	1:A:62:ILE:HD11	1.97	0.47
1:B:33:LEU:HD22	1:B:83:ASN:HB3	1.96	0.46
1:A:45:LYS:HG2	1:A:46:MET:N	2.31	0.46
1:A:81:PRO:HG2	2:A:400[A]:PHP:HZ4	1.97	0.46
2:A:400[A]:PHP:HB61	1:B:48:GLY:O	2.15	0.46
2:A:400[A]:PHP:HD62	1:B:48:GLY:HA3	1.99	0.45
1:A:99:PHE:HE1	1:B:3:ILE:CD1	2.27	0.45
1:B:11:VAL:HG22	1:B:67:CYS:SG	2.58	0.44
1:A:97:LEU:HD21	1:B:24:LEU:HB3	2.00	0.44
1:B:54:ILE:HD12	1:B:79:PRO:HG2	2.00	0.43
1:B:13:ILE:HG21	1:B:85:ILE:HD11	2.00	0.43
1:A:13:ILE:HG22	1:A:66:ILE:HA	2.01	0.42
2:A:400[B]:PHP:CZ6	1:B:8:ARG:HG2	2.50	ed on next page



$\alpha \cdots \tau$	r	•	
Continued	trom	nromanne	naae
$\circ$	110116	picolous	puyc

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{aligned}$	Clash overlap (Å)	
1:A:24:LEU:HD12	1:A:24:LEU:HA	1.88	0.42	
1:A:13:ILE:HG21	1:A:85:ILE:HD11	2.03	0.41	
1:A:50:ILE:HD13	1:A:50:ILE:HG21	1.86	0.41	
1:B:42:TRP:HE3	1:B:44:PRO:HD3	1.85	0.40	
1:A:45:LYS:HG2	1:A:46:MET:H	1.85	0.40	

All (15) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic	Clash
		$\operatorname{distance}\left( \mathrm{\AA}\right)$	overlap(A)
1:A:53:PHE:CZ	1:B:53:PHE:CE2[6_555]	0.32	1.88
1:A:53:PHE:CE2	1:B:53:PHE:CZ[6_555]	0.55	1.65
1:A:53:PHE:CE2	1:B:53:PHE:CE2[6_555]	1.07	1.13
1:A:53:PHE:CZ	1:B:53:PHE:CD2[6_555]	1.27	0.93
1:A:53:PHE:CE1	1:B:53:PHE:CD2[6_555]	1.29	0.91
1:A:53:PHE:CE2	1:B:53:PHE:CE1[6_555]	1.43	0.77
1:A:53:PHE:CE1	1:B:53:PHE:CE2[6_555]	1.59	0.61
1:A:53:PHE:CD2	1:B:53:PHE:CE1[6_555]	1.65	0.55
1:A:53:PHE:CZ	1:B:53:PHE:CZ[6_555]	1.68	0.52
1:A:53:PHE:CD2	1:B:53:PHE:CZ[6_555]	1.74	0.46
1:A:53:PHE:CE2	1:B:53:PHE:CD2[6_555]	1.95	0.25
1:A:53:PHE:CD1	1:B:53:PHE:CD2[6_555]	1.98	0.22
1:A:53:PHE:CD2	1:B:53:PHE:CD1[6_555]	2.03	0.17
1:A:53:PHE:CD2	1:B:53:PHE:CE2[6_555]	2.15	0.05
1:A:53:PHE:CE2	1:B:53:PHE:CD1[6_555]	2.18	0.02

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Allowed   Outliers		Percentiles		
1	A	97/99 (98%)	95 (98%)	2 (2%)	0	100 100			



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	В	97/99 (98%)	92 (95%)	5 (5%)	0	100	100
All	All	194/198 (98%)	187 (96%)	7 (4%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	83/83 (100%)	75 (90%)	8 (10%)	8 10		
1	В	83/83 (100%)	71 (86%)	12 (14%)	3 3		
All	All	166/166 (100%)	146 (88%)	20 (12%)	5 5		

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	11	VAL
1	A	24	LEU
1	A	44	PRO
1	A	55	LYS
1	A	61	GLN
1	A	91	THR
1	A	98	ASN
1	В	1	PRO
1	В	7	GLN
1	В	11	VAL
1	В	23	LEU
1	В	24	LEU
1	В	33	LEU
1	В	35	GLU
1	В	36	MET
1	В	41	ARG
1	В	44	PRO



Continued from previous page...

Mol	Chain	Res	Type
1	В	46	MET
1	В	55	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	92	GLN
1	A	98	ASN
1	В	2	GLN
1	В	92	GLN
1	В	98	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain		$_{ m ain} \mid _{ m Res} \mid$	Link	Bo	Bond lengths		Bond angles			
	Type	Chain	i nes	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
	2	РНР	A	400[A]	-	56,58,58	0.65	1 (1%)	70,78,78	1.58	10 (14%)



Mol	Type	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain Res	Link	Bond lengths			Bond angles		
10101	Type		nes	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2								
2	PHP	A	400[B]	_	56,58,58	0.64	2 (3%)	70,78,78	1.55	12 (17%)								

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

$\mathbf{Mol}$	Type	Chain	$\operatorname{Res}$	Link	Chirals	${f Torsions}$	Rings
2	PHP	A	400[A]	-	ı	9/54/62/62	0/4/4/4
2	PHP	A	400[B]	-	-	10/54/62/62	0/4/4/4

#### All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
2	A	400[A]	PHP	P3-O3'	-2.74	1.49	1.54
2	A	400[B]	PHP	CB6-CG6	2.11	1.55	1.50
2	A	400[B]	PHP	P3-O3'	-2.04	1.50	1.54

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}({}^o)$	$\operatorname{Ideal}({}^{o})$
2	A	400[A]	РНР	OA6-C6-N5	5.45	121.59	110.50
2	A	400[A]	PHP	CB6-OA6-C6	4.92	126.92	115.93
2	A	400[B]	PHP	OA6-CB6-CG6	4.82	120.97	109.39
2	A	400[B]	PHP	C5-CA5-N5	-4.47	98.20	110.36
2	A	400[A]	PHP	C2-CA2-N2	-4.42	98.35	110.36
2	A	400[A]	PHP	C5-CA5-N5	-3.96	99.59	110.36
2	A	400[B]	PHP	CB6-OA6-C6	3.73	124.26	115.93
2	A	400[A]	PHP	OA6-C6-O6	-3.61	117.31	124.25
2	A	400[B]	PHP	O3'-P3-CA3	3.50	114.73	106.87
2	A	400[B]	PHP	C2-CA2-N2	-3.49	100.86	110.36
2	A	400[B]	PHP	OA1-C1-N2	3.39	117.39	110.50
2	A	400[A]	PHP	O3'-P3-CA3	3.33	114.35	106.87
2	A	400[A]	PHP	CGM-CB2-CGL	-3.15	101.78	110.59
2	A	400[B]	PHP	OA6-C6-N5	2.95	116.51	110.50
2	A	400[B]	РНР	CB1-OA1-C1	2.74	122.06	115.93
2	A	400[B]	PHP	CA5-N5-C6	2.59	126.53	120.86
2	A	400[B]	PHP	CGG-CB5-CGF	-2.51	103.56	110.59
2	A	400[A]	PHP	CG4-CB4-CA4	-2.45	107.98	113.38
2	A	400[B]	PHP	CG4-CB4-CA4	-2.40	108.08	113.38
2	A	400[A]	PHP	OA1-C1-N2	2.34	115.27	110.50



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^o)$
2	A	400[A]	PHP	O6-C6-N5	-2.29	121.10	124.85
2	A	400[B]	PHP	O1-C1-N2	-2.21	121.22	124.85

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	400[A]	PHP	CB3-CA3-P3-O3
2	A	400[A]	PHP	N4-CA4-P3-O3
2	A	400[A]	PHP	CB4-CA4-P3-O3
2	A	400[A]	PHP	N5-C6-OA6-CB6
2	A	400[A]	PHP	O6-C6-OA6-CB6
2	A	400[B]	PHP	CB3-CA3-P3-O3
2	A	400[B]	PHP	N4-CA4-P3-O3
2	A	400[B]	PHP	CB4-CA4-P3-O3
2	A	400[A]	PHP	N2-C1-OA1-CB1
2	A	400[A]	PHP	O1-C1-OA1-CB1
2	A	400[A]	PHP	N3-CA3-P3-O3
2	A	400[B]	PHP	N3-CA3-P3-O3
2	A	400[B]	PHP	N4-C5-CA5-N5
2	A	400[B]	PHP	CG6-CB6-OA6-C6
2	A	400[B]	PHP	O5-C5-CA5-N5
2	A	400[B]	PHP	CG1-CB1-OA1-C1
2	A	400[B]	PHP	O5-C5-CA5-CB5
2	A	400[B]	PHP	N4-C5-CA5-CB5
2	A	400[A]	PHP	CG6-CB6-OA6-C6

There are no ring outliers.

2 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	400[A]	PHP	4	0
2	A	400[B]	PHP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the



average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

