



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 23, 2021 – 02:19 PM EST

PDB ID : 1JSS  
Title : Crystal structure of the Mus musculus cholesterol-regulated START protein 4 (StarD4).  
Authors : Romanowski, M.J.; Soccio, R.E.; Breslow, J.L.; Burley, S.K.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2001-08-17  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	 63% 25% 11%
1	B	224	 63% 23% 11%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3498 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cholesterol-regulated START protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	199	1604	1017	272	305	10	0	0	0
1	B	199	1604	1017	272	305	10	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	GLU	LYS	SEE REMARK 999	UNP Q99JV5
A	51	ALA	VAL	SEE REMARK 999	UNP Q99JV5
B	41	GLU	LYS	SEE REMARK 999	UNP Q99JV5
B	51	ALA	VAL	SEE REMARK 999	UNP Q99JV5

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	143	Total	O	0	0
			143	143		
2	B	147	Total	O	0	0
			147	147		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.97Å 69.97Å 85.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.55 – 2.20 28.54 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.5 (28.55-2.20) 99.7 (28.54-2.20)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.49 (at 2.20Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.238 , 0.285 0.239 , 0.284	Depositor DCC
$R_{free}$ test set	4543 reflections (9.64%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.2	Xtrriage
Anisotropy	0.010	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 33.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.022 for -h,-k,l 0.479 for h,-h-k,-l 0.021 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3498	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1644	0.72	3/2234 (0.1%)
1	B	0.41	2/1644 (0.1%)	0.73	3/2234 (0.1%)
All	All	0.41	2/3288 (0.1%)	0.72	6/4468 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	143	GLU	N-CA	6.06	1.58	1.46
1	B	113	CYS	CB-SG	5.47	1.91	1.82

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	143	GLU	N-CA-C	7.74	131.90	111.00
1	B	113	CYS	CA-CB-SG	6.44	125.59	114.00
1	B	143	GLU	N-CA-CB	-6.39	99.10	110.60
1	A	143	GLU	N-CA-C	6.31	128.04	111.00
1	A	113	CYS	CA-CB-SG	5.46	123.83	114.00
1	A	143	GLU	N-CA-CB	-5.41	100.86	110.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1604	0	1550	51	0
1	B	1604	0	1550	56	0
2	A	143	0	0	4	0
2	B	147	0	0	8	0
All	All	3498	0	3100	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:143:GLU:HG3	1:B:175:PRO:HD2	1.23	1.14
1:B:179:SER:OG	1:B:180:PRO:HD2	1.70	0.90
1:B:177:LYS:HG2	1:B:178:ASP:H	1.42	0.84
1:B:143:GLU:CG	1:B:175:PRO:HD2	2.10	0.77
1:B:34:LEU:HD21	1:B:170:GLY:HA3	1.67	0.76
1:B:116:ARG:HD3	1:B:158:ARG:NH2	2.01	0.75
1:B:176:LEU:HD12	1:B:182:GLN:HE21	1.49	0.74
1:B:176:LEU:HD22	1:B:177:LYS:HD2	1.74	0.69
1:A:98:LEU:HG	1:A:119:THR:OG1	1.94	0.68
1:B:44:GLU:HG2	1:B:59:LYS:HE2	1.75	0.67
1:A:46:ARG:HG3	1:A:46:ARG:HH11	1.58	0.67
1:A:40:ILE:HG21	1:A:59:LYS:HZ1	1.60	0.66
1:B:176:LEU:HD12	1:B:182:GLN:NE2	2.12	0.65
1:B:176:LEU:O	1:B:177:LYS:C	2.36	0.64
1:B:189:ILE:HG13	1:B:210:LEU:HD11	1.81	0.63
1:B:45:TRP:HB3	1:B:57:TRP:HB3	1.81	0.63
1:B:47:VAL:HG11	1:B:50:LYS:HE3	1.81	0.63
1:A:40:ILE:HD13	1:A:59:LYS:NZ	2.15	0.62
1:B:177:LYS:HG2	1:B:178:ASP:N	2.15	0.61
1:B:34:LEU:CD2	1:B:170:GLY:HA3	2.32	0.59
1:B:44:GLU:OE2	1:B:59:LYS:HE3	2.02	0.59
1:B:34:LEU:HD22	1:B:148:CYS:HB3	1.84	0.58
1:B:46:ARG:HD2	1:B:46:ARG:N	2.19	0.58
1:A:119:THR:CG2	1:A:130:ARG:HB2	2.34	0.58
1:B:91:TRP:HE3	1:B:94:ASP:OD2	1.87	0.58
1:A:108:PHE:HE1	1:A:114:VAL:HG23	1.68	0.58
1:B:176:LEU:O	1:B:178:ASP:N	2.37	0.58
1:A:91:TRP:HE3	1:A:94:ASP:OD2	1.87	0.57
1:B:34:LEU:HD21	1:B:170:GLY:CA	2.35	0.57
1:B:24:ALA:HB3	2:B:233:HOH:O	2.04	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:SER:HG	1:B:180:PRO:HD2	1.70	0.57
1:B:219:LYS:HE3	2:B:294:HOH:O	2.04	0.57
1:B:106:GLU:HB2	1:B:114:VAL:HB	1.85	0.56
1:B:123:LEU:HB2	1:B:126:ILE:HB	1.87	0.56
1:A:33:THR:O	1:A:36:GLN:HB3	2.06	0.56
1:A:119:THR:HG22	1:A:130:ARG:HB2	1.87	0.56
1:A:119:THR:HG23	2:A:239:HOH:O	2.07	0.55
1:A:45:TRP:CZ3	1:A:59:LYS:HG2	2.42	0.55
1:A:87:ARG:HB3	1:A:88:PRO:HD2	1.88	0.55
1:B:206:MET:HA	1:B:209:THR:HG22	1.88	0.54
1:A:221:LEU:O	1:A:222:ARG:HB2	2.08	0.54
1:B:176:LEU:HD22	1:B:177:LYS:CD	2.37	0.54
1:B:88:PRO:HB3	1:B:102:LEU:HD21	1.90	0.53
1:A:41:GLU:HA	1:A:41:GLU:OE1	2.09	0.52
1:A:130:ARG:CZ	1:A:162:VAL:HG11	2.40	0.52
1:A:53:ASP:OD1	1:A:74:VAL:HG12	2.10	0.51
1:A:88:PRO:HB3	1:A:102:LEU:HD21	1.93	0.51
1:A:176:LEU:HD23	1:A:182:GLN:HE21	1.75	0.50
1:A:105:LEU:HD11	1:A:116:ARG:HB2	1.92	0.50
1:A:158:ARG:HB3	2:A:283:HOH:O	2.10	0.50
1:B:158:ARG:HB3	2:B:248:HOH:O	2.12	0.50
1:A:61:SER:OG	1:A:190:GLN:NE2	2.42	0.49
1:B:87:ARG:NH2	2:B:260:HOH:O	2.44	0.49
1:A:93:LEU:CD2	1:A:99:MET:HG2	2.42	0.49
1:A:177:LYS:HG2	1:A:178:ASP:OD1	2.13	0.49
1:A:108:PHE:CE1	1:A:114:VAL:HG23	2.47	0.49
1:A:154:TRP:HB3	1:A:163:ARG:HH12	1.77	0.48
1:B:87:ARG:HB3	1:B:88:PRO:HD2	1.95	0.48
1:B:177:LYS:N	1:B:177:LYS:HD2	2.29	0.47
1:B:43:ASP:OD1	1:B:43:ASP:O	2.33	0.47
1:A:46:ARG:HG3	1:A:46:ARG:NH1	2.27	0.47
1:A:119:THR:HG22	1:A:130:ARG:H	1.80	0.46
1:B:221:LEU:O	1:B:222:ARG:HB2	2.15	0.46
1:A:176:LEU:O	1:A:177:LYS:C	2.53	0.46
1:B:79:VAL:HG23	2:B:307:HOH:O	2.16	0.46
1:A:40:ILE:CG2	1:A:59:LYS:HZ1	2.25	0.45
1:B:80:ASN:N	1:B:80:ASN:HD22	2.13	0.45
1:B:206:MET:HA	1:B:209:THR:CG2	2.46	0.45
1:B:142:GLU:OE1	1:B:142:GLU:HA	2.17	0.45
1:A:222:ARG:HG3	1:A:222:ARG:HH11	1.82	0.45
1:B:126:ILE:HD11	2:B:334:HOH:O	2.15	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:LEU:CD2	1:A:182:GLN:HE21	2.30	0.45
1:A:38:HIS:HA	1:A:188:TYR:OH	2.17	0.45
1:B:80:ASN:HD22	1:B:80:ASN:H	1.65	0.44
1:A:163:ARG:HD3	2:A:228:HOH:O	2.17	0.44
1:A:24:ALA:HB1	2:A:242:HOH:O	2.16	0.44
1:B:44:GLU:HG2	1:B:59:LYS:CE	2.43	0.44
1:B:221:LEU:O	1:B:222:ARG:CB	2.65	0.44
1:B:206:MET:O	1:B:209:THR:HG22	2.18	0.44
1:A:116:ARG:HA	1:A:132:PHE:O	2.18	0.43
1:A:54:VAL:HG21	1:A:214:TYR:CB	2.48	0.43
1:B:160:GLU:HG2	2:B:248:HOH:O	2.18	0.43
1:A:91:TRP:CD1	1:A:91:TRP:N	2.86	0.43
1:B:54:VAL:HG21	1:B:214:TYR:CB	2.48	0.43
1:A:83:ILE:HG22	1:A:87:ARG:HD2	2.01	0.43
1:B:217:LEU:O	1:B:221:LEU:HD13	2.18	0.43
1:B:61:SER:OG	1:B:190:GLN:NE2	2.43	0.43
1:A:139:VAL:HG22	1:A:140:GLY:N	2.34	0.42
1:B:46:ARG:N	1:B:46:ARG:CD	2.82	0.42
1:B:179:SER:HB3	2:B:289:HOH:O	2.18	0.42
1:B:41:GLU:HG2	1:B:43:ASP:H	1.84	0.42
1:A:69:TYR:O	1:A:188:TYR:HA	2.20	0.42
1:A:80:ASN:H	1:A:80:ASN:HD22	1.68	0.42
1:A:40:ILE:HD13	1:A:59:LYS:HZ2	1.83	0.42
1:B:206:MET:CA	1:B:209:THR:HG22	2.50	0.42
1:A:119:THR:HG22	1:A:130:ARG:CB	2.50	0.41
1:A:133:VAL:HG11	1:A:154:TRP:HB2	2.02	0.41
1:A:154:TRP:O	1:A:163:ARG:NH1	2.52	0.41
1:A:116:ARG:HG3	1:A:133:VAL:HG22	2.02	0.41
1:A:42:GLU:HA	1:A:45:TRP:HD1	1.85	0.41
1:B:116:ARG:HA	1:B:132:PHE:O	2.21	0.41
1:A:45:TRP:HB3	1:A:57:TRP:HB3	2.04	0.40
1:B:124:LEU:O	1:B:125:ASN:HB2	2.20	0.40
1:A:155:SER:O	1:A:157:THR:HG23	2.21	0.40
1:A:167:HIS:O	1:A:168:PRO:C	2.57	0.40
1:A:222:ARG:HG3	1:A:222:ARG:NH1	2.37	0.40
1:B:24:ALA:C	1:B:26:ILE:H	2.24	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/224 (88%)	180 (91%)	14 (7%)	3 (2%)	10	8
1	B	197/224 (88%)	178 (90%)	17 (9%)	2 (1%)	15	14
All	All	394/448 (88%)	358 (91%)	31 (8%)	5 (1%)	12	9

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	177	LYS
1	B	177	LYS
1	B	180	PRO
1	A	42	GLU
1	A	90	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/199 (89%)	173 (97%)	5 (3%)	43	56
1	B	178/199 (89%)	167 (94%)	11 (6%)	18	21
All	All	356/398 (89%)	340 (96%)	16 (4%)	27	34

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	34	LEU
1	A	58	ARG
1	A	94	ASP
1	A	162	VAL
1	B	30	LEU
1	B	36	GLN
1	B	46	ARG
1	B	80	ASN
1	B	81	ASN
1	B	93	LEU
1	B	94	ASP
1	B	143	GLU
1	B	177	LYS
1	B	193	LEU
1	B	210	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	80	ASN
1	A	81	ASN
1	A	85	HIS
1	A	111	ASN
1	A	182	GLN
1	A	190	GLN
1	B	36	GLN
1	B	80	ASN
1	B	81	ASN
1	B	85	HIS
1	B	111	ASN
1	B	182	GLN
1	B	190	GLN

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	199/224 (88%)	-0.38	2 (1%) 82 81	22, 34, 53, 60	0
1	B	199/224 (88%)	-0.34	4 (2%) 65 63	21, 33, 53, 63	0
All	All	398/448 (88%)	-0.36	6 (1%) 73 72	21, 33, 53, 63	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	62	GLU	3.7
1	A	62	GLU	3.2
1	A	113	CYS	3.0
1	B	113	CYS	2.5
1	B	91	TRP	2.1
1	B	26	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.