



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 6, 2023 – 10:48 PM EDT

PDB ID : 1KZG
Title : DbsCdc42(Y889F)
Authors : Rossman, K.L.; Worthylake, D.K.; Snyder, J.T.; Siderovski, D.P.; Campbell, S.L.; Sondek, J.
Deposited on : 2002-02-06
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	 6% (poor fit), 62% (0 outliers), 31% (1 outlier), 5% (2 outliers)
1	C	353	 11% (poor fit), 60% (0 outliers), 32% (1 outlier), 7% (2 outliers)
2	B	188	 6% (poor fit), 69% (0 outliers), 30% (1 outlier)
2	D	188	 6% (poor fit), 63% (0 outliers), 31% (1 outlier), 6% (2 outliers)

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8396 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GUANINE NUCLEOTIDE EXCHANGE FACTOR DBS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	337	2751	1743	474	513	21	0	0	0
1	C	330	2680	1702	457	500	21	0	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	622	MET	-	initiating methionine	UNP Q64096
A	696	ILE	ASN	SEE REMARK 999	UNP Q64096
A	697	PHE	ILE	SEE REMARK 999	UNP Q64096
A	698	LEU	PRO	SEE REMARK 999	UNP Q64096
A	699	ARG	ALA	SEE REMARK 999	UNP Q64096
A	700	GLU	GLY	SEE REMARK 999	UNP Q64096
A	701	LEU	VAL	SEE REMARK 999	UNP Q64096
A	889	PHE	TYR	engineered mutation	UNP Q64096
A	968	GLU	-	cloning artifact	UNP Q64096
A	969	HIS	-	expression tag	UNP Q64096
A	970	HIS	-	expression tag	UNP Q64096
A	971	HIS	-	expression tag	UNP Q64096
A	972	HIS	-	expression tag	UNP Q64096
A	973	HIS	-	expression tag	UNP Q64096
A	974	HIS	-	expression tag	UNP Q64096
C	1622	MET	-	initiating methionine	UNP Q64096
C	1696	ILE	ASN	SEE REMARK 999	UNP Q64096
C	1697	PHE	ILE	SEE REMARK 999	UNP Q64096
C	1698	LEU	PRO	SEE REMARK 999	UNP Q64096
C	1699	ARG	ALA	SEE REMARK 999	UNP Q64096
C	1700	GLU	GLY	SEE REMARK 999	UNP Q64096
C	1701	LEU	VAL	SEE REMARK 999	UNP Q64096
C	1889	PHE	TYR	engineered mutation	UNP Q64096
C	1968	GLU	-	cloning artifact	UNP Q64096
C	1969	HIS	-	expression tag	UNP Q64096

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1970	HIS	-	expression tag	UNP Q64096
C	1971	HIS	-	expression tag	UNP Q64096
C	1972	HIS	-	expression tag	UNP Q64096
C	1973	HIS	-	expression tag	UNP Q64096
C	1974	HIS	-	expression tag	UNP Q64096

- Molecule 2 is a protein called CDC42 HOMOLOG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	188	1471	943	239	282	7	0	0	0
2	D	177	1379	888	220	264	7	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	188	SER	CYS	engineered mutation	UNP P60953
D	1188	SER	CYS	engineered mutation	UNP P60953


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	39	Total 39	O 39	0	0
3	B	34	Total 34	O 34	0	0
3	C	24	Total 24	O 24	0	0
3	D	18	Total 18	O 18	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: GUANINE NUCLEOTIDE EXCHANGE FACTOR DBS




• Molecule 1: GUANINE NUCLEOTIDE EXCHANGE FACTOR DBS



- Molecule 2: CDC42 HOMOLOG



- Molecule 2: CDC42 HOMOLOG



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.32Å 88.19Å 232.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.60 31.16 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-2.60) 99.7 (31.16-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 2.61Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.217 , 0.259 0.219 , 0.261	Depositor DCC
R_{free} test set	2165 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	51.4	Xtrriage
Anisotropy	0.031	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8396	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2804	0.61	0/3767
1	C	0.36	0/2731	0.59	0/3672
2	B	0.42	0/1504	0.68	1/2044 (0.0%)
2	D	0.40	0/1409	0.66	0/1918
All	All	0.39	0/8448	0.63	1/11401 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	19	LEU	CA-CB-CG	5.33	127.55	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2751	0	2733	118	0
1	C	2680	0	2655	101	0
2	B	1471	0	1494	49	0
2	D	1379	0	1396	42	0
3	A	39	0	0	2	0
3	B	34	0	0	2	0
3	C	24	0	0	4	0
3	D	18	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8396	0	8278	302	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 302 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:719:MET:HE2	1:A:801:SER:HB3	1.26	1.13
1:A:685:ASN:HD22	1:A:685:ASN:H	1.21	0.88
1:C:1685:ASN:HD22	1:C:1685:ASN:H	1.25	0.85
1:C:1664:ASN:HD21	1:C:1666:LEU:HG	1.43	0.83
1:C:1691:HIS:CG	1:C:1695:ARG:HH12	1.95	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/353 (94%)	317 (95%)	16 (5%)	0	100	100
1	C	326/353 (92%)	308 (94%)	15 (5%)	3 (1%)	17	35
2	B	186/188 (99%)	171 (92%)	12 (6%)	3 (2%)	9	19
2	D	175/188 (93%)	158 (90%)	17 (10%)	0	100	100
All	All	1020/1082 (94%)	954 (94%)	60 (6%)	6 (1%)	25	47

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	180	PRO
1	C	1842	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	1884	GLU
1	C	1875	LYS
2	B	183	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/316 (95%)	284 (94%)	17 (6%)	21	42
1	C	291/316 (92%)	279 (96%)	12 (4%)	30	56
2	B	168/168 (100%)	164 (98%)	4 (2%)	49	74
2	D	157/168 (94%)	154 (98%)	3 (2%)	57	79
All	All	917/968 (95%)	881 (96%)	36 (4%)	32	58

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	1767	LYS
2	D	1177	LEU
1	C	1790	GLU
1	C	1893	GLN
1	A	855	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1731	ASN
1	C	1948	GLN
2	D	1134	GLN
2	D	1104	HIS
1	C	1893	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	337/353 (95%)	0.14	22 (6%) 18 14	28, 51, 102, 143	0
1	C	330/353 (93%)	0.56	40 (12%) 4 2	31, 58, 114, 138	0
2	B	188/188 (100%)	0.19	11 (5%) 22 17	29, 44, 90, 140	0
2	D	177/188 (94%)	0.26	12 (6%) 17 12	33, 50, 93, 106	0
All	All	1032/1082 (95%)	0.31	85 (8%) 11 8	28, 52, 101, 143	0

The worst 5 of 85 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	182	PRO	9.0
1	C	1853	LEU	8.7
1	A	880	GLY	7.7
1	C	1908	GLY	7.6
1	C	1854	ALA	7.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.