



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2021 – 10:01 AM EDT

PDB ID : 1MDP
Title : REFINED STRUCTURES OF TWO INSERTION(SLASH)DELETION MUTANTS PROBE FUNCTION OF THE MALTODEXTRIN BINDING PROTEIN
Authors : Sharff, A.J.; Quioco, F.A.
Deposited on : 1994-08-10
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	1	363	
1	2	363	
2	A	2	
2	B	2	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5851 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MALTODEXTRIN BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	1	363	2822	1817	459	540	6	0	0	0
1	2	363	2822	1817	459	540	6	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	134	ASP	LYS	engineered mutation	UNP P02928
1	135	PRO	LYS	engineered mutation	UNP P02928
1	?	-	ALA	deletion	UNP P02928
1	?	-	LEU	deletion	UNP P02928
1	?	-	LYS	deletion	UNP P02928
1	?	-	GLU	deletion	UNP P02928
1	?	-	LEU	deletion	UNP P02928
1	?	-	LYS	deletion	UNP P02928
1	?	-	ALA	deletion	UNP P02928
2	134	ASP	LYS	engineered mutation	UNP P02928
2	135	PRO	LYS	engineered mutation	UNP P02928
2	?	-	ALA	deletion	UNP P02928
2	?	-	LEU	deletion	UNP P02928
2	?	-	LYS	deletion	UNP P02928
2	?	-	GLU	deletion	UNP P02928
2	?	-	LEU	deletion	UNP P02928
2	?	-	LYS	deletion	UNP P02928
2	?	-	ALA	deletion	UNP P02928

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	A	2	Total	C	O	0	0	0
			23	12	11			
2	B	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	1	76	Total	O	0	0
			76	76		
3	2	85	Total	O	0	0
			85	85		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: MALTODEXTRIN BINDING PROTEIN

Chain 1: 



- Molecule 1: MALTODEXTRIN BINDING PROTEIN

Chain 2: 



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain A: 



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain B: 

GLC1
GLC2

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.51Å 88.33Å 75.79Å 90.00° 95.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ, X-PLOR	Depositor
R, R_{free}	0.187 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5851	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	1	0.74	0/2891	1.27	14/3925 (0.4%)
1	2	0.75	0/2891	1.33	14/3925 (0.4%)
All	All	0.75	0/5782	1.30	28/7850 (0.4%)

There are no bond length outliers.

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	2	316	ARG	CD-NE-CZ	15.50	145.30	123.60
1	2	316	ARG	NE-CZ-NH1	13.18	126.89	120.30
1	2	66	ARG	NE-CZ-NH1	8.06	124.33	120.30
1	1	341	TYR	CB-CG-CD1	7.63	125.58	121.00
1	2	98	ARG	NE-CZ-NH1	7.38	123.99	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	2822	0	2785	37	0
1	2	2822	0	2785	39	0
2	A	23	0	21	0	0
2	B	23	0	20	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	1	76	0	0	1	0
3	2	85	0	0	0	0
All	All	5851	0	5611	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 75 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:2:122:LEU:HD21	1:2:126:PRO:HD3	1.70	0.73
1:1:333:ILE:HD12	1:1:335:GLN:HE21	1.54	0.73
1:2:333:ILE:HD12	1:2:335:GLN:HE21	1.56	0.70
1:2:189:LYS:HG2	1:2:361:LEU:HD12	1.73	0.70
1:2:1:LYS:HE2	1:2:4:GLU:HG2	1.76	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	359/363 (99%)	338 (94%)	16 (4%)	5 (1%)	11	11
1	2	359/363 (99%)	335 (93%)	19 (5%)	5 (1%)	11	11
All	All	718/726 (99%)	673 (94%)	35 (5%)	10 (1%)	11	11

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1	2	ILE
1	1	3	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	2	2	ILE
1	2	100	ASN
1	2	173	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	291/292 (100%)	266 (91%)	25 (9%)	10	12
1	2	291/292 (100%)	268 (92%)	23 (8%)	12	15
All	All	582/584 (100%)	534 (92%)	48 (8%)	11	14

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	2	73	SER
1	2	202	LYS
1	2	115	LEU
1	2	147	LEU
1	2	222	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	1	335	GLN
1	2	218	ASN
1	2	365	GLN
1	2	335	GLN
1	1	218	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	A	1	2	12,12,12	1.10	1 (8%)	17,17,17	1.99	3 (17%)
2	GLC	A	2	2	11,11,12	0.49	0	15,15,17	0.74	0
2	GLC	B	1	2	12,12,12	1.21	1 (8%)	17,17,17	2.81	3 (17%)
2	GLC	B	2	2	11,11,12	0.52	0	15,15,17	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	1	2	-	2/2/22/22	0/1/1/1
2	GLC	A	2	2	-	1/2/19/22	0/1/1/1
2	GLC	B	1	2	-	1/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	GLC	O5-C1	-3.53	1.34	1.42
2	A	1	GLC	O5-C1	-3.47	1.34	1.42

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	O1-C1-O5	-7.05	89.23	110.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	C1-O5-C5	6.23	125.41	113.66
2	B	1	GLC	O5-C1-C2	5.65	120.37	110.28
2	A	1	GLC	C1-O5-C5	5.17	123.42	113.66
2	A	1	GLC	O1-C1-O5	-4.27	97.57	110.38

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1	GLC	C4-C5-C6-O6
2	A	1	GLC	O5-C5-C6-O6
2	B	1	GLC	O5-C5-C6-O6
2	A	2	GLC	C4-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	1	1
1	2	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	1	135:PRO	C	143:GLY	N	3.54
1	2	135:PRO	C	143:GLY	N	3.48

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.