wwPDB X-ray Structure Validation Summary Report (i)

Mar 4, 2024 – 03:35 AM EST

PDB ID	:	1PK8
Title	:	Crystal Structure of Rat Synapsin I C Domain Complexed to Ca.ATP
Authors	:	Brautigam, C.A.; Chelliah, Y.; Deisenhofer, J.
Deposited on	:	2003-06-05
Resolution	:	2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	2% 61% 11% 27%
1	B	422	% 58% 11% 31%
1	C	422	4% 54% 12% • 32%
1	D	422	5% 57% 14% 29%
1	E	422	2% 55% 12% 32%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	422	
1	G	422	
1	H	422	

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 19708 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called rat synapsin I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	25	0	0
			2414	1535	417	448	14			
1	B	292	Total	C	N	O	S	8	0	0
			2317	1476	397	430	14			
1	C	285	Total	C	N	O	S	18	0	0
			2272	1451	389	419	13			
1	D	300	Total	C	N	O	S	7	0	0
			2367	1506	406	441	14			
1	E	286	Total	C	N	O	S	40	0	0
			2279	1455	390	421	13			
1	F	286	Total	C	N	O	S	24	0	0
			2279	1455	390	421	13			
1	G	293	Total	C	N	O	S	5	0	0
			2326	1482	399	431	14			
1	H	286	Total	C	N	O	S	25	0	0
			2277	1454	390	420	13			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	cloning artifact	UNP P09951
A	1	SER	MET	cloning artifact	UNP P09951
B	0	GLY	-	cloning artifact	UNP P09951
B	1	SER	MET	cloning artifact	UNP P09951
C	0	GLY	-	cloning artifact	UNP P09951
C	1	SER	MET	cloning artifact	UNP P09951
D	0	GLY	-	cloning artifact	UNP P09951
D	1	SER	MET	cloning artifact	UNP P09951
E	0	GLY	-	cloning artifact	UNP P09951
E	1	SER	MET	cloning artifact	UNP P09951
F	0	GLY	-	cloning artifact	UNP P09951
F	1	SER	MET	cloning artifact	UNP P09951
G	0	GLY	-	cloning artifact	UNP P09951

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	1	SER	MET	cloning artifact	UNP P09951
H	0	GLY	-	cloning artifact	UNP P09951
H	1	SER	MET	cloning artifact	UNP P09951

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0
2	E	1	Total Ca 1 1	0	0
2	F	1	Total Ca 1 1	0	0
2	G	1	Total Ca 1 1	0	0
2	H	1	Total Ca 1 1	0	0

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	1
			62	20	10	26	6		
3	B	1	Total	C	N	O	P	0	1
			62	20	10	26	6		
3	C	1	Total	C	N	O	P	0	1
			62	20	10	26	6		
3	D	1	Total	C	N	O	P	0	1
			62	20	10	26	6		
3	E	1	Total	C	N	O	P	0	1
			62	20	10	26	6		
3	F	1	Total	C	N	O	P	0	1
			62	20	10	26	6		
3	G	1	Total	C	N	O	P	0	1
			62	20	10	26	6		
3	H	1	Total	C	N	O	P	0	1
			62	20	10	26	6		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	117	Total O 117 117	0	0
5	B	127	Total O 127 127	0	0
5	C	81	Total O 81 81	0	0
5	D	77	Total O 77 77	0	0
5	E	80	Total O 80 80	0	0
5	F	39	Total O 39 39	0	0
5	G	68	Total O 68 68	0	0
5	H	76	Total O 76 76	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: rat synapsin I

Chain A:



• Molecule 1: rat synapsin I

Chain B:



• Molecule 1: rat synapsin I

Chain C:



• Molecule 1: rat synapsin I



• Molecule 1: rat synapsin I



• Molecule 1: rat synapsin I

Chain F:



• Molecule 1: rat synapsin I

Chain G:



• Molecule 1: rat synapsin I

Chain H:




L259
T262
P265
K269
K279
T298
K299
T300
Y301
F307
I308
Y326
T329
SER
VAL
SER
GLY
ASN
W335
K336
THR
ASN
THR
GLY
SER
ALA
MET
L344
E345
Q346
T359
D368
L375
R381
I384
I385
E386
M392
I395
D400
V407
E408
L409

M412
Q416
A417
LEU
PRO
ARG
GLN

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	70.60Å 78.40Å 135.00Å 80.60° 76.90° 71.80°	Depositor
Resolution (Å)	20.00 – 2.10 37.16 – 2.09	Depositor EDS
% Data completeness (in resolution range)	91.9 (20.00-2.10) 91.5 (37.16-2.09)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) >$ [1]	3.25 (at 2.10Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.222 , 0.259 0.210 , 0.246	Depositor DCC
R_{free} test set	14970 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å^2)	27.7	Xtriage
Anisotropy	0.333	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 47.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	19708	wwPDB-VP
Average B, all atoms (Å^2)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ATP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.38	0/2466	0.62	0/3334
1	B	0.39	0/2367	0.64	0/3197
1	C	0.35	0/2321	0.62	0/3133
1	D	0.33	0/2418	0.61	0/3269
1	E	0.34	0/2328	0.60	0/3143
1	F	0.37	0/2328	0.61	0/3143
1	G	0.35	0/2375	0.62	0/3207
1	H	0.35	0/2326	0.60	0/3140
All	All	0.36	0/18929	0.62	0/25566

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2414	0	2397	36	0
1	B	2317	0	2289	33	0
1	C	2272	0	2246	34	0
1	D	2367	0	2341	39	0
1	E	2279	0	2253	33	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2279	0	2253	29	0
1	G	2326	0	2298	31	0
1	H	2277	0	2251	39	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	62	0	24	1	0
3	B	62	0	24	1	0
3	C	62	0	24	0	0
3	D	62	0	24	2	0
3	E	62	0	24	0	0
3	F	62	0	24	0	0
3	G	62	0	24	0	0
3	H	62	0	24	0	0
4	A	4	0	6	0	0
4	B	4	0	6	0	0
5	A	117	0	0	1	0
5	B	127	0	0	2	0
5	C	81	0	0	0	0
5	D	77	0	0	1	0
5	E	80	0	0	2	0
5	F	39	0	0	0	0
5	G	68	0	0	1	0
5	H	76	0	0	1	0
All	All	19708	0	18532	264	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 264 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:248:ASP:O	1:D:306:PRO:HD3	1.80	0.82
1:E:328:ARG:HD3	1:E:337:THR:HB	1.64	0.78
1:D:407:VAL:O	1:D:411:VAL:HG23	1.94	0.68
1:D:399:GLN:O	1:D:403:LYS:HG3	1.93	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:336:LYS:HE2	1:G:290:ASP:OD1	1.94	0.67

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/422 (72%)	297 (98%)	7 (2%)	0	100	100
1	B	286/422 (68%)	282 (99%)	4 (1%)	0	100	100
1	C	277/422 (66%)	272 (98%)	5 (2%)	0	100	100
1	D	296/422 (70%)	288 (97%)	7 (2%)	1 (0%)	41	41
1	E	278/422 (66%)	271 (98%)	7 (2%)	0	100	100
1	F	278/422 (66%)	270 (97%)	8 (3%)	0	100	100
1	G	285/422 (68%)	279 (98%)	6 (2%)	0	100	100
1	H	278/422 (66%)	271 (98%)	7 (2%)	0	100	100
All	All	2282/3376 (68%)	2230 (98%)	51 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	337	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/340 (77%)	260 (99%)	3 (1%)	73	79
1	B	252/340 (74%)	248 (98%)	4 (2%)	62	69
1	C	247/340 (73%)	238 (96%)	9 (4%)	35	36
1	D	258/340 (76%)	248 (96%)	10 (4%)	32	33
1	E	248/340 (73%)	241 (97%)	7 (3%)	43	47
1	F	248/340 (73%)	241 (97%)	7 (3%)	43	47
1	G	253/340 (74%)	243 (96%)	10 (4%)	31	32
1	H	247/340 (73%)	239 (97%)	8 (3%)	39	41
All	All	2016/2720 (74%)	1958 (97%)	58 (3%)	42	46

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	138	GLU
1	H	335	TRP
1	F	222	PHE
1	H	329	THR
1	G	400	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	399	GLN
1	G	338	ASN
1	H	383	HIS
1	G	334	ASN
1	D	195	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 26 ligands modelled in this entry, 8 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
3	ATP	A	800[A]	2	26,33,33	0.78	0	31,52,52	0.98	2 (6%)
3	ATP	B	801[B]	2	26,33,33	0.79	0	31,52,52	0.99	4 (12%)
3	ATP	A	800[B]	2	26,33,33	0.84	0	31,52,52	0.98	2 (6%)
3	ATP	D	803[A]	2	26,33,33	0.77	0	31,52,52	0.98	3 (9%)
3	ATP	E	804[A]	2	26,33,33	0.83	0	31,52,52	1.01	4 (12%)
4	EDO	A	818	-	3,3,3	0.91	0	2,2,2	0.41	0
3	ATP	E	804[B]	2	26,33,33	0.80	0	31,52,52	0.97	3 (9%)
3	ATP	B	801[A]	2	26,33,33	0.79	0	31,52,52	0.98	3 (9%)
3	ATP	H	807[B]	2	26,33,33	0.80	0	31,52,52	0.97	4 (12%)
3	ATP	C	802[A]	2	26,33,33	0.77	0	31,52,52	1.00	4 (12%)
3	ATP	C	802[B]	2	26,33,33	0.80	0	31,52,52	0.98	4 (12%)
3	ATP	G	806[B]	2	26,33,33	0.80	0	31,52,52	0.99	3 (9%)
3	ATP	H	807[A]	2	26,33,33	0.78	0	31,52,52	0.98	4 (12%)
4	EDO	B	819	-	3,3,3	0.63	0	2,2,2	0.43	0
3	ATP	F	805[B]	2	26,33,33	0.82	0	31,52,52	0.99	4 (12%)
3	ATP	G	806[A]	2	26,33,33	0.81	0	31,52,52	0.97	3 (9%)
3	ATP	F	805[A]	2	26,33,33	0.80	0	31,52,52	0.97	3 (9%)
3	ATP	D	803[B]	2	26,33,33	0.80	0	31,52,52	0.98	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	800[A]	2	-	0/18/38/38	0/3/3/3
3	ATP	B	801[B]	2	-	7/18/38/38	0/3/3/3
3	ATP	A	800[B]	2	-	6/18/38/38	0/3/3/3
3	ATP	D	803[A]	2	-	0/18/38/38	0/3/3/3
3	ATP	E	804[A]	2	-	0/18/38/38	0/3/3/3
4	EDO	A	818	-	-	0/1/1/1	-
3	ATP	E	804[B]	2	-	3/18/38/38	0/3/3/3
3	ATP	B	801[A]	2	-	0/18/38/38	0/3/3/3
3	ATP	H	807[B]	2	-	2/18/38/38	0/3/3/3
3	ATP	C	802[A]	2	-	2/18/38/38	0/3/3/3
3	ATP	C	802[B]	2	-	1/18/38/38	0/3/3/3
3	ATP	G	806[B]	2	-	5/18/38/38	0/3/3/3
3	ATP	H	807[A]	2	-	2/18/38/38	0/3/3/3
4	EDO	B	819	-	-	1/1/1/1	-
3	ATP	F	805[B]	2	-	3/18/38/38	0/3/3/3
3	ATP	G	806[A]	2	-	0/18/38/38	0/3/3/3
3	ATP	F	805[A]	2	-	0/18/38/38	0/3/3/3
3	ATP	D	803[B]	2	-	4/18/38/38	0/3/3/3

There are no bond length outliers.

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	800[A]	ATP	C5-C6-N6	2.55	124.23	120.35
3	E	804[A]	ATP	O3G-PG-O2G	-2.35	98.66	107.64
3	A	800[B]	ATP	C5-C6-N6	2.32	123.88	120.35
3	D	803[A]	ATP	C5-C6-N6	2.31	123.86	120.35
3	E	804[B]	ATP	C5-C6-N6	2.27	123.81	120.35

There are no chirality outliers.

5 of 36 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	800[B]	ATP	PB-O3B-PG-O3G
3	B	801[B]	ATP	PB-O3B-PG-O3G
3	C	802[A]	ATP	PB-O3B-PG-O2G
3	E	804[B]	ATP	PB-O3B-PG-O3G
3	F	805[B]	ATP	PB-O3B-PG-O3G

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	801[B]	ATP	1	0
3	A	800[B]	ATP	1	0
3	D	803[B]	ATP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ATP A 800 (A)



Ligand ATP B 801 (B)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP A 800 (B)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP D 803 (A)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP E 804 (A)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP E 804 (B)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP B 801 (A)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP H 807 (B)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP C 802 (A)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP C 802 (B)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP G 806 (B)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP H 807 (A)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP F 805 (B)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP G 806 (A)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP F 805 (A)



Bond lengths



Bond angles



Torsions



Rings

Ligand ATP D 803 (B)



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	306/422 (72%)	0.16	9 (2%)	51	57	11, 24, 47, 69	6 (1%)
1	B	292/422 (69%)	0.03	6 (2%)	63	68	12, 22, 43, 60	2 (0%)
1	C	285/422 (67%)	0.29	15 (5%)	26	32	17, 31, 56, 82	5 (1%)
1	D	300/422 (71%)	0.53	20 (6%)	17	22	19, 35, 59, 72	2 (0%)
1	E	286/422 (67%)	0.22	7 (2%)	59	64	20, 33, 56, 72	10 (3%)
1	F	286/422 (67%)	0.48	16 (5%)	24	29	19, 38, 66, 83	7 (2%)
1	G	293/422 (69%)	0.31	17 (5%)	23	28	15, 33, 58, 73	1 (0%)
1	H	286/422 (67%)	0.16	5 (1%)	70	74	15, 30, 50, 61	6 (2%)
All	All	2334/3376 (69%)	0.27	95 (4%)	37	43	11, 31, 58, 83	39 (1%)

The worst 5 of 95 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	174	VAL	7.3
1	A	168	LEU	7.0
1	D	167	VAL	6.6
1	A	172	VAL	6.3
1	C	139	ILE	5.9

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	EDO	A	818	4/4	0.79	0.18	20,22,26,26	0
2	CA	E	817	1/1	0.83	0.12	56,56,56,56	0
2	CA	F	817	1/1	0.84	0.15	48,48,48,48	0
3	ATP	C	802[B]	31/31	0.92	0.15	30,33,36,37	31
3	ATP	C	802[A]	31/31	0.92	0.15	21,27,36,37	31
3	ATP	E	804[A]	31/31	0.94	0.15	28,32,37,39	31
3	ATP	E	804[B]	31/31	0.94	0.15	33,35,38,40	31
3	ATP	F	805[A]	31/31	0.94	0.14	26,35,39,41	31
3	ATP	F	805[B]	31/31	0.94	0.14	33,35,37,37	31
3	ATP	H	807[A]	31/31	0.94	0.14	28,33,43,43	31
3	ATP	H	807[B]	31/31	0.94	0.14	28,31,34,36	31
2	CA	D	817	1/1	0.94	0.11	39,39,39,39	0
3	ATP	D	803[B]	31/31	0.95	0.16	27,29,33,33	31
2	CA	G	817	1/1	0.95	0.10	39,39,39,39	0
2	CA	H	817	1/1	0.95	0.10	38,38,38,38	0
3	ATP	D	803[A]	31/31	0.95	0.16	23,27,30,31	31
4	EDO	B	819	4/4	0.95	0.13	26,28,30,32	0
3	ATP	B	801[A]	31/31	0.96	0.13	15,20,23,23	31
3	ATP	B	801[B]	31/31	0.96	0.13	28,31,32,32	31
3	ATP	G	806[A]	31/31	0.96	0.12	23,26,30,31	31
3	ATP	G	806[B]	31/31	0.96	0.12	24,26,28,28	31
3	ATP	A	800[A]	31/31	0.97	0.15	11,19,21,22	31
3	ATP	A	800[B]	31/31	0.97	0.15	21,30,33,33	31
2	CA	C	817	1/1	0.98	0.09	44,44,44,44	0
2	CA	A	817	1/1	0.98	0.14	29,29,29,29	0
2	CA	B	817	1/1	0.98	0.11	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ATP C 802 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around ATP C 802 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around ATP E 804 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around ATP E 804 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around ATP F 805 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around ATP F 805 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around ATP H 807 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around ATP H 807 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around ATP D 803 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around ATP D 803 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around ATP B 801 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around ATP B 801 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around ATP G 806 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around ATP G 806 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around ATP A 800 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around ATP A 800 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.