



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 14, 2023 – 05:11 PM EDT

PDB ID : 1QZR
Title : CRYSTAL STRUCTURE OF THE ATPASE REGION OF SACCHAROMYCES CEREVISIAE TOPOISOMERASE II BOUND TO ICRF-187 (DEXRAZOXANE)
Authors : Classen, S.; Olland, S.; Berger, J.M.
Deposited on : 2003-09-17
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	 4% 80% 11% 9%
1	B	418	 6% 80% 9% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CDX	B	901[A]	X	-	-	-
4	CDX	B	901[B]	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA topoisomerase II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	382	Total	C	N	O	S	0	0	0
			3069	1954	520	580	15			
1	B	378	Total	C	N	O	S	0	0	0
			3041	1940	513	573	15			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	cloning artifact	UNP P06786
A	-3	HIS	-	cloning artifact	UNP P06786
A	-2	MET	-	cloning artifact	UNP P06786
A	-1	VAL	-	cloning artifact	UNP P06786
A	0	THR	-	cloning artifact	UNP P06786
B	-4	GLY	-	cloning artifact	UNP P06786
B	-3	HIS	-	cloning artifact	UNP P06786
B	-2	MET	-	cloning artifact	UNP P06786
B	-1	VAL	-	cloning artifact	UNP P06786
B	0	THR	-	cloning artifact	UNP P06786

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	419	Total 419	O 419	0	0
5	B	347	Total 347	O 347	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA topoisomerase II



- Molecule 1: DNA topoisomerase II



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.55Å 71.44Å 215.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.18 – 1.90 45.40 – 1.90	Depositor EDS
% Data completeness (in resolution range)	90.8 (45.18-1.90) 90.8 (45.40-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.196 , 0.237 0.194 , 0.236	Depositor DCC
R_{free} test set	5512 reflections (8.41%)	wwPDB-VP
Wilson B-factor (Å ²)	27.8	Xtrriage
Anisotropy	0.473	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 64.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6978	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, MG, CDX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/3124	0.66	5/4216 (0.1%)
1	B	0.46	0/3096	0.66	5/4179 (0.1%)
All	All	0.48	0/6220	0.66	10/8395 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	73	ASP	CB-CG-OD2	5.84	123.55	118.30
1	B	85	ASP	CB-CG-OD2	5.56	123.30	118.30
1	A	43	ASP	CB-CG-OD2	5.38	123.15	118.30
1	A	65	ASP	CB-CG-OD2	5.33	123.09	118.30
1	B	296	ASP	CB-CG-OD2	5.32	123.08	118.30
1	B	26	ASP	CB-CG-OD2	5.29	123.06	118.30
1	A	47	ASP	CB-CG-OD2	5.21	122.99	118.30
1	B	405	ASP	CB-CG-OD2	5.07	122.87	118.30
1	A	133	ASP	CB-CG-OD2	5.07	122.86	118.30
1	B	162	ASP	CB-CG-OD2	5.06	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3069	0	3059	22	0
1	B	3041	0	3035	20	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	0	13	0	0
3	B	31	0	13	0	0
4	B	38	0	32	8	0
5	A	419	0	0	0	0
5	B	347	0	0	4	0
All	All	6978	0	6152	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:149:CYS:SG	5:B:1226:HOH:O	2.28	0.85
1:A:245:ARG:H	1:A:249:ASN:HD22	1.30	0.79
1:B:91:GLU:H	1:B:91:GLU:CD	1.94	0.69
1:A:134:GLU:HG3	1:A:136:LYS:HE2	1.83	0.59
1:A:17:GLN:NE2	1:A:144:TYR:OH	2.38	0.56
4:B:901[B]:CDX:HAQ2	4:B:901[B]:CDX:HAE2	1.88	0.55
1:A:131:ASP:O	1:A:134:GLU:HG3	2.06	0.55
1:A:94:THR:HG22	1:A:200:LYS:HE2	1.89	0.54
1:A:121:PHE:CD2	1:A:158:LEU:HD22	2.43	0.53
1:B:392:MET:HA	1:B:392:MET:CE	2.38	0.52
1:B:251:VAL:CG1	1:B:279:ILE:HG21	2.39	0.52
4:B:901[B]:CDX:HAE2	4:B:901[B]:CDX:CAQ	2.39	0.52
1:A:323:GLN:HE22	1:A:383:ILE:H	1.58	0.51
1:B:11:LYS:HE3	1:B:12:TYR:CE2	2.46	0.50
1:B:222:ARG:O	1:B:225:ASP:HB2	2.11	0.49
1:B:63:ILE:O	5:B:1226:HOH:O	2.20	0.49
1:A:223:VAL:HG12	1:A:242:LEU:HD11	1.94	0.48
1:B:210:GLU:HG3	5:B:1162:HOH:O	2.14	0.48
1:A:220:ARG:HG2	1:A:237:LEU:HD21	1.95	0.47
1:B:291:ALA:HB3	1:B:352:PHE:HB2	1.97	0.46
1:B:284:ILE:HG13	1:B:381:CYS:SG	2.55	0.46
1:A:280:LEU:HG	1:A:387:TYR:CE1	2.51	0.46
1:A:96:GLU:HG2	1:A:198:THR:HG23	1.98	0.45
1:A:323:GLN:NE2	1:A:383:ILE:H	2.13	0.45
1:A:323:GLN:HE22	1:A:382:GLU:HA	1.81	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:301:GLN:HE21	1:B:301:GLN:HB2	1.61	0.45
1:A:291:ALA:HB3	1:A:352:PHE:HB2	1.98	0.45
4:B:901[B]:CDX:HAF1	4:B:901[B]:CDX:HAK1	1.72	0.45
1:A:341:LYS:HD2	1:A:343:PHE:CE2	2.52	0.44
4:B:901[B]:CDX:HAK2	5:B:1252:HOH:O	2.17	0.44
1:A:317:VAL:O	1:A:321:THR:HG23	2.18	0.43
4:B:901[A]:CDX:HAF1	4:B:901[A]:CDX:HAK1	1.69	0.43
1:B:251:VAL:HG12	1:B:279:ILE:HG21	2.00	0.42
1:A:94:THR:CG2	1:A:200:LYS:HE2	2.49	0.42
1:A:231:ARG:HD3	1:A:283:ARG:NH2	2.35	0.42
1:B:144:TYR:CZ	4:B:901[A]:CDX:HAQ1	2.55	0.41
1:A:308:ILE:HD13	1:A:367:LYS:HD3	2.01	0.41
1:B:251:VAL:HG11	1:B:279:ILE:HG21	2.02	0.41
1:A:222:ARG:O	1:A:225:ASP:HB2	2.21	0.41
1:A:65:ASP:OD1	1:A:222:ARG:NE	2.49	0.41
1:B:91:GLU:CD	1:B:91:GLU:N	2.70	0.41
1:B:117:PRO:HB3	1:B:158:LEU:CD2	2.50	0.40
1:A:27:THR:OG1	4:B:901[B]:CDX:HAE1	2.22	0.40
1:B:141:ARG:HD3	1:B:360:PRO:HD2	2.03	0.40
1:B:175:ASN:N	1:B:175:ASN:HD22	2.20	0.40
1:B:280:LEU:O	1:B:291:ALA:HA	2.21	0.40
1:B:82:LYS:HD2	1:B:82:LYS:HA	1.97	0.40
4:B:901[A]:CDX:CAQ	4:B:901[A]:CDX:HAE2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/418 (90%)	368 (98%)	8 (2%)	0	100	100
1	B	372/418 (89%)	362 (97%)	9 (2%)	1 (0%)	41	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	748/836 (90%)	730 (98%)	17 (2%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	190	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/381 (90%)	335 (98%)	7 (2%)	55	51
1	B	339/381 (89%)	322 (95%)	17 (5%)	24	15
All	All	681/762 (89%)	657 (96%)	24 (4%)	36	27

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ASP
1	A	77	ARG
1	A	221	ARG
1	A	227	ASN
1	A	334	LYS
1	A	349	MET
1	A	350	PHE
1	B	52	LYS
1	B	83	ARG
1	B	91	GLU
1	B	111	LYS
1	B	162	ASP
1	B	163	LEU
1	B	164	ASN
1	B	175	ASN
1	B	205	ARG
1	B	227	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	257	SER
1	B	301	GLN
1	B	350	PHE
1	B	351	ILE
1	B	380	ARG
1	B	382	GLU
1	B	402	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	35	GLN
1	A	93	HIS
1	A	129	ASN
1	A	249	ASN
1	A	323	GLN
1	A	389	ASN
1	B	175	ASN
1	B	234	ASN
1	B	249	ASN
1	B	301	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ANP	B	903	2	29,33,33	1.80	8 (27%)	31,52,52	1.27	4 (12%)
4	CDX	B	901[A]	-	19,20,20	0.94	0	21,28,28	2.14	7 (33%)
4	CDX	B	901[B]	-	19,20,20	0.89	0	21,28,28	2.10	6 (28%)
3	ANP	A	902	2	29,33,33	1.65	8 (27%)	31,52,52	1.26	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	902	2	-	4/14/38/38	0/3/3/3
4	CDX	B	901[A]	-	1/1/7/9	5/8/32/32	0/2/2/2
3	ANP	B	903	2	-	4/14/38/38	0/3/3/3
4	CDX	B	901[B]	-	1/1/7/9	3/8/32/32	0/2/2/2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	903	ANP	PB-O3A	-5.58	1.52	1.59
3	A	902	ANP	PB-O3A	-4.19	1.53	1.59
3	B	903	ANP	PB-O1B	3.43	1.51	1.46
3	A	902	ANP	PG-O2G	-2.68	1.49	1.56
3	A	902	ANP	PG-O1G	2.63	1.50	1.46
3	A	902	ANP	PB-O2B	-2.57	1.49	1.56
3	A	902	ANP	C2-N3	2.42	1.36	1.32
3	A	902	ANP	PG-O3G	-2.34	1.50	1.56
3	B	903	ANP	PB-O2B	-2.34	1.50	1.56
3	B	903	ANP	PG-O2G	-2.28	1.50	1.56
3	B	903	ANP	C5-C4	2.27	1.46	1.40
3	B	903	ANP	C2-N3	2.20	1.35	1.32
3	B	903	ANP	PG-O3G	-2.18	1.50	1.56
3	A	902	ANP	C5-C4	2.07	1.46	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	902	ANP	PB-O1B	2.02	1.49	1.46
3	B	903	ANP	PG-O1G	2.02	1.49	1.46

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	901[A]	CDX	CAC-CAD-NAJ	-4.56	106.13	112.58
4	B	901[B]	CDX	CAC-CAD-NAJ	-4.50	106.21	112.58
4	B	901[B]	CDX	CAP-CAQ-NAJ	-4.14	106.73	112.58
4	B	901[A]	CDX	CAE-CAL-NAM	-4.03	107.05	114.31
4	B	901[A]	CDX	CAP-CAQ-NAJ	-3.95	107.00	112.58
4	B	901[B]	CDX	CAE-CAL-NAM	-3.94	107.21	114.31
4	B	901[A]	CDX	CAL-CAK-NAJ	-3.85	108.37	113.56
3	B	903	ANP	N3-C2-N1	-3.71	122.88	128.68
3	A	902	ANP	N3-C2-N1	-3.61	123.03	128.68
4	B	901[B]	CDX	CAQ-NAJ-CAD	-2.65	106.87	109.90
4	B	901[B]	CDX	CAL-CAK-NAJ	-2.57	110.10	113.56
4	B	901[B]	CDX	CAF-NAM-CAR	-2.42	107.24	109.89
3	B	903	ANP	C3'-C2'-C1'	2.32	104.48	100.98
4	B	901[A]	CDX	CAF-NAM-CAR	-2.32	107.35	109.89
4	B	901[A]	CDX	CAQ-NAJ-CAD	-2.10	107.51	109.90
3	B	903	ANP	C4-C5-N7	-2.08	107.23	109.40
3	B	903	ANP	O2B-PB-O3A	2.06	111.52	104.64
4	B	901[A]	CDX	OAO-CAP-NAI	2.04	123.52	120.28

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	901[A]	CDX	CAL
4	B	901[B]	CDX	CAL

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	902	ANP	PB-N3B-PG-O1G
3	A	902	ANP	PG-N3B-PB-O1B
3	A	902	ANP	PA-O3A-PB-O1B
3	A	902	ANP	PA-O3A-PB-O2B
3	B	903	ANP	PB-N3B-PG-O1G
3	B	903	ANP	PG-N3B-PB-O1B
3	B	903	ANP	PA-O3A-PB-O1B
3	B	903	ANP	PA-O3A-PB-O2B

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	B	901[A]	CDX	NAJ-CAK-CAL-CAE
4	B	901[A]	CDX	CAK-CAL-NAM-CAF
4	B	901[B]	CDX	NAJ-CAK-CAL-CAE
4	B	901[B]	CDX	CAK-CAL-NAM-CAF
4	B	901[A]	CDX	CAE-CAL-NAM-CAF
4	B	901[A]	CDX	CAK-CAL-NAM-CAR
4	B	901[A]	CDX	CAE-CAL-NAM-CAR
4	B	901[B]	CDX	CAE-CAL-NAM-CAF


There are no ring outliers.


2 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	901[A]	CDX	3	0
4	B	901[B]	CDX	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	382/418 (91%)	0.24	16 (4%) 36 39	13, 20, 32, 79	0
1	B	378/418 (90%)	0.32	23 (6%) 21 24	14, 22, 31, 52	0
All	All	760/836 (90%)	0.28	39 (5%) 28 31	13, 21, 31, 79	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	104	ILE	4.9
1	B	165	VAL	4.5
1	B	405	ASP	4.2
1	B	186	THR	4.1
1	B	162	ASP	3.9
1	A	47	ASP	3.7
1	B	191	GLY	3.7
1	A	276	ILE	3.7
1	B	189	LYS	3.6
1	A	333	LYS	3.6
1	A	385	LEU	3.4
1	B	164	ASN	3.3
1	B	187	SER	3.0
1	A	402	GLU	3.0
1	A	255	LEU	2.9
1	B	161	ALA	2.8
1	A	409	GLU	2.8
1	B	403	ILE	2.7
1	A	210	GLU	2.7
1	B	106	ILE	2.7
1	A	407	ASN	2.6
1	A	44	GLU	2.6
1	B	113	ASN	2.5
1	A	334	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	29	ILE	2.5
1	B	398	THR	2.5
1	B	296	ASP	2.4
1	A	408	GLU	2.4
1	B	276	ILE	2.4
1	B	190	LYS	2.3
1	B	108	ILE	2.3
1	B	185	ILE	2.3
1	B	209	LYS	2.2
1	B	395	ASP	2.2
1	A	113	ASN	2.2
1	A	280	LEU	2.1
1	B	340	VAL	2.1
1	B	167	GLN	2.0
1	A	93	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CDX	B	901[A]	19/19	0.87	0.25	54,55,57,58	19
4	CDX	B	901[B]	19/19	0.87	0.25	54,56,57,58	19
2	MG	B	905	1/1	0.94	0.10	25,25,25,25	0
3	ANP	B	903	31/31	0.97	0.11	24,30,38,39	0
3	ANP	A	902	31/31	0.98	0.14	20,23,26,29	0
2	MG	A	904	1/1	0.98	0.12	23,23,23,23	0




The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around CDX B 901 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ANP B 903:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.