



wwPDB X-ray Structure Validation Summary Report

Feb 4, 2024 – 03:58 PM EST


PDB ID : 1R27
Title : Crystal Structure of NarGH complex
Authors : Jormakka, M.; Richardson, D.; Byrne, B.; Iwata, S.
Deposited on : 2003-09-26
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	1246	
1	C	1246	
2	B	512	
2	D	512	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SF4	B	5800	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 29331 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Respiratory nitrate reductase 1 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1218	9660	6100	1691	1821	48	0	0	0
1	C	1218	9660	6100	1691	1821	48	0	0	0


- Molecule 2 is a protein called Respiratory nitrate reductase 1 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	465	3711	2354	640	687	30	0	0	0
2	D	465	3711	2354	640	687	30	0	0	0

- Molecule 3 is MOLYBDENUM ATOM (three-letter code: MO) (formula: Mo).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mo	0	0
			1	1		
3	C	1	Total	Mo	0	0
			1	1		

- Molecule 4 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Fe S	0	0
			8	4 4		
4	B	1	Total	Fe S	0	0
			8	4 4		
4	B	1	Total	Fe S	0	0
			8	4 4		
4	B	1	Total	Fe S	0	0
			8	4 4		
4	C	1	Total	Fe S	0	0
			8	4 4		
4	D	1	Total	Fe S	0	0
			8	4 4		
4	D	1	Total	Fe S	0	0
			8	4 4		

- Molecule 5 is 2-AMINO-5,6-DIMERCAPTO-7-METHYL-3,7,8A,9-TETRAHYDRO-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-4-ONE GUANOSINE DINUCLEOTIDE (three-letter code: MGD) (formula: C₂₀H₂₆N₁₀O₁₃P₂S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
5	A	1	Total	C	N	O	P	S	0	0
			47	20	10	13	2	2		
5	A	1	Total	C	N	O	P	S	0	0
			47	20	10	13	2	2		
5	C	1	Total	C	N	O	P	S	0	0
			47	20	10	13	2	2		
5	C	1	Total	C	N	O	P	S	0	0
			47	20	10	13	2	2		

- Molecule 6 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe₃S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	Fe	S	0	0
			7	3	4		
6	D	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	815	Total	O	0	0
			815	815		
7	B	376	Total	O	0	0
			376	376		
7	C	784	Total	O	0	0
			784	784		
7	D	346	Total	O	0	0
			346	346		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: Respiratory nitrate reductase 1 alpha chain





- Molecule 1: Respiratory nitrate reductase 1 alpha chain






• Molecule 2: Respiratory nitrate reductase 1 beta chain



• Molecule 2: Respiratory nitrate reductase 1 beta chain



GLY
SER
ASP
THR
LYS
PHE
ASN
LEU
PHE
ASN
SER
SER
ARG
ARG
ILE
ASP
ALA
ILE
ASP
VAL
THR
SER
LYS
THR
GLU
PRO
HIS
PRO

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	127.40Å 298.30Å 296.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.86 – 2.00	Depositor
% Data completeness (in resolution range)	88.2 (39.86-2.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.204 , 0.230	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	29331	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: F3S, SF4, MO, MGD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/9914	0.62	2/13462 (0.0%)
1	C	0.32	0/9914	0.61	2/13462 (0.0%)
2	B	0.32	0/3800	0.61	1/5145 (0.0%)
2	D	0.32	0/3800	0.60	0/5145
All	All	0.32	0/27428	0.61	5/37214 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	809	LEU	N-CA-C	-5.66	95.73	111.00
2	B	11	LEU	CA-CB-CG	5.41	127.75	115.30
1	A	1097	ILE	N-CA-C	-5.38	96.47	111.00
1	C	990	GLU	N-CA-C	-5.36	96.54	111.00
1	C	809	LEU	N-CA-C	-5.13	97.15	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9660	0	9330	239	0
1	C	9660	0	9330	227	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	3711	0	3659	57	0
2	D	3711	0	3659	74	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
4	A	8	0	0	0	0
4	B	24	0	0	2	0
4	C	8	0	0	0	0
4	D	24	0	0	1	0
5	A	94	0	44	6	0
5	C	94	0	44	7	0
6	B	7	0	0	0	0
6	D	7	0	0	0	0
7	A	815	0	0	39	0
7	B	376	0	0	8	0
7	C	784	0	0	26	0
7	D	346	0	0	9	0
All	All	29331	0	26066	577	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 577 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1098:HIS:O	1:A:1164:ALA:HB1	1.50	1.11
1:A:454:LYS:HD3	1:A:487:LEU:HB3	1.40	1.00
1:C:990:GLU:O	1:C:991:ASP:HB2	1.66	0.96
1:C:106:ASN:HD22	2:D:112:HIS:HE1	1.17	0.93
1:A:1164:ALA:HA	7:A:6567:HOH:O	1.69	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1216/1246 (98%)	1150 (95%)	59 (5%)	7 (1%)	25	19
1	C	1216/1246 (98%)	1144 (94%)	62 (5%)	10 (1%)	19	13
2	B	463/512 (90%)	450 (97%)	10 (2%)	3 (1%)	25	19
2	D	463/512 (90%)	448 (97%)	14 (3%)	1 (0%)	47	44
All	All	3358/3516 (96%)	3192 (95%)	145 (4%)	21 (1%)	25	19

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1164	ALA
1	A	1244	SER
1	C	748	GLY
1	A	578	VAL
1	A	719	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1019/1043 (98%)	996 (98%)	23 (2%)	50	53
1	C	1019/1043 (98%)	995 (98%)	24 (2%)	49	51
2	B	400/439 (91%)	391 (98%)	9 (2%)	50	53
2	D	400/439 (91%)	389 (97%)	11 (3%)	43	44
All	All	2838/2964 (96%)	2771 (98%)	67 (2%)	49	51

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	27	LYS
2	D	58	ASN
2	D	307	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	37	GLU
2	B	11	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 77 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	862	GLN
2	D	112	HIS
1	C	921	GLN
1	C	1165	GLN
2	D	165	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SF4	B	5806	2	0,12,12	-	-	-		
4	SF4	D	7800	2	0,12,12	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MGD	C	7801	3	41,52,52	2.35	12 (29%)	40,81,81	1.60	4 (10%)
5	MGD	A	5802	3	41,52,52	1.90	7 (17%)	40,81,81	2.20	4 (10%)
4	SF4	D	7806	2	0,12,12	-	-	-	-	-
4	SF4	B	5800	2	0,12,12	-	-	-	-	-
4	SF4	B	5805	2	0,12,12	-	-	-	-	-
5	MGD	A	5801	3	41,52,52	2.55	13 (31%)	40,81,81	1.71	5 (12%)
6	F3S	D	7808	2	0,9,9	-	-	-	-	-
6	F3S	B	5808	2	0,9,9	-	-	-	-	-
5	MGD	C	7802	3	41,52,52	1.93	9 (21%)	40,81,81	2.31	4 (10%)
4	SF4	A	5807	1	0,12,12	-	-	-	-	-
4	SF4	D	7805	2	0,12,12	-	-	-	-	-
4	SF4	C	7807	1	0,12,12	-	-	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	B	5806	2	-	-	0/6/5/5
4	SF4	D	7800	2	-	-	0/6/5/5
5	MGD	C	7801	3	-	6/18/66/66	0/6/6/6
5	MGD	A	5802	3	-	5/18/66/66	0/6/6/6
4	SF4	D	7806	2	-	-	0/6/5/5
4	SF4	B	5800	2	-	-	0/6/5/5
4	SF4	B	5805	2	-	-	0/6/5/5
4	SF4	D	7805	2	-	-	0/6/5/5
6	F3S	D	7808	2	-	-	0/3/3/3
6	F3S	B	5808	2	-	-	0/3/3/3
5	MGD	C	7802	3	-	5/18/66/66	0/6/6/6
4	SF4	A	5807	1	-	-	0/6/5/5
5	MGD	A	5801	3	-	6/18/66/66	0/6/6/6
4	SF4	C	7807	1	-	-	0/6/5/5

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	7802	MGD	C14-N15	7.73	1.55	1.46
5	C	7801	MGD	O11-C11	7.26	1.53	1.43
5	A	5801	MGD	O11-C11	7.16	1.53	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	5801	MGD	C21-N22	7.01	1.43	1.35
5	A	5802	MGD	C14-N15	6.85	1.54	1.46

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	7802	MGD	O11-C23-C14	-11.57	101.25	108.96
5	A	5802	MGD	O11-C23-C14	-10.98	101.64	108.96
5	A	5801	MGD	O11-C23-N22	6.30	115.04	108.57
5	C	7801	MGD	O11-C23-N22	5.75	114.48	108.57
5	C	7802	MGD	O11-C23-N22	-5.40	103.02	108.57

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	5801	MGD	C5'-O5'-PB-O1B
5	A	5802	MGD	C5'-O5'-PB-O1B
5	C	7801	MGD	C5'-O5'-PB-O1B
5	C	7802	MGD	PA-O3B-PB-O5'
5	C	7802	MGD	C5'-O5'-PB-O1B


There are no ring outliers.


6 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	7800	SF4	1	0
5	C	7801	MGD	3	0
5	A	5802	MGD	5	0
4	B	5800	SF4	2	0
5	A	5801	MGD	1	0
5	C	7802	MGD	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.