



Full wwPDB X-ray Structure Validation Report i

Oct 17, 2021 – 01:21 AM EDT

PDB ID : 1R6F

Title : The structure of Yersinia pestis V-antigen, an essential virulence factor and mediator of immunity against plague

Authors : Derewenda, U.; Mateja, A.; Devedjiev, Y.; Routzahn, K.M.; Evdokimov, A.G.; Derewenda, Z.S.; Waugh, D.S.

Deposited on : 2003-10-15

Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Xtriage (Phenix) : 1.13

EDS : 2.23.2

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.



2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 2139 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Virulence-associated V antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	272	2139	1343	365	425	6	0	0	1


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	GLY	-	cloning artifact	UNP P21206
A	40	ALA	LYS	engineered mutation	UNP P21206
A	41	ALA	ASP	engineered mutation	UNP P21206
A	42	ALA	LYS	engineered mutation	UNP P21206
A	327	HIS	-	expression tag	UNP P21206
A	328	HIS	-	expression tag	UNP P21206
A	329	HIS	-	expression tag	UNP P21206
A	330	HIS	-	expression tag	UNP P21206
A	331	HIS	-	expression tag	UNP P21206
A	332	HIS	-	expression tag	UNP P21206

3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Virulence-associated V antigen



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	35.88Å 45.09Å 46.91Å 76.20° 78.42° 77.17°	Depositor
Resolution (Å)	18.50 – 2.17 17.64 – 2.16	Depositor EDS
% Data completeness (in resolution range)	96.2 (18.50-2.17) 95.2 (17.64-2.16)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$< I/\sigma(I) >$ ¹	4.42 (at 2.15Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R , R_{free}	0.217 , 0.284 0.266 , 0.264	Depositor DCC
R_{free} test set	1383 reflections (9.80%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtriage
Anisotropy	0.572	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 29.6	EDS
L-test for twinning ²	$< L > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.001 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2139	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2163	1.02	11/2909 (0.4%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	282	ASP	CB-CG-OD2	6.75	124.37	118.30
1	A	228	GLY	N-CA-C	6.64	129.70	113.10
1	A	45	ASP	CB-CG-OD2	6.58	124.22	118.30
1	A	275	ASP	CB-CG-OD2	6.24	123.92	118.30
1	A	67	ASP	CB-CG-OD2	6.00	123.70	118.30
1	A	132	ASP	CB-CG-OD2	5.83	123.55	118.30
1	A	278	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	A	262	ASP	CB-CG-OD2	5.38	123.15	118.30
1	A	227	ASP	CB-CG-OD2	5.23	123.01	118.30
1	A	203	ASP	CB-CG-OD2	5.22	122.99	118.30
1	A	195	ASP	CB-CG-OD2	5.09	122.88	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2139	0	2141	58	0
All	All	2139	0	2141	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:PHE:HD2	1:A:119:MET:HE2	1.42	0.85
1:A:284:VAL:O	1:A:288:THR:HG23	1.76	0.84
1:A:153:LEU:HG	1:A:302:ILE:CD1	2.09	0.82
1:A:118:PHE:HD2	1:A:119:MET:CE	1.94	0.81
1:A:278:ARG:HG2	1:A:278:ARG:HH11	1.48	0.78
1:A:173:GLU:HG2	1:A:191:ILE:HG12	1.69	0.73
1:A:153:LEU:HG	1:A:302:ILE:HD13	1.72	0.72
1:A:35:LEU:O	1:A:38:LEU:HB2	1.94	0.67
1:A:118:PHE:CD2	1:A:119:MET:CE	2.78	0.66
1:A:174:ILE:HA	1:A:184:ILE:HD13	1.78	0.66
1:A:254:LYS:HE2	1:A:257:TYR:CE2	2.31	0.65
1:A:322:ASP:OD1	1:A:322:ASP:N	2.27	0.65
1:A:78:PHE:CZ	1:A:123:HIS:HB2	2.36	0.60
1:A:153:LEU:CG	1:A:302:ILE:HD13	2.31	0.60
1:A:186:ILE:HG13	1:A:237:LYS:HB2	1.85	0.59
1:A:318:ARG:CG	1:A:318:ARG:HH11	2.16	0.59
1:A:318:ARG:HG3	1:A:318:ARG:NH1	2.17	0.58
1:A:319:LEU:O	1:A:319:LEU:HG	2.04	0.58
1:A:78:PHE:HZ	1:A:123:HIS:HB2	1.70	0.57
1:A:47:SER:HB3	1:A:63:VAL:HA	1.87	0.56
1:A:135:ILE:HG21	1:A:301:ALA:CB	2.36	0.55
1:A:187:HIS:CG	1:A:188:ASP:H	2.25	0.54
1:A:318:ARG:HH11	1:A:318:ARG:HG3	1.73	0.54
1:A:222:THR:HG23	1:A:238:ASP:OD1	2.08	0.53
1:A:153:LEU:CD2	1:A:302:ILE:HD13	2.38	0.53
1:A:118:PHE:CD2	1:A:119:MET:HE1	2.42	0.53
1:A:221:GLN:NE2	1:A:232:LYS:HB3	2.24	0.52
1:A:121:VAL:HG22	1:A:140:VAL:HG21	1.92	0.52
1:A:121:VAL:CG2	1:A:140:VAL:HG21	2.39	0.51
1:A:163:GLU:OE2	1:A:247:THR:OG1	2.23	0.51
1:A:163:GLU:HB3	1:A:291:LEU:HD21	1.93	0.51
1:A:143:MET:HE1	1:A:308:PHE:HE2	1.76	0.50
1:A:254:LYS:HG3	1:A:257:TYR:CZ	2.46	0.50
1:A:47:SER:HB2	1:A:62:ARG:O	2.13	0.49
1:A:104:PHE:CZ	1:A:136:LEU:HB3	2.48	0.48
1:A:118:PHE:CZ	1:A:122:MET:HE2	2.47	0.48
1:A:318:ARG:CG	1:A:318:ARG:NH1	2.74	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:HIS:CG	1:A:188:ASP:N	2.81	0.47
1:A:152:LYS:NZ	1:A:155:GLU:OE2	2.39	0.47
1:A:226:VAL:HB	1:A:227:ASP:H	1.57	0.46
1:A:118:PHE:CD2	1:A:119:MET:HE2	2.34	0.46
1:A:135:ILE:HG21	1:A:301:ALA:HB2	1.99	0.44
1:A:116:ARG:HG3	1:A:312:TYR:CE1	2.52	0.44
1:A:117:ALA:HB2	1:A:143:MET:HB3	2.00	0.43
1:A:132:ASP:HB2	1:A:297:ARG:NH1	2.34	0.43
1:A:221:GLN:HE22	1:A:232:LYS:HB3	1.81	0.43
1:A:29:SER:O	1:A:33:GLU:HG2	2.18	0.43
1:A:254:LYS:HE2	1:A:257:TYR:CD2	2.53	0.43
1:A:118:PHE:CZ	1:A:122:MET:CE	3.02	0.42
1:A:174:ILE:O	1:A:178:LEU:HG	2.20	0.42
1:A:77:TYR:HE2	1:A:319:LEU:HD13	1.85	0.41
1:A:227:ASP:OD1	1:A:228:GLY:N	2.51	0.41
1:A:254:LYS:HG3	1:A:257:TYR:CE1	2.55	0.41
1:A:254:LYS:HD2	1:A:256:SER:O	2.21	0.41
1:A:39:VAL:O	1:A:42:ALA:O	2.39	0.41
1:A:177:HIS:CD2	1:A:184:ILE:HD12	2.56	0.41
1:A:278:ARG:HG2	1:A:282:ASP:OD2	2.22	0.40
1:A:116:ARG:HG3	1:A:312:TYR:CD1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	264/310 (85%)	249 (94%)	12 (4%)	3 (1%)	14 11

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	228	GLY
1	A	260	ASN
1	A	226	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	238/277 (86%)	200 (84%)	38 (16%)	2 1

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	SER
1	A	32	LEU
1	A	37	GLN
1	A	44	ILE
1	A	63	VAL
1	A	65	THR
1	A	68	ILE
1	A	70	LEU
1	A	86	LYS
1	A	89	HIS
1	A	100	ARG
1	A	145	HIS
1	A	156	GLU
1	A	165	LYS
1	A	184	ILE
1	A	191	ILE
1	A	196	LYS
1	A	197	ASN
1	A	217	GLU
1	A	226	VAL
1	A	231	LYS
1	A	253	LEU
1	A	256	SER
1	A	258	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	259	TYR
1	A	260	ASN
1	A	275	ASP
1	A	278	ARG
1	A	288	THR
1	A	290	GLN
1	A	299	ASN
1	A	303	GLU
1	A	305	LEU
1	A	316	MET
1	A	317	GLN
1	A	318	ARG
1	A	321	ASP
1	A	322	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	197	ASN
1	A	221	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.