



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 4, 2024 – 01:01 PM EST

PDB ID : 1RQJ  
Title : Active Conformation of Farnesyl Pyrophosphate Synthase Bound to Isopentyl Pyrophosphate and Risedronate  
Authors : Hosfield, D.J.; Zhang, Y.; Dougan, D.R.; Brooun, A.; Tari, L.W.; Swanson, R.V.; Finn, J.  
Deposited on : 2003-12-05  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	299	
1	B	299	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5022 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Geranyltranstransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	299	Total 2263	C 1409	N 404	O 438	S 12	0	2	0
1	B	298	Total 2278	C 1420	N 408	O 439	S 11	0	9	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total 3	Mg 3	0	0
2	B	3	Total 3	Mg 3	0	0

- Molecule 3 is 3-METHYLBUT-3-ENYL TRIHYDROGEN DIPHOSPHATE (three-letter code: IPE) (formula: C<sub>5</sub>H<sub>12</sub>O<sub>7</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			14	5	7	2		
3	B	1	Total	C	O	P	0	0
			14	5	7	2		

- Molecule 4 is 1-HYDROXY-2-(3-PYRIDINYL)ETHYLIDENE BIS-PHOSPHONIC ACID (three-letter code: RIS) (formula: C<sub>7</sub>H<sub>11</sub>NO<sub>7</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			17	7	1	7	2		
4	B	1	Total	C	N	O	P	0	0
			17	7	1	7	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	197	Total	O	0	0
			197	197		
5	B	216	Total	O	0	0
			216	216		

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Geranyltranstransferase



- Molecule 1: Geranyltranstransferase



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.80Å 88.80Å 174.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95	Depositor
% Data completeness (in resolution range)	99.3 (20.00-1.95)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.206 , 0.239	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5022	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: IPE, RIS, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/2303	0.80	14/3113 (0.4%)
1	B	0.47	0/2351	0.81	13/3178 (0.4%)
All	All	0.46	0/4654	0.80	27/6291 (0.4%)

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	193	ASP	CB-CG-OD2	8.15	125.64	118.30
1	B	114	ASP	CB-CG-OD2	7.58	125.12	118.30
1	B	289	ASP	CB-CG-OD2	7.42	124.98	118.30
1	B	156	ASP	CB-CG-OD2	7.00	124.60	118.30
1	A	289	ASP	CB-CG-OD2	6.77	124.39	118.30
1	A	113	ASP	CB-CG-OD2	6.59	124.23	118.30
1	A	286	ASP	CB-CG-OD2	6.45	124.10	118.30
1	B	252	ASP	CB-CG-OD2	6.41	124.07	118.30
1	B	23	ASP	CB-CG-OD2	6.07	123.76	118.30
1	A	150	ASP	CB-CG-OD2	5.96	123.66	118.30
1	A	252	ASP	CB-CG-OD2	5.94	123.65	118.30
1	B	113	ASP	CB-CG-OD2	5.91	123.62	118.30
1	A	244	ASP	CB-CG-OD2	5.84	123.56	118.30
1	A	158	ASP	CB-CG-OD2	5.80	123.53	118.30
1	A	112	ASP	CB-CG-OD2	5.80	123.52	118.30
1	A	114	ASP	CB-CG-OD2	5.72	123.44	118.30
1	B	286	ASP	CB-CG-OD2	5.63	123.37	118.30
1	A	220	ASP	CB-CG-OD2	5.58	123.33	118.30
1	A	290	ASP	CB-CG-OD2	5.58	123.32	118.30
1	A	156	ASP	CB-CG-OD2	5.54	123.28	118.30
1	B	148	ASP	CB-CG-OD2	5.36	123.13	118.30
1	B	184	ASP	CB-CG-OD2	5.29	123.06	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	112	ASP	CB-CG-OD2	5.23	123.01	118.30
1	A	248	ASP	CB-CG-OD2	5.20	122.98	118.30
1	B	88	ASP	CB-CG-OD1	5.18	122.96	118.30
1	B	106	ASP	CB-CG-OD2	5.03	122.82	118.30
1	A	148	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2263	0	2270	18	0
1	B	2278	0	2291	18	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	14	0	9	2	0
3	B	14	0	9	1	0
4	A	17	0	7	0	0
4	B	17	0	7	1	0
5	A	197	0	0	5	1
5	B	216	0	0	2	1
All	All	5022	0	4593	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:PRO:HD3	1:A:123:HIS:CD2	2.29	0.67
1:B:110:MET:SD	1:B:183:LEU:HD21	2.37	0.65
1:A:98:HIS:HE1	3:A:900:IPE:O3B	1.85	0.59
1:B:249[A]:VAL:CG2	1:B:277:LEU:HD21	2.35	0.56
1:B:82[A]:VAL:CG2	1:B:87:LEU:HD11	2.36	0.54

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:HIS:CE1	1:A:135:GLY:HA3	2.43	0.54
1:B:98:HIS:HE1	3:B:902:IPE:O3B	1.91	0.53
1:B:249[A]:VAL:HG21	1:B:277:LEU:HD21	1.93	0.51
1:A:125:LYS:NZ	5:A:1053:HOH:O	2.42	0.51
1:A:239:ALA:HA	1:A:242:VAL:CG1	2.42	0.50
1:A:317:GLN:NE2	5:A:1030:HOH:O	2.46	0.49
1:B:249[B]:VAL:CG1	1:B:277:LEU:HD21	2.42	0.49
1:B:98:HIS:HD2	5:B:923:HOH:O	1.97	0.48
1:A:104:HIS:O	1:A:110:MET:HG3	2.14	0.48
1:A:158:ASP:OD2	1:A:219:GLY:HA2	2.13	0.48
1:A:104:HIS:HE1	1:A:135:GLY:HA3	1.79	0.47
1:A:260:GLN:NE2	5:A:1099:HOH:O	2.43	0.47
1:B:23:ASP:OD1	1:B:25:PRO:HD2	2.14	0.47
1:B:202:LYS:O	4:B:903:RIS:HC5	2.16	0.45
1:B:71:PHE:CD1	1:B:311:LEU:HD22	2.52	0.45
1:A:208:ARG:HA	1:A:233:ALA:HB1	1.98	0.45
1:B:102:LEU:HD13	1:B:116:ARG:HD3	1.99	0.45
1:B:249[B]:VAL:HG11	1:B:277:LEU:HD21	1.98	0.45
1:A:98:HIS:HD2	5:A:927:HOH:O	2.01	0.44
1:A:247:LEU:HB3	1:A:256:LEU:HD21	2.00	0.43
1:B:208:ARG:HA	1:B:233:ALA:HB1	2.00	0.43
1:A:55:GLU:HA	1:A:58:GLN:HE21	1.84	0.43
1:B:72:LEU:CD2	1:B:311:LEU:HD23	2.48	0.43
1:A:113:ASP:O	1:A:123:HIS:HE1	2.01	0.43
1:A:58:GLN:NE2	5:A:940:HOH:O	2.48	0.43
1:B:29:GLU:HG3	5:B:1028:HOH:O	2.19	0.42
1:A:203:THR:HG23	3:A:900:IPE:H42	2.00	0.42
1:B:77:GLY:C	1:B:82[A]:VAL:HG22	2.40	0.41
1:B:245:ASP:HB3	1:B:272:PRO:CD	2.51	0.41
1:B:242:VAL:HG22	1:B:284:ALA:HB2	2.02	0.41
1:A:83:SER:HB3	1:A:86:THR:HG22	2.01	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1096:HOH:O	5:B:1105:HOH:O[5_665]	2.08	0.12

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/299 (100%)	297 (99%)	2 (1%)	0	100	100
1	B	305/299 (102%)	301 (99%)	4 (1%)	0	100	100
All	All	604/598 (101%)	598 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/234 (101%)	224 (95%)	12 (5%)	24	11
1	B	242/234 (103%)	225 (93%)	17 (7%)	15	5
All	All	478/468 (102%)	449 (94%)	29 (6%)	20	7

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	MET
1	A	82	VAL
1	A	86	THR
1	A	100	TYR
1	A	183	LEU
1	A	195	LEU
1	A	197	ARG
1	A	224	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	231	LYS
1	A	285	ARG
1	A	300	GLU
1	A	320	LYS
1	B	63	LEU
1	B	100	TYR
1	B	157	ARG
1	B	182	ASP
1	B	183	LEU
1	B	192	LEU
1	B	199	HIS
1	B	242	VAL
1	B	248	ASP
1	B	249[A]	VAL
1	B	249[B]	VAL
1	B	259	ARG
1	B	264[A]	GLN
1	B	264[B]	GLN
1	B	265	GLN
1	B	296	LYS
1	B	302	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN
1	A	34	GLN
1	A	49	GLN
1	A	58	GLN
1	A	98	HIS
1	A	104	HIS
1	A	123	HIS
1	A	130	ASN
1	A	260	GLN
1	A	317	GLN
1	B	26	GLN
1	B	34	GLN
1	B	58	GLN
1	B	98	HIS
1	B	130	ASN
1	B	317	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	IPE	B	902	-	11,13,13	0.62	0	15,19,19	0.97	0
4	RIS	A	901	2	17,17,17	6.25	6 (35%)	27,27,27	1.50	5 (18%)
3	IPE	A	900	-	11,13,13	0.56	0	15,19,19	1.00	1 (6%)
4	RIS	B	903	2	17,17,17	6.26	7 (41%)	27,27,27	1.64	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IPE	B	902	-	-	0/13/13/13	-
4	RIS	A	901	2	-	3/23/23/23	0/1/1/1
3	IPE	A	900	-	-	0/13/13/13	-
4	RIS	B	903	2	-	3/23/23/23	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	901	RIS	P14-C8	-13.92	1.75	1.85
4	A	901	RIS	P9-C8	-13.65	1.75	1.85
4	B	903	RIS	P14-C8	-12.73	1.76	1.85
4	B	903	RIS	P9-O11	11.69	1.68	1.50
4	A	901	RIS	P9-O11	11.69	1.68	1.50
4	B	903	RIS	P14-O17	11.58	1.68	1.50
4	B	903	RIS	P9-C8	-10.87	1.77	1.85
4	A	901	RIS	P14-O16	7.20	1.68	1.54
4	A	901	RIS	P14-O15	7.18	1.67	1.54
4	B	903	RIS	P14-O15	7.16	1.67	1.54
4	B	903	RIS	P9-O10	7.03	1.67	1.54
4	A	901	RIS	P9-O10	6.33	1.66	1.54
4	B	903	RIS	P14-O16	-2.33	1.50	1.54

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	903	RIS	C7-C2-C1	-3.98	115.75	121.07
4	A	901	RIS	C7-C2-C1	-3.35	116.60	121.07
4	A	901	RIS	O16-P14-C8	3.33	113.64	106.17
4	B	903	RIS	O10-P9-O11	-3.05	106.21	113.06
4	B	903	RIS	O15-P14-C8	2.82	112.50	106.17
4	B	903	RIS	C7-C2-C3	2.73	125.73	121.46
4	A	901	RIS	C5-N4-C3	2.62	121.38	116.85
4	B	903	RIS	O12-P9-C8	2.32	111.38	106.17
4	A	901	RIS	C7-C2-C3	2.22	124.93	121.46
4	B	903	RIS	C8-C7-C2	2.16	119.81	116.10
4	A	901	RIS	O16-P14-O17	-2.12	108.30	113.06
3	A	900	IPE	C5-C3-C2	2.02	121.58	115.24
4	B	903	RIS	C5-N4-C3	2.00	120.32	116.85

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	901	RIS	C1-C2-C7-C8
4	A	901	RIS	C3-C2-C7-C8
4	B	903	RIS	C1-C2-C7-C8
4	B	903	RIS	C3-C2-C7-C8
4	A	901	RIS	C7-C8-P9-O11
4	B	903	RIS	C7-C8-P9-O11


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	902	IPE	1	0
3	A	900	IPE	2	0
4	B	903	RIS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.