

wwPDB X-ray Structure Validation Summary Report (i)

Feb 4, 2024 – 08:21 PM EST

PDB ID : 1STM

Title : SATELLITE PANICUM MOSAIC VIRUS

Authors : Ban, N.; McPherson, A.

Deposited on : 1995-07-12

Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain			
1	A	157	76%	11%	•••	10%
1	В	157	75%	12%	•••	10%
1	С	157	76%	11%	•••	10%
1	D	157	76%	11%	•••	10%
1	Е	157	76%	11%	•••	10%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5633 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called SATELLITE PANICUM MOSAIC VIRUS.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	141	Total	С	N	О	S	0	0	0
1	A	141	1061	664	184	208	5	0	U	0
1	В	141	Total	С	N	О	S	0	0	0
1	Ъ	141	1061	664	184	208	5	0	U	
1	С	141	Total	С	N	О	S	0	0	0
1		141	1061	664	184	208	5	0	0	
1	D	141	Total	С	N	О	S	0	0	0
1	ט	141	1061	664	184	208	5	0	U	0
1	Е	141	Total	С	N	О	S	0	0	0
1	<u> 1</u> 2	141	1061	664	184	208	5		U	0

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	66	Total O 66 66	0	0
2	В	66	Total O 66 66	0	0
2	С	66	Total O 66 66	0	0
2	D	66	Total O 66 66	0	0
2	E	64	Total O 64 64	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: SATELLITE PANICUM MOSAIC VIRUS





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 42 3 2	Depositor
Cell constants	183.00Å 183.00Å 183.00Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 - 1.90	Depositor
% Data completeness	(Not available) (30.00-1.90)	Depositor
(in resolution range)	(1100 available) (90.00 1.50)	Берозног
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.210 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5633	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.50	0/1081	0.91	4/1474~(0.3%)	
1	В	0.50	0/1081	0.91	4/1474 (0.3%)	
1	С	0.50	0/1081	0.91	4/1474 (0.3%)	
1	D	0.50	0/1081	0.91	4/1474 (0.3%)	
1	Е	0.50	0/1081	0.91	4/1474 (0.3%)	
All	All	0.50	0/5405	0.91	$20/7370 \ (0.3\%)$	

There are no bond length outliers.

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	В	65	VAL	CB-CA-C	-5.63	100.70	111.40
1	A	65	VAL	CB-CA-C	-5.61	100.73	111.40
1	D	65	VAL	CB-CA-C	-5.61	100.73	111.40
1	Е	65	VAL	CB-CA-C	-5.60	100.75	111.40
1	С	65	VAL	CB-CA-C	-5.59	100.77	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1061	0	1045	16	0
1	В	1061	0	1045	17	0
1	С	1061	0	1045	16	0

Continued on next page...



$\alpha \cdots$, r	•	
Continued	trom	mromonie	maaa
-	110116	DICULUUS	Duuc
	J	1	1

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1061	0	1045	16	0
1	Е	1061	0	1045	16	1
2	A	66	0	0	2	0
2	В	66	0	0	2	0
2	С	66	0	0	2	0
2	D	66	0	0	2	0
2	Е	64	0	0	2	0
All	All	5633	0	5225	71	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 71 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:E:113:PHE:O	1:E:116:THR:HG23	1.76	0.86
1:C:65:VAL:HG22	1:D:92:LEU:HD13	1.58	0.85
1:B:113:PHE:O	1:B:116:THR:HG23	1.76	0.85
1:A:113:PHE:O	1:A:116:THR:HG23	1.76	0.85
1:D:65:VAL:HG22	1:E:92:LEU:HD13	1.58	0.85

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:117:GLU:OE1	1:E:117:GLU:OE1[13_455]	1.72	0.48

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	139/157~(88%)	137 (99%)	2 (1%)	0	100	100
1	В	139/157~(88%)	137 (99%)	2 (1%)	0	100	100
1	С	139/157~(88%)	137 (99%)	2 (1%)	0	100	100
1	D	139/157 (88%)	137 (99%)	2 (1%)	0	100	100
1	E	139/157~(88%)	137 (99%)	2 (1%)	0	100	100
All	All	695/785~(88%)	685 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perc	entiles
1	A	115/128~(90%)	109~(95%)	6 (5%)	23	14
1	В	115/128 (90%)	109 (95%)	6 (5%)	23	14
1	С	115/128 (90%)	109 (95%)	6 (5%)	23	14
1	D	115/128 (90%)	109 (95%)	6 (5%)	23	14
1	Е	115/128 (90%)	109 (95%)	6 (5%)	23	14
All	All	575/640 (90%)	545 (95%)	30 (5%)	23	14

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	37	THR
1	Е	65	VAL
1	С	78	ASN
1	Е	78	ASN
1	Е	26	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:



Mol	Chain	Res	Type
1	A	78	ASN
1	В	78	ASN
1	С	78	ASN
1	D	78	ASN
1	Ε	78	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

