



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 5, 2024 – 12:26 AM EST

PDB ID : 1TDF
Title : CRYSTAL STRUCTURE OF ESCHERICHIA COLI THIOREDOXIN REDUCTASE REFINED AT 2 ANGSTROM RESOLUTION: IMPLICATIONS FOR A LARGE CONFORMATIONAL CHANGE DURING CATALYSIS
Authors : Waksman, G.; Krishna, T.S.R.; Williams Junior, C.H.; Kuriyan, J.
Deposited on : 1994-01-14
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	316	85% 12%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3530 atoms, of which 877 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called THIOREDOXIN REDUCTASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	316	2941	1491	547	424	469	10	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	138	SER	CYS	conflict	UNP P0A9P4

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	P		
2	A	1	60	27	7	9	15	2	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
3	A	1	55	21	7	7	17	3	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
4	A	158	474	316	158	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: THIOREDOXIN REDUCTASE

Chain A:  85% 12%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	123.80Å 123.80Å 81.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	6.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.179 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3530	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/2433	1.13	8/3295 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	52	TRP	CD1-CG-CD2	7.27	112.12	106.30
1	A	311	ARG	NE-CZ-NH1	7.27	123.93	120.30
1	A	67	ARG	NE-CZ-NH1	7.13	123.86	120.30
1	A	52	TRP	CE2-CD2-CG	-7.05	101.66	107.30
1	A	67	ARG	NE-CZ-NH2	-6.69	116.95	120.30
1	A	198	ASN	N-CA-C	5.54	125.97	111.00
1	A	197	GLY	CA-C-N	-5.33	105.47	117.20
1	A	130	ARG	NE-CZ-NH1	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2394	547	2356	20	1
2	A	53	7	31	1	0
3	A	48	7	25	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	158	316	0	2	1
All	All	2653	877	2412	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:THR:HG22	1:A:228:SER:HB3	1.76	0.67
3:A:481:NAP:N1A	4:A:827:HOH:O	2.29	0.66
1:A:293:ARG:HH22	3:A:481:NAP:H6N	1.73	0.53
1:A:176:ARG:HG3	3:A:481:NAP:C2A	2.40	0.50
1:A:59:LEU:HD21	1:A:64:LEU:HG	1.94	0.49
1:A:85:ASN:HA	1:A:251:ILE:HD11	1.95	0.48
1:A:174:ILE:HD13	1:A:222:LEU:HD11	1.96	0.48
1:A:87:VAL:HG13	1:A:89:LEU:HD13	1.97	0.47
1:A:178:ASP:HA	1:A:203:THR:HB	1.98	0.46
1:A:119:LEU:HD22	3:A:481:NAP:N6A	2.31	0.46
1:A:94:PHE:O	1:A:104:THR:HA	2.15	0.46
1:A:184:LYS:HD2	1:A:185:ILE:N	2.32	0.44
1:A:34:ILE:HD13	1:A:84:ILE:HD11	1.99	0.44
1:A:43:LEU:HB2	4:A:615:HOH:O	2.17	0.44
1:A:6:SER:O	1:A:105:CYS:HA	2.19	0.43
1:A:86:LYS:NZ	1:A:86:LYS:HB2	2.35	0.42
1:A:183:GLU:HB3	1:A:184:LYS:NZ	2.34	0.42
1:A:59:LEU:HD13	1:A:67:ARG:NH2	2.35	0.42
1:A:153:GLY:HA3	1:A:175:HIS:CE1	2.54	0.41
2:A:500:FAD:H9	2:A:500:FAD:H1'1	1.90	0.41
1:A:160:GLU:O	1:A:164:LEU:HB2	2.21	0.41
1:A:59:LEU:HD13	1:A:67:ARG:HH21	1.87	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:ARG:HH12	4:A:701:HOH:H1[12_554]	1.31	0.29

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	314/316 (99%)	298 (95%)	14 (4%)	2 (1%)	25 31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	228	SER
1	A	231	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	251/251 (100%)	235 (94%)	16 (6%)	17 23

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LEU
1	A	43	LEU
1	A	55	ASP
1	A	64	LEU
1	A	66	GLU
1	A	87	VAL
1	A	100	ASN
1	A	164	LEU
1	A	173	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	181	ARG
1	A	184	LYS
1	A	196	ASN
1	A	198	ASN
1	A	201	LEU
1	A	236	VAL
1	A	251	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	155	ASN
1	A	166	ASN
1	A	214	GLN
1	A	248	ASN
1	A	266	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAP	A	481	-	45,52,52	1.44	7 (15%)	56,80,80	1.33	5 (8%)
2	FAD	A	500	-	53,58,58	0.88	1 (1%)	68,89,89	0.81	2 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	481	-	-	7/31/67/67	0/5/5/5
2	FAD	A	500	-	-	4/30/50/50	0/6/6/6

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	481	NAP	C3N-C7N	5.24	1.58	1.50
3	A	481	NAP	C2N-N1N	3.12	1.38	1.35
3	A	481	NAP	O4D-C1D	2.84	1.45	1.41
3	A	481	NAP	C4N-C3N	-2.66	1.34	1.39
3	A	481	NAP	C2N-C3N	-2.51	1.35	1.39
3	A	481	NAP	C8A-N7A	-2.44	1.30	1.34
2	A	500	FAD	C8A-N7A	-2.34	1.30	1.34
3	A	481	NAP	O4B-C1B	2.26	1.44	1.41

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	481	NAP	C3N-C7N-N7N	4.11	122.68	117.75
3	A	481	NAP	C6N-N1N-C2N	-3.69	118.61	121.97
3	A	481	NAP	O7N-C7N-C3N	-3.37	115.60	119.63
3	A	481	NAP	PN-O3-PA	-2.56	124.06	132.83
2	A	500	FAD	C5A-C6A-N6A	2.24	123.75	120.35
2	A	500	FAD	O3'-C3'-C4'	-2.15	103.61	108.81
3	A	481	NAP	O4B-C4B-C3B	-2.07	101.03	105.11

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	FAD	PA-O3P-P-O5'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	481	NAP	C5D-O5D-PN-O3
3	A	481	NAP	C2D-C1D-N1N-C2N
3	A	481	NAP	C2D-C1D-N1N-C6N
2	A	500	FAD	P-O3P-PA-O1A
3	A	481	NAP	C5D-O5D-PN-O1N
3	A	481	NAP	C5D-O5D-PN-O2N
2	A	500	FAD	P-O3P-PA-O2A
3	A	481	NAP	O4B-C4B-C5B-O5B
3	A	481	NAP	C4D-C5D-O5D-PN
2	A	500	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	481	NAP	4	0
2	A	500	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.