



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 17, 2023 – 06:00 PM EST

PDB ID : 1VKN  
Title : Crystal structure of N-acetyl-gamma-glutamyl-phosphate reductase (TM1782) from *Thermotoga maritima* at 1.80 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2004-06-11  
Resolution : 1.80 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	88% 9% .
1	B	351	88% 8% ..
1	C	351	86% 10% .
1	D	351	12% 87% 8% . 5%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 11278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-acetyl-gamma-glutamyl-phosphate reductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	340	2654	1707	438	498	3	8	0	0	0
1	B	340	2667	1712	440	504	3	8	0	2	0
1	C	336	2607	1676	428	492	3	8	0	0	0
1	D	335	2594	1672	426	485	3	8	0	0	0

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	UNP Q9X2A2
A	-10	GLY	-	expression tag	UNP Q9X2A2
A	-9	SER	-	expression tag	UNP Q9X2A2
A	-8	ASP	-	expression tag	UNP Q9X2A2
A	-7	LYS	-	expression tag	UNP Q9X2A2
A	-6	ILE	-	expression tag	UNP Q9X2A2
A	-5	HIS	-	expression tag	UNP Q9X2A2
A	-4	HIS	-	expression tag	UNP Q9X2A2
A	-3	HIS	-	expression tag	UNP Q9X2A2
A	-2	HIS	-	expression tag	UNP Q9X2A2
A	-1	HIS	-	expression tag	UNP Q9X2A2
A	0	HIS	-	expression tag	UNP Q9X2A2
A	1	MSE	MET	modified residue	UNP Q9X2A2
A	210	MSE	MET	modified residue	UNP Q9X2A2
A	233	MSE	MET	modified residue	UNP Q9X2A2
A	271	MSE	MET	modified residue	UNP Q9X2A2
A	290	MSE	MET	modified residue	UNP Q9X2A2
A	292	MSE	MET	modified residue	UNP Q9X2A2
A	302	MSE	MET	modified residue	UNP Q9X2A2
A	320	MSE	MET	modified residue	UNP Q9X2A2
A	323	MSE	MET	modified residue	UNP Q9X2A2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-11	MET	-	expression tag	UNP Q9X2A2
B	-10	GLY	-	expression tag	UNP Q9X2A2
B	-9	SER	-	expression tag	UNP Q9X2A2
B	-8	ASP	-	expression tag	UNP Q9X2A2
B	-7	LYS	-	expression tag	UNP Q9X2A2
B	-6	ILE	-	expression tag	UNP Q9X2A2
B	-5	HIS	-	expression tag	UNP Q9X2A2
B	-4	HIS	-	expression tag	UNP Q9X2A2
B	-3	HIS	-	expression tag	UNP Q9X2A2
B	-2	HIS	-	expression tag	UNP Q9X2A2
B	-1	HIS	-	expression tag	UNP Q9X2A2
B	0	HIS	-	expression tag	UNP Q9X2A2
B	1	MSE	MET	modified residue	UNP Q9X2A2
B	210	MSE	MET	modified residue	UNP Q9X2A2
B	233	MSE	MET	modified residue	UNP Q9X2A2
B	271	MSE	MET	modified residue	UNP Q9X2A2
B	290	MSE	MET	modified residue	UNP Q9X2A2
B	292	MSE	MET	modified residue	UNP Q9X2A2
B	302	MSE	MET	modified residue	UNP Q9X2A2
B	320	MSE	MET	modified residue	UNP Q9X2A2
B	323	MSE	MET	modified residue	UNP Q9X2A2
C	-11	MET	-	expression tag	UNP Q9X2A2
C	-10	GLY	-	expression tag	UNP Q9X2A2
C	-9	SER	-	expression tag	UNP Q9X2A2
C	-8	ASP	-	expression tag	UNP Q9X2A2
C	-7	LYS	-	expression tag	UNP Q9X2A2
C	-6	ILE	-	expression tag	UNP Q9X2A2
C	-5	HIS	-	expression tag	UNP Q9X2A2
C	-4	HIS	-	expression tag	UNP Q9X2A2
C	-3	HIS	-	expression tag	UNP Q9X2A2
C	-2	HIS	-	expression tag	UNP Q9X2A2
C	-1	HIS	-	expression tag	UNP Q9X2A2
C	0	HIS	-	expression tag	UNP Q9X2A2
C	1	MSE	MET	modified residue	UNP Q9X2A2
C	210	MSE	MET	modified residue	UNP Q9X2A2
C	233	MSE	MET	modified residue	UNP Q9X2A2
C	271	MSE	MET	modified residue	UNP Q9X2A2
C	290	MSE	MET	modified residue	UNP Q9X2A2
C	292	MSE	MET	modified residue	UNP Q9X2A2
C	302	MSE	MET	modified residue	UNP Q9X2A2
C	320	MSE	MET	modified residue	UNP Q9X2A2
C	323	MSE	MET	modified residue	UNP Q9X2A2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-11	MET	-	expression tag	UNP Q9X2A2
D	-10	GLY	-	expression tag	UNP Q9X2A2
D	-9	SER	-	expression tag	UNP Q9X2A2
D	-8	ASP	-	expression tag	UNP Q9X2A2
D	-7	LYS	-	expression tag	UNP Q9X2A2
D	-6	ILE	-	expression tag	UNP Q9X2A2
D	-5	HIS	-	expression tag	UNP Q9X2A2
D	-4	HIS	-	expression tag	UNP Q9X2A2
D	-3	HIS	-	expression tag	UNP Q9X2A2
D	-2	HIS	-	expression tag	UNP Q9X2A2
D	-1	HIS	-	expression tag	UNP Q9X2A2
D	0	HIS	-	expression tag	UNP Q9X2A2
D	1	MSE	MET	modified residue	UNP Q9X2A2
D	210	MSE	MET	modified residue	UNP Q9X2A2
D	233	MSE	MET	modified residue	UNP Q9X2A2
D	271	MSE	MET	modified residue	UNP Q9X2A2
D	290	MSE	MET	modified residue	UNP Q9X2A2
D	292	MSE	MET	modified residue	UNP Q9X2A2
D	302	MSE	MET	modified residue	UNP Q9X2A2
D	320	MSE	MET	modified residue	UNP Q9X2A2
D	323	MSE	MET	modified residue	UNP Q9X2A2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	263	Total O 263 263	0	0
2	B	213	Total O 213 213	0	0
2	C	191	Total O 191 191	0	0
2	D	89	Total O 89 89	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: N-acetyl-gamma-glutamyl-phosphate reductase




- Molecule 1: N-acetyl-gamma-glutamyl-phosphate reductase




- Molecule 1: N-acetyl-gamma-glutamyl-phosphate reductase



- Molecule 1: N-acetyl-gamma-glutamyl-phosphate reductase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.10Å 92.10Å 100.88Å 90.00° 112.76° 90.00°	Depositor
Resolution (Å)	46.51 – 1.80 46.51 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.1 (46.51-1.80) 98.1 (46.51-1.80)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.20 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.2.0001	Depositor
R, $R_{free}$	0.178 , 0.212 0.187 , 0.218	Depositor DCC
$R_{free}$ test set	6565 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.1	Xtrriage
Anisotropy	0.252	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 47.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.012 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	11278	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/2705	0.82	1/3651 (0.0%)
1	B	0.76	0/2726	0.87	8/3681 (0.2%)
1	C	0.76	0/2658	0.82	6/3593 (0.2%)
1	D	0.60	0/2645	0.76	6/3574 (0.2%)
All	All	0.73	0/10734	0.82	21/14499 (0.1%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	235	ARG	NE-CZ-NH2	-7.95	116.33	120.30
1	B	197	ARG	NE-CZ-NH2	-7.61	116.50	120.30
1	C	93	ASP	CB-CG-OD2	6.69	124.32	118.30
1	B	68	ASP	CB-CG-OD2	6.60	124.24	118.30
1	A	93	ASP	CB-CG-OD2	6.53	124.17	118.30
1	B	197	ARG	NE-CZ-NH1	6.26	123.43	120.30
1	D	97	ASP	CB-CG-OD2	6.12	123.81	118.30
1	D	93	ASP	CB-CG-OD2	6.08	123.78	118.30
1	D	80	TYR	N-CA-C	5.91	126.96	111.00
1	C	101	ASP	CB-CG-OD2	5.87	123.58	118.30
1	D	246	ASP	CB-CG-OD2	5.87	123.58	118.30
1	C	68	ASP	CB-CG-OD2	5.77	123.50	118.30
1	D	327	ASP	CB-CG-OD2	5.51	123.26	118.30
1	C	327	ASP	CB-CG-OD2	5.42	123.18	118.30
1	C	306	ASP	CB-CG-OD2	5.34	123.11	118.30
1	D	81	ASP	CB-CG-OD2	5.24	123.02	118.30
1	C	97	ASP	CB-CG-OD2	5.10	122.89	118.30
1	B	171	ASP	CB-CG-OD2	5.03	122.83	118.30
1	B	164[A]	ASP	CB-CG-OD2	5.03	122.83	118.30
1	B	164[B]	ASP	CB-CG-OD2	5.03	122.83	118.30
1	B	81	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2654	0	2649	17	0
1	B	2667	0	2655	14	0
1	C	2607	0	2572	18	0
1	D	2594	0	2566	15	0
2	A	263	0	0	3	0
2	B	213	0	0	3	0
2	C	191	0	0	4	0
2	D	89	0	0	0	0
All	All	11278	0	10442	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:129:LEU:HD21	1:D:217:ILE:HD11	1.69	0.74
1:A:129:LEU:HD21	1:A:217:ILE:HD11	1.72	0.71
1:C:278:LYS:HG2	2:C:431:HOH:O	1.92	0.69
1:C:286:VAL:HG22	1:C:305:ILE:HG22	1.74	0.69
1:B:79:SER:O	1:B:83:VAL:HG23	2.01	0.60
1:B:35:ARG:HG3	2:B:427:HOH:O	2.02	0.58
1:A:82:LEU:C	1:A:82:LEU:HD12	2.27	0.55
1:B:197:ARG:NE	2:B:406:HOH:O	2.41	0.54
1:A:201:VAL:HG13	1:A:227:THR:HG23	1.91	0.53
1:A:286:VAL:HG22	1:A:305:ILE:HG22	1.92	0.51
1:C:41:LYS:HB2	1:C:44:GLU:HG3	1.93	0.51
1:C:127:PRO:HB3	1:C:322:ILE:CD1	2.41	0.51
1:B:327:ASP:HB2	1:B:330:LYS:HG2	1.94	0.50
1:C:82:LEU:HD12	1:C:82:LEU:C	2.32	0.49
1:A:41:LYS:HG2	1:A:54:ILE:HD13	1.93	0.49
1:A:123:VAL:CG1	1:A:140:VAL:HG22	2.43	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:7:ILE:HD11	1:C:70:LEU:HD11	1.95	0.48
1:A:41:LYS:HB2	1:A:44:GLU:HG3	1.96	0.47
1:A:146:TYR:HB2	1:A:147:PRO:HD3	1.95	0.47
1:B:186:ASP:HB3	1:B:197:ARG:HH22	1.78	0.47
1:A:87:LYS:NZ	2:A:529:HOH:O	2.47	0.47
1:B:83:VAL:HG22	1:B:91:ILE:CD1	2.45	0.47
1:C:156:PRO:HG3	1:C:259:PHE:CD1	2.50	0.47
1:D:79:SER:HA	1:D:81:ASP:N	2.30	0.47
1:A:173:LYS:HE3	1:D:173:LYS:HE3	1.97	0.47
1:C:156:PRO:HG3	1:C:259:PHE:CG	2.51	0.46
1:C:278:LYS:CG	2:C:431:HOH:O	2.59	0.46
1:C:248:SER:OG	1:C:251:GLU:HB2	2.16	0.46
1:C:251:GLU:CG	2:C:446:HOH:O	2.64	0.46
1:D:237:ILE:HG13	1:D:307:ASN:HB3	1.98	0.45
1:B:126:LEU:HD23	1:B:129:LEU:HD22	1.97	0.45
1:A:327:ASP:HB3	2:A:586:HOH:O	2.17	0.45
1:B:82:LEU:C	1:B:82:LEU:HD12	2.37	0.44
1:B:2:ILE:HD11	1:B:326:LEU:HD11	1.98	0.44
1:A:176:VAL:HG21	1:A:192:VAL:HG11	1.99	0.44
1:C:176:VAL:HG21	1:C:192:VAL:HG11	2.01	0.43
1:D:98:PHE:O	1:D:122:ARG:HD2	2.18	0.43
1:B:238:LEU:HD21	1:C:198:PRO:HG3	2.00	0.43
1:B:197:ARG:HD3	2:B:379:HOH:O	2.17	0.43
1:C:207:VAL:N	1:C:208:PRO:HD2	2.34	0.43
1:B:59:ASP:HB3	1:B:62:LYS:HD2	2.01	0.43
1:A:16:GLU:OE1	2:A:390:HOH:O	2.21	0.42
1:D:83:VAL:HG13	1:D:91:ILE:HG13	2.00	0.42
1:C:333:GLU:HG3	2:C:383:HOH:O	2.18	0.42
1:D:79:SER:HB2	1:D:83:VAL:HG23	2.01	0.42
1:B:282:GLY:O	1:B:334:PHE:HE2	2.02	0.42
1:D:79:SER:O	1:D:79:SER:OG	2.32	0.42
1:D:90:LYS:HD2	1:D:323:MSE:HE2	2.01	0.42
1:D:127:PRO:HB3	1:D:322:ILE:HD11	2.01	0.42
1:D:176:VAL:HG21	1:D:192:VAL:HG11	2.02	0.42
1:B:158:LEU:HD22	1:B:165:PRO:HB3	2.02	0.42
1:C:176:VAL:HA	1:C:230:LEU:HD11	2.02	0.41
1:A:142:ASN:HA	1:A:143:PRO:HD3	1.93	0.41
1:A:127:PRO:HB3	1:A:322:ILE:HD11	2.03	0.41
1:A:158:LEU:HD22	1:A:165:PRO:HB3	2.03	0.41
1:C:124:TYR:CE2	1:C:213:GLU:HG3	2.55	0.41
1:D:207:VAL:N	1:D:208:PRO:HD2	2.36	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:249:LEU:HD23	1:C:271:MSE:HE3	2.03	0.40
1:D:335:THR:HB	1:D:336:PRO:HD2	2.03	0.40
1:D:237:ILE:O	1:D:304:ALA:HA	2.22	0.40
1:A:290:MSE:HA	1:A:300:ILE:O	2.21	0.40
1:D:142:ASN:HA	1:D:143:PRO:HD3	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	338/351 (96%)	332 (98%)	5 (2%)	1 (0%)	41 27
1	B	340/351 (97%)	334 (98%)	5 (2%)	1 (0%)	41 27
1	C	332/351 (95%)	326 (98%)	5 (2%)	1 (0%)	41 27
1	D	331/351 (94%)	322 (97%)	8 (2%)	1 (0%)	41 27
All	All	1341/1404 (96%)	1314 (98%)	23 (2%)	4 (0%)	41 27

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	80	TYR
1	A	312	ALA
1	B	312	ALA
1	C	312	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/295 (97%)	285 (99%)	2 (1%)	84	81
1	B	290/295 (98%)	285 (98%)	5 (2%)	60	51
1	C	279/295 (95%)	275 (99%)	4 (1%)	67	59
1	D	277/295 (94%)	274 (99%)	3 (1%)	73	68
All	All	1133/1180 (96%)	1119 (99%)	14 (1%)	71	65

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	LEU
1	A	122	ARG
1	B	82	LEU
1	B	115	SER
1	B	121	LYS
1	B	122	ARG
1	B	244	LYS
1	C	54	ILE
1	C	82	LEU
1	C	84	ARG
1	C	122	ARG
1	D	80	TYR
1	D	122	ARG
1	D	238	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	262	ASN
1	D	138	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	331/351 (94%)	-0.25	2 (0%) 89 87	25, 32, 50, 67	0
1	B	331/351 (94%)	-0.18	6 (1%) 68 64	25, 34, 58, 78	0
1	C	327/351 (93%)	-0.00	8 (2%) 59 54	26, 35, 55, 73	0
1	D	326/351 (92%)	0.62	42 (12%) 3 2	30, 43, 66, 97	0
All	All	1315/1404 (93%)	0.04	58 (4%) 34 28	25, 36, 58, 97	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	82	LEU	6.7
1	D	84	ARG	5.8
1	B	0	HIS	5.5
1	D	58	PHE	5.2
1	D	83	VAL	5.1
1	D	80	TYR	4.7
1	D	86	LEU	4.4
1	D	76	ALA	4.1
1	D	114	LEU	4.0
1	D	77	GLY	3.7
1	C	86	LEU	3.5
1	D	70	LEU	3.4
1	C	118	GLU	3.1
1	D	196	LEU	3.0
1	D	89	VAL	2.9
1	D	85	GLU	2.9
1	D	81	ASP	2.9
1	D	117	TYR	2.8
1	D	74	LEU	2.8
1	D	111	GLY	2.8
1	D	79	SER	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	91	ILE	2.7
1	D	120	ILE	2.7
1	B	86	LEU	2.7
1	D	57	GLU	2.7
1	D	118	GLU	2.7
1	C	185	VAL	2.6
1	C	119	ASN	2.6
1	D	192	VAL	2.6
1	D	119	ASN	2.6
1	D	189	PHE	2.5
1	D	56	SER	2.5
1	C	84	ARG	2.5
1	D	231	VAL	2.5
1	D	7	ILE	2.4
1	A	181	ARG	2.4
1	D	178	GLY	2.4
1	D	115	SER	2.4
1	D	185	VAL	2.4
1	B	87	LYS	2.4
1	D	31	TYR	2.4
1	D	187	TYR	2.4
1	C	187	TYR	2.3
1	B	189	PHE	2.3
1	D	139	VAL	2.2
1	B	192	VAL	2.2
1	D	38	ALA	2.2
1	D	59	ASP	2.2
1	D	116	GLY	2.2
1	A	189	PHE	2.2
1	D	67	CYS	2.2
1	D	3	ARG	2.1
1	B	119	ASN	2.1
1	C	87	LYS	2.1
1	D	195	SER	2.1
1	D	63	VAL	2.0
1	C	82	LEU	2.0
1	D	326	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.