

# Full wwPDB X-ray Structure Validation Report (i)

#### Nov 2, 2021 – 04:33 AM EDT

PDB ID	:	1XEX
Title	:	Structural biochemistry of ATP-driven dimerization and DNA stimulated ac-
		tivation of SMC ATPases.
Authors	:	Lammens, A.; Schele, A.; Hopfner, KP.
Deposited on		
Resolution	:	2.50  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
$\mathrm{EDS}$	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chai	n	
1	А	182	58%	31%	• 9%
2	В	172	63%	30%	• 6%



#### $1 \mathrm{XEX}$

# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2706 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called SMC protein.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	166	Total 1296	C 829	N 224	0 240	${ m S} { m 3}$	33	0	0

• Molecule 2 is a protein called SMC protein.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	В	161	Total 1249	C 800	N 214	O 229	${ m S}{ m 6}$	28	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
В	1098	GLN	GLU	engineered mutation	GB 28375557

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Mg 1 1	0	0

• Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
4	А	1	Total 31	C 10	N 5	0 13	Р 3	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	60	Total         O           60         60	0	0
5	В	69	Total         O           69         69	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: SMC protein





## 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants	102.60Å 102.60Å 87.20Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	20.00 - 2.50	Depositor
% Data completeness	(Not available) (20.00-2.50)	Depositor
(in resolution range)	(1007 available) (20.00-2.00)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
$R, R_{free}$	0.207 , $0.261$	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2706	wwPDB-VP
Average B, all atoms $(Å^2)$	53.0	wwPDB-VP



# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.42	0/1320	0.59	0/1777	
2	В	0.42	0/1265	0.54	0/1689	
All	All	0.42	0/2585	0.57	0/3466	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1296	0	1323	48	0
2	В	1249	0	1307	46	1
3	А	1	0	0	0	0
4	А	31	0	12	0	0
5	А	60	0	0	4	0
5	В	69	0	0	4	1
All	All	2706	0	2642	88	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

A + a 1	A + a	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
2:B:1169:LYS:HB2	2:B:1171:GLN:HE22	1.19	1.02
2:B:1128:THR:HG21	2:B:1133:MET:HB3	1.48	0.95
2:B:1064:LYS:HD2	5:B:21:HOH:O	1.71	0.89
2:B:1026:GLU:OE1	2:B:1087:LYS:HE3	1.75	0.86
2:B:1043:ASN:HD22	2:B:1049:SER:H	1.24	0.84
1:A:135:ILE:CG1	2:B:1092:PRO:HG3	2.09	0.82
2:B:1017:PHE:HZ	2:B:1039:LEU:O	1.64	0.80
2:B:1169:LYS:HB2	2:B:1171:GLN:NE2	1.97	0.78
1:A:161:LEU:O	1:A:164:ASP:HB2	1.87	0.75
2:B:1128:THR:HG21	2:B:1133:MET:CB	2.19	0.73
1:A:104:ARG:HG3	1:A:114:TYR:CE2	2.25	0.71
1:A:135:ILE:HG12	2:B:1092:PRO:CG	2.24	0.67
1:A:70:GLY:HA3	5:A:1040:HOH:O	1.93	0.67
2:B:1161:MET:O	2:B:1165:GLU:HG3	1.97	0.65
1:A:119:ARG:HG2	5:A:1058:HOH:O	1.96	0.65
1:A:135:ILE:HG12	2:B:1092:PRO:HG3	1.78	0.65
1:A:119:ARG:NE	5:A:1058:HOH:O	2.29	0.65
2:B:1151:LYS:HB2	5:B:106:HOH:O	1.98	0.64
2:B:1023:ASN:CG	2:B:1087:LYS:HD2	2.16	0.64
2:B:1076:LEU:HD22	2:B:1107:ASN:HD21	1.66	0.61
2:B:1017:PHE:CZ	2:B:1039:LEU:O	2.51	0.60
2:B:1024:PHE:CD2	2:B:1039:LEU:HG	2.37	0.59
1:A:2:PRO:HD3	1:A:92:GLY:HA3	1.85	0.58
1:A:4:ILE:HD13	1:A:84:ILE:HD11	1.85	0.58
1:A:155:SER:OG	1:A:158:GLU:HG3	2.03	0.58
2:B:1023:ASN:O	2:B:1026:GLU:HG2	2.04	0.57
2:B:1021:SER:HA	2:B:1039:LEU:HD12	1.86	0.56
1:A:135:ILE:HG13	2:B:1092:PRO:HG3	1.85	0.55
1:A:119:ARG:CD	5:A:1058:HOH:O	2.55	0.55
1:A:49:VAL:HG13	1:A:50:LEU:HD13	1.87	0.55
2:B:1100:ASP:OD2	2:B:1128:THR:OG1	2.22	0.55
2:B:1128:THR:HG21	2:B:1133:MET:CG	2.38	0.54
1:A:33:GLY:O	1:A:39:LYS:HE2	2.08	0.54
2:B:1076:LEU:HD22	2:B:1107:ASN:ND2	2.23	0.54
2:B:1116:LYS:O	2:B:1119:SER:HB3	2.06	0.54
2:B:1138:ASP:O	2:B:1139:LYS:HD3	2.08	0.54
1:A:154:MET:HE3	1:A:162:LEU:HD13	1.90	0.54
1:A:124:SER:O	1:A:128:ASP:OD1	2.25	0.54
2:B:1141:ILE:HG23	2:B:1152:VAL:HG13	1.91	0.53
2:B:1010:LYS:HB2	2:B:1048:PHE:CE2	2.44	0.52
2:B:1069:MET:O	2:B:1074:LYS:HE3	2.10	0.52

Continued on next page...



Continued from previo		Interatomic	Clash	
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)	
1:A:160:ARG:O	1:A:163:ILE:HG22	2.10	0.52	
1:A:162:LEU:O	1:A:166:ILE:HG13	2.10	0.52	
1:A:7:LEU:HD23	1:A:9:LEU:HD11	1.92	0.51	
2:B:1081:PHE:O	2:B:1085:ILE:HG13	2.10	0.50	
1:A:154:MET:HE1	1:A:162:LEU:HD22	1.93	0.50	
2:B:1058:PRO:HG2	2:B:1061:LYS:HB2	1.94	0.49	
2:B:1043:ASN:ND2	2:B:1049:SER:H	2.03	0.48	
1:A:80:ALA:HB3	1:A:106:VAL:HG12	1.96	0.48	
1:A:28:PHE:C	1:A:28:PHE:CD1	2.88	0.47	
1:A:103:ARG:HB3	1:A:115:TRP:HB2	1.96	0.47	
1:A:74:GLU:HA	1:A:75:PRO:HD3	1.83	0.46	
2:B:1065:ARG:HB3	2:B:1067:GLU:OE1	2.15	0.46	
1:A:48:PHE:CZ	1:A:104:ARG:HD2	2.50	0.46	
2:B:1010:LYS:O	2:B:1010:LYS:HG2	2.15	0.46	
1:A:3:TYR:CZ	1:A:87:ASN:OD1	2.69	0.46	
1:A:88:ASN:O	1:A:98:ASP:HA	2.16	0.46	
1:A:159:ARG:HD3	2:B:1054:ILE:HD11	1.97	0.45	
1:A:7:LEU:HD12	1:A:84:ILE:HG13	1.98	0.45	
2:B:1036:SER:HA	5:B:37:HOH:O	2.17	0.44	
1:A:17:ASN:HD21	1:A:77:ALA:HA	1.80	0.44	
1:A:142:ILE:HG22	1:A:144:LEU:CD1	2.47	0.44	
1:A:4:ILE:CD1	1:A:84:ILE:HD11	2.47	0.44	
2:B:1010:LYS:HB2	2:B:1048:PHE:HE2	1.83	0.44	
1:A:61:SER:HB3	1:A:65:ASP:OD2	2.17	0.44	
1:A:49:VAL:HG22	1:A:49:VAL:O	2.18	0.43	
1:A:3:TYR:CE2	1:A:87:ASN:OD1	2.71	0.43	
2:B:1067:GLU:H	2:B:1067:GLU:CD	2.21	0.43	
2:B:1024:PHE:HD2	2:B:1039:LEU:HG	1.83	0.43	
1:A:39:LYS:HZ2	1:A:39:LYS:HB2	1.83	0.43	
2:B:1100:ASP:OD1	2:B:1128:THR:OG1	2.36	0.43	
1:A:5:GLU:OE2	1:A:99:GLU:OE2	2.35	0.43	
1:A:161:LEU:HA	1:A:164:ASP:OD2	2.19	0.43	
1:A:28:PHE:HD1	1:A:29:THR:N	2.17	0.42	
1:A:81:GLU:HB2	1:A:105:ARG:HD2	2.02	0.42	
1:A:68:PHE:HE2	1:A:74:GLU:HG2	1.84	0.42	
1:A:28:PHE:C	1:A:28:PHE:HD1	2.23	0.42	
2:B:1157:LEU:O	2:B:1161:MET:HG2	2.21	0.41	
2:B:1017:PHE:CE2	2:B:1041:LEU:HG	2.55	0.41	
1:A:166:ILE:HG21	2:B:1020:ILE:HD11	2.02	0.41	
2:B:1129:LEU:HA	5:B:8:HOH:O	2.19	0.41	
1:A:88:ASN:HB3	1:A:97:GLU:O	2.20	0.41	

Continued from previous page...

Continued on next page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1128:THR:HG21	2:B:1133:MET:HG2	2.03	0.41
2:B:1104:ASP:O	2:B:1108:VAL:HG23	2.20	0.41
2:B:1107:ASN:OD1	2:B:1110:ARG:NH2	2.54	0.41
1:A:7:LEU:HD12	1:A:84:ILE:CG1	2.51	0.40
1:A:32:VAL:C	1:A:39:LYS:HD3	2.42	0.40
1:A:160:ARG:O	1:A:160:ARG:HD2	2.21	0.40

Continued from previous page...

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1 Atom-2		Interatomic distance (Å)	Clash overlap (Å)	
2:B:1105:ASP:OD1	5:B:111:HOH:O[4_645]	2.17	0.03	

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	164/182~(90%)	145 (88%)	15~(9%)	4 (2%)	6 9
2	В	157/172~(91%)	152 (97%)	5(3%)	0	100 100
All	All	321/354 (91%)	297 (92%)	20~(6%)	4 (1%)	13 24

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	16	GLY
1	А	52	GLY
1	А	53	LEU
1	А	72	LYS



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	138/151~(91%)	129~(94%)	9~(6%)	17 33		
2	В	134/144~(93%)	128 (96%)	6 (4%)	27 51		
All	All	272/295~(92%)	257 (94%)	15 (6%)	21 41		

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	25	SER
1	А	28	PHE
1	А	73	ASN
1	А	104	ARG
1	А	105	ARG
1	А	116	LEU
1	А	120	ARG
1	А	128	ASP
1	А	144	LEU
2	В	1011	ASN
2	В	1016	THR
2	В	1087	LYS
2	В	1134	MET
2	В	1136	ASN
2	В	1158	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	17	ASN
2	В	1011	ASN
2	В	1023	ASN
2	В	1043	ASN
2	В	1171	GLN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths		Bond angles		les	
Moi Typ	туре	Ullaili	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
4	ATP	А	1001	3	26,33,33	1.26	4 (15%)	31,52,52	1.33	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	А	1001	3	-	0/18/38/38	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	А	1001	ATP	C2-N3	2.49	1.36	1.32
4	А	1001	ATP	C8-N7	-2.39	1.30	1.34
4	А	1001	ATP	PG-O2G	-2.23	1.46	1.54
4	А	1001	ATP	C4-N3	2.18	1.38	1.35



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	А	1001	ATP	O4'-C1'-C2'	-4.41	100.48	106.93
4	А	1001	ATP	C3'-C2'-C1'	3.59	106.38	100.98
4	А	1001	ATP	C4-C5-N7	2.35	111.85	109.40

All (3) bond angle outliers are listed below:

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

