



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 15, 2023 – 09:51 PM EDT

PDB ID : 1XF4  
Title : Structure of ligand-free Fab DNA-1 in space group P321 solved from crystals with perfect hemihedral twinning  
Authors : Schuermann, J.P.; Prewitt, S.P.; Deutscher, S.L.; Tanner, J.J.  
Deposited on : 2004-09-13  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	71% (green), 25% (yellow), 4% (orange), 0% (red), 0% (grey)
1	L	214	79% (green), 17% (yellow), 4% (orange), 0% (red), 0% (grey)
2	B	230	64% (green), 27% (yellow), 7% (orange), 0% (red), 2% (grey)
2	H	230	70% (green), 20% (yellow), 7% (orange), 0% (red), 3% (grey)

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	213	Total 1618	C 1012	N 270	O 330	S 6	0	0	0
1	A	213	Total 1611	C 1006	N 270	O 329	S 6	0	0	0

- Molecule 2 is a protein called Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	213	Total 1564	C 997	N 250	O 308	S 9	20	0	0
2	B	213	Total 1567	C 1000	N 250	O 308	S 9	20	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	1	GLN	-	cloning artifact	GB 3399661
H	2	VAL	-	cloning artifact	GB 3399661
H	3	LYS	-	cloning artifact	GB 3399661
H	4	LEU	-	cloning artifact	GB 3399661
H	218	HIS	-	expression tag	GB 3399661
H	219	HIS	-	expression tag	GB 3399661
H	220	HIS	-	expression tag	GB 3399661
H	221	HIS	-	expression tag	GB 3399661
H	222	HIS	-	expression tag	GB 3399661
H	223	HIS	-	expression tag	GB 3399661
B	1	GLN	-	cloning artifact	GB 3399661
B	2	VAL	-	cloning artifact	GB 3399661
B	3	LYS	-	cloning artifact	GB 3399661
B	4	LEU	-	cloning artifact	GB 3399661
B	218	HIS	-	expression tag	GB 3399661
B	219	HIS	-	expression tag	GB 3399661

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	220	HIS	-	expression tag	GB 3399661
B	221	HIS	-	expression tag	GB 3399661
B	222	HIS	-	expression tag	GB 3399661
B	223	HIS	-	expression tag	GB 3399661

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	L	1	Total	O	S	0	0
			5	4	1		
3	L	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	11	Total	O	0	0
			11	11		
4	H	4	Total	O	0	0
			4	4		
4	A	14	Total	O	0	0
			14	14		
4	B	6	Total	O	0	0
			6	6		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab light chain




- Molecule 1: Fab light chain



- Molecule 2: Fab heavy chain



- Molecule 2: Fab heavy chain





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	179.18Å 179.18Å 91.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.50 – 2.50 49.45 – 2.50	Depositor EDS
% Data completeness (in resolution range)	77.8 (49.50-2.50) 77.8 (49.45-2.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.00 (at 2.48Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.183 , 0.224 0.181 , 0.218	Depositor DCC
$R_{free}$ test set	4796 reflections (9.32%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.4	Xtrriage
Anisotropy	0.226	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 41.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.106 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6410	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	0/1647	0.86	6/2238 (0.3%)
1	L	0.66	0/1655	0.86	8/2249 (0.4%)
2	B	0.57	0/1609	0.82	4/2200 (0.2%)
2	H	0.61	0/1605	0.83	4/2194 (0.2%)
All	All	0.62	0/6516	0.84	22/8881 (0.2%)

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	61	ARG	NE-CZ-NH2	-8.51	116.05	120.30
2	B	207	ASP	CB-CG-OD2	7.24	124.82	118.30
1	A	151	ASP	CB-CG-OD2	7.00	124.60	118.30
1	L	110	ASP	CB-CG-OD2	6.99	124.59	118.30
2	H	207	ASP	CB-CG-OD2	6.87	124.48	118.30
1	A	82	ASP	CB-CG-OD2	6.60	124.24	118.30
1	L	167	ASP	CB-CG-OD2	6.25	123.93	118.30
1	L	143	ASP	CB-CG-OD2	6.16	123.84	118.30
1	A	143	ASP	CB-CG-OD2	6.03	123.72	118.30
1	A	184	ASP	CB-CG-OD2	5.90	123.61	118.30
1	L	151	ASP	CB-CG-OD2	5.73	123.46	118.30
2	B	72	ASP	CB-CG-OD2	5.49	123.24	118.30
1	A	110	ASP	CB-CG-OD2	5.40	123.16	118.30
1	L	165	ASP	CB-CG-OD2	5.39	123.15	118.30
2	B	177	LEU	CA-CB-CG	5.38	127.69	115.30
2	H	86	ASP	CB-CG-OD2	5.36	123.12	118.30
2	B	55	ASP	CB-CG-OD2	5.33	123.09	118.30
2	H	72	ASP	CB-CG-OD2	5.30	123.07	118.30
2	H	177	LEU	CA-CB-CG	5.26	127.39	115.30
1	L	61	ARG	NE-CZ-NH1	5.17	122.89	120.30

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	167	ASP	CB-CG-OD2	5.04	122.84	118.30
1	L	184	ASP	CB-CG-OD2	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1611	0	1519	37	0
1	L	1618	0	1522	21	0
2	B	1567	0	1504	32	0
2	H	1564	0	1495	25	0
3	A	5	0	0	0	0
3	L	10	0	0	0	0
4	A	14	0	0	0	0
4	B	6	0	0	0	0
4	H	4	0	0	0	0
4	L	11	0	0	0	0
All	All	6410	0	6040	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:LEU:HD13	1:A:175:MET:CE	2.04	0.88
1:L:163:TRP:CE2	1:L:175:MET:HG3	2.18	0.78
2:H:34:MET:HE2	2:H:92:CYS:HB2	1.65	0.78
1:A:136:LEU:HD13	1:A:175:MET:HE2	1.64	0.78
2:B:22:CYS:SG	2:B:34:MET:HE3	2.28	0.73
1:A:94:THR:HG21	2:B:58:LYS:HG2	1.71	0.71
1:A:136:LEU:HD13	1:A:175:MET:HE1	1.72	0.70
1:L:212:ASN:H	1:L:212:ASN:HD22	1.40	0.69

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:TRP:CE2	1:A:175:MET:HG3	2.28	0.68
1:L:163:TRP:CD2	1:L:175:MET:HG3	2.29	0.67
1:L:151:ASP:HA	1:L:191:SER:HB3	1.75	0.66
2:B:22:CYS:SG	2:B:34:MET:CE	2.84	0.65
2:B:93:VAL:HG11	2:B:100(C):MET:HE3	1.78	0.65
1:L:14:SER:O	1:L:17:GLU:HB3	1.98	0.64
1:A:149:LYS:HB2	1:A:193:THR:HB	1.79	0.63
2:B:39:GLN:HG3	2:B:44:GLY:O	2.00	0.62
1:L:116:SER:O	1:L:134:CYS:HA	2.00	0.61
1:L:161:ASN:HD22	1:L:177:SER:HA	1.66	0.61
2:B:187:THR:O	2:B:191:GLU:HB2	2.01	0.60
2:H:119:PRO:HB3	2:H:145:TYR:HB3	1.84	0.60
1:A:148:TRP:O	1:A:154:GLU:HA	2.02	0.60
1:L:185:GLU:HA	1:L:188:ARG:HH21	1.67	0.59
2:B:124:LEU:HB2	2:B:139:GLY:CA	2.33	0.59
2:H:188:TRP:CD1	2:H:189:PRO:HA	2.40	0.57
1:A:124:GLN:HG2	1:A:129:GLY:O	2.05	0.57
1:A:147:LYS:CG	1:A:154:GLU:HG3	2.36	0.56
2:B:155:ASN:O	2:B:158:SER:HB2	2.05	0.56
2:B:188:TRP:CD1	2:B:189:PRO:HA	2.40	0.56
1:L:115:VAL:HA	1:L:135:PHE:O	2.06	0.56
2:B:15:GLY:N	2:B:82(C):LEU:O	2.31	0.55
2:B:47:TRP:CZ2	2:B:49:GLY:HA2	2.41	0.55
2:H:4:LEU:HD22	2:H:34:MET:HE1	1.89	0.54
2:B:124:LEU:HB2	2:B:139:GLY:C	2.28	0.54
1:A:147:LYS:HG3	1:A:154:GLU:HG3	1.90	0.53
2:H:140:CYS:HB2	2:H:154:TRP:CH2	2.43	0.53
2:B:136:VAL:HG22	2:B:185:SER:HA	1.90	0.53
1:L:142:LYS:HG3	1:L:173:TYR:CD1	2.44	0.52
1:A:161:ASN:HD22	1:A:177:SER:HA	1.73	0.52
1:L:39:LYS:NZ	1:L:81:GLU:O	2.38	0.51
1:A:163:TRP:CD2	1:A:175:MET:HG3	2.47	0.50
2:H:155:ASN:O	2:H:158:SER:HB2	2.12	0.50
1:L:170:ASP:O	1:L:171:SER:HB2	2.12	0.49
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.46	0.49
1:L:150:ILE:HA	1:L:191:SER:O	2.12	0.49
2:H:124:LEU:HD12	2:H:140:CYS:N	2.28	0.49
1:A:29:ILE:HD11	1:A:71:PHE:CE1	2.48	0.49
2:B:154:TRP:CD1	2:B:163:VAL:HG21	2.47	0.48
1:L:13:ALA:HA	1:L:107:LYS:HD2	1.94	0.48
2:H:38:LYS:HB2	2:H:48:ILE:HD11	1.96	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:ASN:O	1:A:51:ALA:HB3	2.14	0.48
1:L:123:GLU:OE2	2:H:208:LYS:NZ	2.47	0.47
1:A:161:ASN:ND2	1:A:177:SER:OG	2.47	0.47
2:B:12:VAL:HG11	2:B:82(C):LEU:HD12	1.96	0.47
2:H:144:GLY:CA	2:H:174:LEU:HD23	2.44	0.47
2:H:196:ASN:HA	2:H:207:ASP:OD2	2.15	0.47
2:B:87:SER:O	2:B:88:ALA:HB2	2.15	0.47
2:H:188:TRP:CG	2:H:189:PRO:HA	2.49	0.47
1:A:151:ASP:HA	1:A:191:SER:HB3	1.97	0.47
1:A:160:LEU:HD11	2:B:171:GLN:HG3	1.96	0.46
2:B:184:PRO:HB2	2:B:187:THR:HG23	1.97	0.46
1:A:117:ILE:HD12	1:A:194:CYS:HB2	1.96	0.46
1:A:13:ALA:HB1	1:A:17:GLU:HG2	1.98	0.46
2:B:188:TRP:CG	2:B:189:PRO:HA	2.50	0.46
1:A:13:ALA:HA	1:A:107:LYS:HD2	1.98	0.46
1:A:116:SER:O	1:A:134:CYS:HA	2.16	0.46
1:A:125:LEU:HD22	1:A:183:LYS:HG3	1.97	0.46
1:A:142:LYS:HG3	1:A:173:TYR:CD1	2.50	0.46
2:B:138:LEU:HD13	2:B:210:ILE:HG21	1.97	0.45
2:H:171:GLN:NE2	2:H:176:THR:OG1	2.47	0.45
2:H:35:HIS:HA	2:H:50:TYR:HB3	1.98	0.44
1:L:118:PHE:HA	1:L:119:PRO:HD3	1.79	0.44
1:A:15:VAL:HG13	1:A:78:LEU:O	2.17	0.44
2:B:96:GLY:HA2	2:B:100(B):ALA:O	2.17	0.44
1:A:183:LYS:O	1:A:187:GLU:HG2	2.18	0.44
2:H:34:MET:HE2	2:H:92:CYS:CB	2.42	0.44
2:B:4:LEU:HD22	2:B:34:MET:CE	2.48	0.44
2:B:29:PHE:CD2	2:B:76:SER:HA	2.53	0.43
1:A:175:MET:HE1	1:A:177:SER:HB2	2.00	0.43
2:B:36:TRP:CD1	2:B:69:LEU:HD13	2.54	0.43
1:L:125:LEU:HD22	1:L:183:LYS:HG3	2.00	0.43
2:B:4:LEU:HD22	2:B:34:MET:HE2	2.00	0.43
2:B:124:LEU:HB2	2:B:139:GLY:HA3	2.01	0.43
2:B:211:VAL:HA	2:B:212:PRO:HD2	1.77	0.43
1:L:163:TRP:CG	1:L:175:MET:HG3	2.54	0.42
1:A:156:GLN:O	1:A:159:VAL:HG23	2.19	0.42
1:A:181:LEU:HD22	1:A:185:GLU:OE2	2.19	0.42
2:H:51:ILE:O	2:H:52(A):PRO:HD3	2.19	0.42
1:A:115:VAL:HA	1:A:135:PHE:O	2.19	0.42
1:A:160:LEU:HD13	2:B:169:VAL:CG2	2.49	0.42
2:H:10:GLU:HG3	2:H:18:VAL:HG21	2.02	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:175:MET:HE3	1:L:177:SER:HB2	2.02	0.42
1:A:175:MET:HE2	1:A:175:MET:HB3	1.81	0.42
2:B:70:THR:OG1	2:B:79:TYR:HB2	2.20	0.41
1:L:175:MET:HB3	1:L:175:MET:HE2	1.81	0.41
2:H:136:VAL:HG22	2:H:185:SER:HA	2.01	0.41
2:H:208:LYS:HD3	2:H:208:LYS:HA	1.84	0.41
2:H:119:PRO:CB	2:H:145:TYR:HB3	2.50	0.41
1:A:175:MET:HE3	1:A:176:SER:C	2.41	0.41
2:B:35:HIS:CD2	2:B:100(C):MET:HG2	2.56	0.41
2:B:146:PHE:HA	2:B:147:PRO:HA	1.91	0.41
1:A:89:GLN:HA	1:A:97:THR:O	2.21	0.41
2:H:14:PRO:HA	2:H:82(C):LEU:O	2.21	0.41
2:H:22:CYS:HB3	2:H:78:ALA:HB3	2.02	0.40
2:H:124:LEU:HB2	2:H:139:GLY:CA	2.52	0.40
2:B:119:PRO:HB3	2:B:145:TYR:HB3	2.04	0.40
1:A:193:THR:HG23	1:A:206:VAL:HG23	2.02	0.40
1:L:122:SER:HA	1:L:125:LEU:HD12	2.03	0.40
2:H:14:PRO:HD3	2:H:112:SER:C	2.41	0.40
1:A:61:ARG:HG2	1:A:61:ARG:NH2	2.36	0.40
1:A:62:PHE:CD1	1:A:75:ILE:HG12	2.56	0.40
1:A:175:MET:CE	1:A:177:SER:HB2	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/214 (99%)	204 (97%)	6 (3%)	1 (0%)	29	48
1	L	211/214 (99%)	194 (92%)	16 (8%)	1 (0%)	29	48
2	B	209/230 (91%)	195 (93%)	9 (4%)	5 (2%)	6	9
2	H	209/230 (91%)	192 (92%)	14 (7%)	3 (1%)	11	20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	840/888 (95%)	785 (94%)	45 (5%)	10 (1%)	13	24

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	98	ARG
2	H	99	PRO
2	B	97	TYR
2	B	98	ARG
2	B	99	PRO
2	B	100	TYR
1	L	30	TYR
2	B	100(A)	TYR
2	H	96	GLY
1	A	30	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/188 (94%)	165 (93%)	12 (7%)	16	30
1	L	177/188 (94%)	166 (94%)	11 (6%)	18	35
2	B	173/200 (86%)	158 (91%)	15 (9%)	10	20
2	H	171/200 (86%)	158 (92%)	13 (8%)	13	25
All	All	698/776 (90%)	647 (93%)	51 (7%)	14	27

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	10	SER
1	L	12	SER
1	L	30	TYR
1	L	33	LEU
1	L	175	MET

Continued on next page...

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	L	185	GLU
1	L	188	ARG
1	L	202	THR
1	L	207	LYS
1	L	208	SER
1	L	212	ASN
2	H	11	LEU
2	H	12	VAL
2	H	57	THR
2	H	76	SER
2	H	82(B)	SER
2	H	83	THR
2	H	112	SER
2	H	136	VAL
2	H	158	SER
2	H	177	LEU
2	H	192	THR
2	H	203	SER
2	H	208	LYS
1	A	17	GLU
1	A	33	LEU
1	A	56	GLU
1	A	70	GLN
1	A	127	SER
1	A	143	ASP
1	A	175	MET
1	A	185	GLU
1	A	202	THR
1	A	206	VAL
1	A	208	SER
1	A	212	ASN
2	B	11	LEU
2	B	51	ILE
2	B	57	THR
2	B	58	LYS
2	B	62	LYS
2	B	74	SER
2	B	82	LEU
2	B	83	THR
2	B	134	SER
2	B	136	VAL
2	B	158	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	163	VAL
2	B	177	LEU
2	B	192	THR
2	B	196	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	137	ASN
1	L	161	ASN
1	L	190	ASN
1	L	212	ASN
2	H	171	GLN
1	A	40	GLN
1	A	50	ASN
1	A	90	HIS
1	A	137	ASN
1	A	161	ASN
1	A	212	ASN
2	B	171	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	L	301	-	4,4,4	0.13	0	6,6,6	0.24	0
3	SO4	L	302	-	4,4,4	1.14	0	6,6,6	0.51	0
3	SO4	A	303	-	4,4,4	0.12	0	6,6,6	0.20	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	213/214 (99%)	-0.33	0 100 100	17, 24, 31, 35	0
1	L	213/214 (99%)	-0.39	0 100 100	17, 24, 30, 34	0
2	B	209/230 (90%)	-0.40	0 100 100	17, 24, 29, 39	0
2	H	209/230 (90%)	-0.46	0 100 100	17, 24, 29, 39	0
All	All	844/888 (95%)	-0.39	0 100 100	17, 24, 30, 39	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	L	301	5/5	0.98	0.14	46,46,47,47	5
3	SO4	A	303	5/5	0.98	0.10	44,44,44,44	5
3	SO4	L	302	5/5	0.99	0.10	37,37,38,39	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.