



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 15, 2023 – 02:45 PM EDT

PDB ID : 1Y3A  
Title : Structure of G-Alpha-I1 bound to a GDP-selective peptide provides insight into guanine nucleotide exchange  
Authors : Johnston, C.A.; Willard, F.S.; Jezyk, M.R.; Fredericks, Z.; Bodor, E.T.; Jones, M.B.; Blaesius, R.; Harden, T.K.; Sondek, J.; Watts, V.J.; Ramer, J.K.; Siderovski, D.P.  
Deposited on : 2004-11-24  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">4%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey); position: relative;"> <span style="position: absolute; top: 50%; left: 0; right: 0; transform: translate(-50%, -50%); font-size: 8px;">71% 19% 9%</span> </div> </div>
1	B	329	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">5%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey); position: relative;"> <span style="position: absolute; top: 50%; left: 0; right: 0; transform: translate(-50%, -50%); font-size: 8px;">73% 19% 7%</span> </div> </div>
1	C	329	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">3%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey); position: relative;"> <span style="position: absolute; top: 50%; left: 0; right: 0; transform: translate(-50%, -50%); font-size: 8px;">70% 19% 9%</span> </div> </div>
1	D	329	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">5%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey); position: relative;"> <span style="position: absolute; top: 50%; left: 0; right: 0; transform: translate(-50%, -50%); font-size: 8px;">70% 20% 8%</span> </div> </div>
2	E	16	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">12%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey); position: relative;"> <span style="position: absolute; top: 50%; left: 0; right: 0; transform: translate(-50%, -50%); font-size: 8px;">31% 44% 25%</span> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	16	 50% 25% 25%
2	G	16	 56% 19% 25%
2	H	16	 12% 44% 31% 25%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10424 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Guanine nucleotide-binding protein G(i), alpha-1 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	299	Total 2413	C 1540	N 407	O 451	S 15	0	0	0
1	B	307	Total 2471	C 1575	N 414	O 467	S 15	0	0	0
1	C	298	Total 2404	C 1534	N 405	O 450	S 15	0	0	0
1	D	304	Total 2452	C 1562	N 411	O 464	S 15	0	0	0

- Molecule 2 is a protein called KB752 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	12	Total 109	C 71	N 16	O 21	S 1	0	0	0
2	F	12	Total 109	C 71	N 16	O 21	S 1	0	0	0
2	G	12	Total 109	C 71	N 16	O 21	S 1	0	0	0
2	H	12	Total 109	C 71	N 16	O 21	S 1	0	0	0

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:  $C_{10}H_{15}N_5O_{11}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	C	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	D	1	Total	C	N	O	P	0	0
			28	10	5	11	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	46	Total	O	0	0
			46	46		
4	B	29	Total	O	0	0
			29	29		
4	C	30	Total	O	0	0
			30	30		
4	D	25	Total	O	0	0
			25	25		
4	E	2	Total	O	0	0
			2	2		
4	F	1	Total	O	0	0
			1	1		
4	G	2	Total	O	0	0
			2	2		
4	H	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Guanine nucleotide-binding protein G(i), alpha-1 subunit




- Molecule 1: Guanine nucleotide-binding protein G(i), alpha-1 subunit




- Molecule 1: Guanine nucleotide-binding protein G(i), alpha-1 subunit







● Molecule 1: Guanine nucleotide-binding protein G(i), alpha-1 subunit




● Molecule 2: KB752 peptide




● Molecule 2: KB752 peptide



● Molecule 2: KB752 peptide



● Molecule 2: KB752 peptide





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.94Å 112.78Å 109.49Å 90.00° 93.75° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 37.59 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.50) 96.1 (37.59-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.49 (at 2.51Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.240 , 0.280 0.244 , 0.250	Depositor DCC
$R_{free}$ test set	3001 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.0	Xtrriage
Anisotropy	0.107	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 42.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	10424	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	1/2456 (0.0%)	0.59	0/3306
1	B	0.45	1/2516 (0.0%)	0.61	0/3389
1	C	0.43	0/2447	0.60	0/3295
1	D	0.43	1/2496 (0.0%)	0.60	1/3363 (0.0%)
2	E	0.35	0/112	0.51	0/151
2	F	0.41	0/112	0.51	0/151
2	G	0.37	0/112	0.48	0/151
2	H	0.38	0/112	0.50	0/151
All	All	0.43	3/10363 (0.0%)	0.60	1/13957 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	60	GLY	N-CA	5.80	1.54	1.46
1	B	241	ASN	N-CA	5.66	1.57	1.46
1	D	241	ASN	N-CA	5.52	1.57	1.46

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	241	ASN	N-CA-C	-7.67	90.29	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2413	0	2411	52	0
1	B	2471	0	2452	43	0
1	C	2404	0	2398	54	1
1	D	2452	0	2437	58	1
2	E	109	0	98	8	0
2	F	109	0	98	5	0
2	G	109	0	98	4	0
2	H	109	0	98	4	0
3	A	28	0	12	2	0
3	B	28	0	12	2	0
3	C	28	0	12	3	0
3	D	28	0	12	3	0
4	A	46	0	0	8	0
4	B	29	0	0	2	0
4	C	30	0	0	3	0
4	D	25	0	0	1	0
4	E	2	0	0	0	0
4	F	1	0	0	0	0
4	G	2	0	0	1	0
4	H	1	0	0	0	0
All	All	10424	0	10138	216	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 216 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:208:ARG:H	1:D:208:ARG:HD2	1.15	1.09
1:D:55:ILE:HA	1:D:60:GLY:HA2	1.39	1.03
1:A:55:ILE:HA	1:A:60:GLY:HA2	1.38	1.03
1:C:55:ILE:HA	1:C:60:GLY:HA2	1.39	1.03
1:B:55:ILE:HA	1:B:60:GLY:HA2	1.38	1.02

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:63:GLU:OE1	1:D:257:LYS:NZ[2_656]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/329 (89%)	291 (99%)	1 (0%)	1 (0%)	41	61
1	B	303/329 (92%)	297 (98%)	2 (1%)	4 (1%)	12	21
1	C	292/329 (89%)	288 (99%)	2 (1%)	2 (1%)	22	39
1	D	300/329 (91%)	292 (97%)	6 (2%)	2 (1%)	22	39
2	E	10/16 (62%)	10 (100%)	0	0	100	100
2	F	10/16 (62%)	10 (100%)	0	0	100	100
2	G	10/16 (62%)	10 (100%)	0	0	100	100
2	H	10/16 (62%)	9 (90%)	1 (10%)	0	100	100
All	All	1228/1380 (89%)	1207 (98%)	12 (1%)	9 (1%)	22	39

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	241	ASN
1	C	241	ASN
1	D	235	ALA
1	A	59	ALA
1	B	59	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/284 (93%)	254 (97%)	9 (3%)	37	63
1	B	268/284 (94%)	258 (96%)	10 (4%)	34	60
1	C	262/284 (92%)	256 (98%)	6 (2%)	50	76
1	D	267/284 (94%)	259 (97%)	8 (3%)	41	68
2	E	12/16 (75%)	11 (92%)	1 (8%)	11	22
2	F	12/16 (75%)	12 (100%)	0	100	100
2	G	12/16 (75%)	12 (100%)	0	100	100
2	H	12/16 (75%)	11 (92%)	1 (8%)	11	22
All	All	1108/1200 (92%)	1073 (97%)	35 (3%)	39	65

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	208	ARG
1	D	239	GLU
1	D	329	THR
1	B	193	ASP
1	B	159	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	204	GLN
1	D	79	GLN
1	C	333	GLN
1	D	147	GLN
1	B	79	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GDP	D	358	-	24,30,30	1.61	6 (25%)	30,47,47	2.22	9 (30%)
3	GDP	A	355	-	24,30,30	1.61	5 (20%)	30,47,47	2.21	9 (30%)
3	GDP	B	356	-	24,30,30	1.72	7 (29%)	30,47,47	2.20	9 (30%)
3	GDP	C	357	-	24,30,30	1.54	4 (16%)	30,47,47	2.24	8 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	D	358	-	-	3/12/32/32	0/3/3/3
3	GDP	A	355	-	-	3/12/32/32	0/3/3/3
3	GDP	B	356	-	-	3/12/32/32	0/3/3/3
3	GDP	C	357	-	-	3/12/32/32	0/3/3/3

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	356	GDP	C6-N1	4.00	1.43	1.37
3	D	358	GDP	C6-N1	3.34	1.42	1.37
3	C	357	GDP	C6-N1	3.15	1.42	1.37
3	A	355	GDP	C6-N1	3.09	1.42	1.37
3	D	358	GDP	C8-N7	-2.96	1.30	1.35

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	357	GDP	O5'-PA-O1A	-7.57	79.48	109.07
3	B	356	GDP	O5'-PA-O1A	-7.50	79.75	109.07
3	D	358	GDP	O5'-PA-O1A	-7.49	79.80	109.07
3	A	355	GDP	O5'-PA-O1A	-7.44	80.01	109.07
3	C	357	GDP	O3B-PB-O3A	4.65	120.24	104.64

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	355	GDP	O4'-C4'-C5'-O5'
3	B	356	GDP	O4'-C4'-C5'-O5'
3	C	357	GDP	O4'-C4'-C5'-O5'
3	D	358	GDP	O4'-C4'-C5'-O5'
3	A	355	GDP	C3'-C4'-C5'-O5'

There are no ring outliers.


4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	358	GDP	3	0
3	A	355	GDP	2	0
3	B	356	GDP	2	0
3	C	357	GDP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	299/329 (90%)	0.28	14 (4%) 31 33	19, 40, 85, 154	0
1	B	307/329 (93%)	0.18	18 (5%) 22 23	16, 37, 78, 136	0
1	C	298/329 (90%)	0.15	10 (3%) 45 48	21, 38, 80, 104	0
1	D	304/329 (92%)	0.31	16 (5%) 26 28	22, 43, 96, 171	0
2	E	12/16 (75%)	1.20	2 (16%) 1 1	43, 64, 84, 142	0
2	F	12/16 (75%)	-0.04	0 100 100	24, 44, 69, 110	0
2	G	12/16 (75%)	0.22	0 100 100	34, 48, 91, 130	0
2	H	12/16 (75%)	0.99	2 (16%) 1 1	43, 66, 100, 148	0
All	All	1256/1380 (91%)	0.25	62 (4%) 29 31	16, 40, 85, 171	0

The worst 5 of 62 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	257	LYS	6.2
1	D	258	TRP	5.5
1	C	59	ALA	4.8
1	A	345	LYS	4.7
2	E	12	ASP	4.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GDP	C	357	28/28	0.93	0.18	25,31,58,58	0
3	GDP	B	356	28/28	0.94	0.18	23,30,48,51	0
3	GDP	A	355	28/28	0.94	0.18	23,31,49,50	0
3	GDP	D	358	28/28	0.95	0.17	28,35,56,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around GDP B 356:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GDP A 355:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.