

wwPDB X-ray Structure Validation Summary Report (i)

Aug 16, 2023 – 03:47 AM EDT

PDB ID : 1Y7O

Title: The structure of Streptococcus pneumoniae A153P ClpP

Authors: Kimber, M.S.; Gribun, A.; Ching, R.; Sprangers, R.; Fiebig, K.M.; Houry,

W.A.

Deposited on : 2004-12-09

Resolution : 2.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
-		010	2%					
1	A	218	64%	12% • 22%				
1	D	010	2%					
1	В	218	67%	11% • 18%				
		210	.% ■					
1	С	218	67%	13% • 19%				
	_		2%					
1	D	218	67%	12% • 18%				
			3%					
1	Е	218	62%	15% • 20%				



Continued from previous page...

Mol	Chain	Length	Quality of chain					
1	F	218	61%	16%	•	21%		
1	G	218	66%	14%		19%		



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	A	171	Total C N O Se 1315 828 217 258 12	0	0	0
1	В	178	Total C N O S Se 1373 862 228 270 1 12	0	0	0
1	С	176	Total C N O S Se 1357 853 226 265 1 12	0	0	0
1	D	178	Total C N O S Se 1371 859 228 271 1 12	0	0	0
1	Е	174	Total C N O S Se 1340 842 223 262 1 12	0	0	0
1	F	173	Total C N O S Se 1328 837 219 259 1 12	0	0	0
1	G	177	Total C N O Se 1365 857 227 269 12	0	0	0

There are 245 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	cloning artifact	UNP P63788
A	-18	GLY	-	cloning artifact	UNP P63788
A	-17	SER	-	cloning artifact	UNP P63788
A	-16	SER	-	cloning artifact	UNP P63788
A	-15	HIS	-	cloning artifact	UNP P63788
A	-14	HIS	-	cloning artifact	UNP P63788
A	-13	HIS	-	cloning artifact	UNP P63788
A	-12	HIS	-	cloning artifact	UNP P63788
A	-11	HIS	-	cloning artifact	UNP P63788
A	-10	HIS	-	cloning artifact	UNP P63788
A	-9	SER	-	cloning artifact	UNP P63788
A	-8	SER	-	cloning artifact	UNP P63788
A	-7	GLY	-	cloning artifact	UNP P63788
A	-6	LEU	-	cloning artifact	UNP P63788
A	-5	VAL	-	cloning artifact	UNP P63788



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	PRO	-	cloning artifact	UNP P63788
A	-3	ARG	-	cloning artifact	UNP P63788
A	-2	GLY	-	cloning artifact	UNP P63788
A	-1	SER	-	cloning artifact	UNP P63788
A	0	HIS	-	cloning artifact	UNP P63788
A	29	MSE	MET	modified residue	UNP P63788
A	38	MSE	MET	modified residue	UNP P63788
A	79	MSE	MET	modified residue	UNP P63788
A	91	MSE	MET	modified residue	UNP P63788
A	93	MSE	MET	modified residue	UNP P63788
A	97	MSE	MET	modified residue	UNP P63788
A	112	MSE	MET	modified residue	UNP P63788
A	119	MSE	MET	modified residue	UNP P63788
A	124	MSE	MET	modified residue	UNP P63788
A	140	PRO	ALA	engineered mutation	UNP P63788
A	162	MSE	MET	modified residue	UNP P63788
A	175	MSE	MET	modified residue	UNP P63788
A	190	MSE	MET	modified residue	UNP P63788
A	197	GLY	-	cloning artifact	UNP P63788
A	198	SER	-	cloning artifact	UNP P63788
В	-19	MET	-	cloning artifact	UNP P63788
В	-18	GLY	-	cloning artifact	UNP P63788
В	-17	SER	-	cloning artifact	UNP P63788
В	-16	SER	-	cloning artifact	UNP P63788
В	-15	HIS	-	cloning artifact	UNP P63788
В	-14	HIS	-	cloning artifact	UNP P63788
В	-13	HIS	-	cloning artifact	UNP P63788
В	-12	HIS	-	cloning artifact	UNP P63788
В	-11	HIS	-	cloning artifact	UNP P63788
В	-10	HIS	-	cloning artifact	UNP P63788
В	-9	SER	-	cloning artifact	UNP P63788
В	-8	SER	-	cloning artifact	UNP P63788
В	-7	GLY	-	cloning artifact	UNP P63788
В	-6	LEU	-	cloning artifact	UNP P63788
В	-5	VAL	-	cloning artifact	UNP P63788
В	-4	PRO	-	cloning artifact	UNP P63788
В	-3	ARG	-	cloning artifact	UNP P63788
В	-2	GLY	-	cloning artifact	UNP P63788
В	-1	SER	-	cloning artifact	UNP P63788
В	0	HIS	-	cloning artifact	UNP P63788
В	29	MSE	MET	modified residue	UNP P63788
В	38	MSE	MET	modified residue	UNP P63788



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
В	79	MSE	MET	modified residue	UNP P63788
В	91	MSE	MET	modified residue	UNP P63788
В	93	MSE	MET	modified residue	UNP P63788
В	97	MSE	MET	modified residue	UNP P63788
В	112	MSE	MET	modified residue	UNP P63788
В	119	MSE	MET	modified residue	UNP P63788
В	124	MSE	MET	modified residue	UNP P63788
В	140	PRO	ALA	engineered mutation	UNP P63788
В	162	MSE	MET	modified residue	UNP P63788
В	175	MSE	MET	modified residue	UNP P63788
В	190	MSE	MET	modified residue	UNP P63788
В	197	GLY	-	cloning artifact	UNP P63788
В	198	SER	_	cloning artifact	UNP P63788
С	-19	MET	-	cloning artifact	UNP P63788
С	-18	GLY	-	cloning artifact	UNP P63788
С	-17	SER	_	cloning artifact	UNP P63788
С	-16	SER	-	cloning artifact	UNP P63788
С	-15	HIS	-	cloning artifact	UNP P63788
С	-14	HIS	-	cloning artifact	UNP P63788
С	-13	HIS	-	cloning artifact	UNP P63788
С	-12	HIS	-	cloning artifact	UNP P63788
С	-11	HIS	-	cloning artifact	UNP P63788
С	-10	HIS	-	cloning artifact	UNP P63788
С	-9	SER	-	cloning artifact	UNP P63788
С	-8	SER	-	cloning artifact	UNP P63788
С	-7	GLY	-	cloning artifact	UNP P63788
С	-6	LEU	-	cloning artifact	UNP P63788
С	-5	VAL	-	cloning artifact	UNP P63788
С	-4	PRO	-	cloning artifact	UNP P63788
С	-3	ARG	-	cloning artifact	UNP P63788
С	-2	GLY	-	cloning artifact	UNP P63788
С	-1	SER		cloning artifact	UNP P63788
С	0	HIS	-	cloning artifact	UNP P63788
С	29	MSE	MET	modified residue	UNP P63788
С	38	MSE	MET	modified residue	UNP P63788
С	79	MSE	MET	modified residue	UNP P63788
С	91	MSE	MET	modified residue	UNP P63788
С	93	MSE	MET	modified residue	UNP P63788
С	97	MSE	MET	modified residue	UNP P63788
С	112	MSE	MET	modified residue	UNP P63788
С	119	MSE	MET	modified residue	UNP P63788
С	124	MSE	MET	modified residue	UNP P63788



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
С	140	PRO	ALA	engineered mutation	UNP P63788
С	162	MSE	MET	modified residue	UNP P63788
С	175	MSE	MET	modified residue	UNP P63788
С	190	MSE	MET	modified residue	UNP P63788
С	197	GLY	_	cloning artifact	UNP P63788
С	198	SER	-	cloning artifact	UNP P63788
D	-19	MET	_	cloning artifact	UNP P63788
D	-18	GLY	-	cloning artifact	UNP P63788
D	-17	SER	-	cloning artifact	UNP P63788
D	-16	SER	-	cloning artifact	UNP P63788
D	-15	HIS	_	cloning artifact	UNP P63788
D	-14	HIS	_	cloning artifact	UNP P63788
D	-13	HIS	_	cloning artifact	UNP P63788
D	-12	HIS	_	cloning artifact	UNP P63788
D	-11	HIS	_	cloning artifact	UNP P63788
D	-10	HIS	_	cloning artifact	UNP P63788
D	-9	SER	_	cloning artifact	UNP P63788
D	-8	SER	-	cloning artifact	UNP P63788
D	-7	GLY	_	cloning artifact	UNP P63788
D	-6	LEU	-	cloning artifact	UNP P63788
D	-5	VAL	-	cloning artifact	UNP P63788
D	-4	PRO	-	cloning artifact	UNP P63788
D	-3	ARG	-	cloning artifact	UNP P63788
D	-2	GLY	-	cloning artifact	UNP P63788
D	-1	SER	-	cloning artifact	UNP P63788
D	0	HIS	-	cloning artifact	UNP P63788
D	29	MSE	MET	modified residue	UNP P63788
D	38	MSE	MET	modified residue	UNP P63788
D	79	MSE	MET	modified residue	UNP P63788
D	91	MSE	MET	modified residue	UNP P63788
D	93	MSE	MET	modified residue	UNP P63788
D	97	MSE	MET	modified residue	UNP P63788
D	112	MSE	MET	modified residue	UNP P63788
D	119	MSE	MET	modified residue	UNP P63788
D	124	MSE	MET	modified residue	UNP P63788
D	140	PRO	ALA	engineered mutation	UNP P63788
D	162	MSE	MET	modified residue	UNP P63788
D	175	MSE	MET	modified residue	UNP P63788
D	190	MSE	MET	modified residue	UNP P63788
D	197	GLY	-	cloning artifact	UNP P63788
D	198	SER	-	cloning artifact	UNP P63788
Е	-19	MET	-	cloning artifact	UNP P63788



 $Continued\ from\ previous\ page...$

Е			Actual	Comment	Reference
	-18	GLY	-	cloning artifact	UNP P63788
E	-17	SER	-	cloning artifact	UNP P63788
Е	-16	SER	-	cloning artifact	UNP P63788
Е	-15	HIS	-	cloning artifact	UNP P63788
Е	-14	HIS	-	cloning artifact	UNP P63788
Е	-13	HIS	-	cloning artifact	UNP P63788
Е	-12	HIS	-	cloning artifact	UNP P63788
Е	-11	HIS	-	cloning artifact	UNP P63788
Е	-10	HIS	-	cloning artifact	UNP P63788
Е	-9	SER	-	cloning artifact	UNP P63788
Е	-8	SER	-	cloning artifact	UNP P63788
Е	-7	GLY	-	cloning artifact	UNP P63788
Е	-6	LEU	-	cloning artifact	UNP P63788
Е	-5	VAL	-	cloning artifact	UNP P63788
Е	-4	PRO	-	cloning artifact	UNP P63788
Е	-3	ARG	-	cloning artifact	UNP P63788
Е	-2	GLY	-	cloning artifact	UNP P63788
Е	-1	SER	-	cloning artifact	UNP P63788
Е	0	HIS	-	cloning artifact	UNP P63788
Е	29	MSE	MET	modified residue	UNP P63788
Е	38	MSE	MET	modified residue	UNP P63788
Е	79	MSE	MET	modified residue	UNP P63788
Е	91	MSE	MET	modified residue	UNP P63788
Е	93	MSE	MET	modified residue	UNP P63788
Е	97	MSE	MET	modified residue	UNP P63788
Е	112	MSE	MET	modified residue	UNP P63788
Е	119	MSE	MET	modified residue	UNP P63788
Е	124	MSE	MET	modified residue	UNP P63788
Е	140	PRO	ALA	engineered mutation	UNP P63788
Е	162	MSE	MET	modified residue	UNP P63788
Е	175	MSE	MET	modified residue	UNP P63788
Е	190	MSE	MET	modified residue	UNP P63788
Е	197	GLY	-	cloning artifact	UNP P63788
Е	198	SER		cloning artifact	UNP P63788
F	-19	MET		cloning artifact	UNP P63788
F	-18	GLY	=	cloning artifact	UNP P63788
F	-17	SER	-	cloning artifact	UNP P63788
F	-16	SER	-	cloning artifact	UNP P63788
F	-15	HIS	=	cloning artifact	UNP P63788
F	-14	HIS	-	cloning artifact	UNP P63788
F	-13	HIS	-	cloning artifact	UNP P63788
F	-12	HIS	-	cloning artifact	UNP P63788



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
F	-11	HIS	-	cloning artifact	UNP P63788
F	-10	HIS	_	cloning artifact	UNP P63788
F	-9	SER	-	cloning artifact	UNP P63788
F	-8	SER	_	cloning artifact	UNP P63788
F	-7	GLY	_	cloning artifact	UNP P63788
F	-6	LEU	-	cloning artifact	UNP P63788
F	-5	VAL	-	cloning artifact	UNP P63788
F	-4	PRO	-	cloning artifact	UNP P63788
F	-3	ARG	-	cloning artifact	UNP P63788
F	-2	GLY	-	cloning artifact	UNP P63788
F	-1	SER	-	cloning artifact	UNP P63788
F	0	HIS	-	cloning artifact	UNP P63788
F	29	MSE	MET	modified residue	UNP P63788
F	38	MSE	MET	modified residue	UNP P63788
F	79	MSE	MET	modified residue	UNP P63788
F	91	MSE	MET	modified residue	UNP P63788
F	93	MSE	MET	modified residue	UNP P63788
F	97	MSE	MET	modified residue	UNP P63788
F	112	MSE	MET	modified residue	UNP P63788
F	119	MSE	MET	modified residue	UNP P63788
F	124	MSE	MET	modified residue	UNP P63788
F	140	PRO	ALA	engineered mutation	UNP P63788
F	162	MSE	MET	modified residue	UNP P63788
F	175	MSE	MET	modified residue	UNP P63788
F	190	MSE	MET	modified residue	UNP P63788
F	197	GLY	-	cloning artifact	UNP P63788
F	198	SER	-	cloning artifact	UNP P63788
G	-19	MET	-	cloning artifact	UNP P63788
G	-18	GLY	-	cloning artifact	UNP P63788
G	-17	SER	-	cloning artifact	UNP P63788
G	-16	SER	-	cloning artifact	UNP P63788
G	-15	HIS	-	cloning artifact	UNP P63788
G	-14	HIS	-	cloning artifact	UNP P63788
G	-13	HIS		cloning artifact	UNP P63788
G	-12	HIS		cloning artifact	UNP P63788
G	-11	HIS		cloning artifact	UNP P63788
G	-10	HIS		cloning artifact	UNP P63788
G	-9	SER		cloning artifact	UNP P63788
G	-8	SER	-	cloning artifact	UNP P63788
G	-7	GLY	-	cloning artifact	UNP P63788
G	-6	LEU		cloning artifact	UNP P63788
G	-5	VAL	-	cloning artifact	UNP P63788



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
G	-4	PRO	-	cloning artifact	UNP P63788
G	-3	ARG	-	cloning artifact	UNP P63788
G	-2	GLY	-	cloning artifact	UNP P63788
G	-1	SER	-	cloning artifact	UNP P63788
G	0	HIS	-	cloning artifact	UNP P63788
G	29	MSE	MET	modified residue	UNP P63788
G	38	MSE	MET	modified residue	UNP P63788
G	79	MSE	MET	modified residue	UNP P63788
G	91	MSE	MET	modified residue	UNP P63788
G	93	MSE	MET	modified residue	UNP P63788
G	97	MSE	MET	modified residue	UNP P63788
G	112	MSE	MET	modified residue	UNP P63788
G	119	MSE	MET	modified residue	UNP P63788
G	124	MSE	MET	modified residue	UNP P63788
G	140	PRO	ALA	engineered mutation	UNP P63788
G	162	MSE	MET	modified residue	UNP P63788
G	175	MSE	MET	modified residue	UNP P63788
G	190	MSE	MET	modified residue	UNP P63788
G	197	GLY	-	cloning artifact	UNP P63788
G	198	SER	-	cloning artifact	UNP P63788

 \bullet Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ca 1 1	0	0
2	В	1	Total Ca 1 1	0	0
2	С	2	Total Ca 2 2	0	0
2	E	1	Total Ca 1 1	0	0
2	F	2	Total Ca 2 2	0	0
2	G	1	Total Ca 1 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	28	Total O 28 28	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	30	Total O 30 30	0	0
3	С	28	Total O 28 28	0	0
3	D	17	Total O 17 17	0	0
3	E	13	Total O 13 13	0	0
3	F	11	Total O 11 11	0	0
3	G	17	Total O 17 17	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP-dependent Clp protease proteolytic subunit











• Molecule 1: ATP-dependent Clp protease proteolytic subunit



• Molecule 1: ATP-dependent Clp protease proteolytic subunit





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants	105.65Å 105.65Å 236.19Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.50 - 2.51	Depositor
resolution (A)	29.84 - 2.50	EDS
% Data completeness	89.0 (29.50-2.51)	Depositor
(in resolution range)	90.6 (29.84-2.50)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) > 1$	2.49 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.1.19, CNX 2000.1	Depositor
R, R_{free}	0.186 , 0.248	Depositor
it, it _{free}	0.194 , 0.251	DCC
R_{free} test set	4780 reflections (9.81%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 36.7	EDS
L-test for twinning ²	$< L > = 0.46, < L^2> = 0.29$	Xtriage
Estimated twinning fraction	0.043 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9601	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.81% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	В	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.58	0/1321	0.80	4/1764 (0.2%)
1	В	0.59	0/1379	0.82	6/1841 (0.3%)
1	С	0.61	0/1363	0.82	6/1819 (0.3%)
1	D	0.55	0/1377	0.79	4/1838 (0.2%)
1	Е	0.53	0/1346	0.77	5/1796 (0.3%)
1	F	0.60	0/1334	0.80	3/1781 (0.2%)
1	G	0.52	0/1371	0.78	4/1831 (0.2%)
All	All	0.57	0/9491	0.80	$32/12670 \ (0.3\%)$

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	С	0	1

There are no bond length outliers.

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
1	F	17	ASP	CB-CG-OD2	7.88	125.39	118.30
1	A	85	ASP	CB-CG-OD2	7.54	125.08	118.30
1	В	17	ASP	CB-CG-OD2	6.95	124.55	118.30
1	G	172	ASP	CB-CG-OD2	6.64	124.28	118.30
1	D	57	ASP	CB-CG-OD2	6.53	124.18	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	С	2	ILE	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1315	0	1311	19	0
1	В	1373	0	1373	15	0
1	С	1357	0	1360	12	0
1	D	1371	0	1367	10	0
1	Ε	1340	0	1341	20	0
1	F	1328	0	1333	20	0
1	G	1365	0	1361	24	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	2	0	0	0	0
2	Ε	1	0	0	0	0
2	F	2	0	0	0	0
2	G	1	0	0	0	0
3	A	28	0	0	0	0
3	В	30	0	0	3	0
3	С	28	0	0	1	0
3	D	17	0	0	0	0
3	Ε	13	0	0	0	0
3	F	11	0	0	0	0
3	G	17	0	0	1	0
All	All	9601	0	9446	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 104 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)	
1:F:76:VAL:HA	1:F:79:MSE:HE3	1.47	0.94	
1:B:26:ARG:HG2	3:B:1011:HOH:O	1.72	0.88	
1:B:26:ARG:NE	3:B:1011:HOH:O	2.02	0.86	



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:A:97:MSE:HE1	1:A:150:LEU:HD13	1.58	0.83
1:G:30:LEU:CD1	1:G:38:MSE:HE1	2.14	0.77

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	165/218 (76%)	161 (98%)	3 (2%)	1 (1%)	25	43
1	В	172/218 (79%)	164 (95%)	7 (4%)	1 (1%)	25	43
1	С	170/218 (78%)	163 (96%)	7 (4%)	0	100	100
1	D	172/218 (79%)	164 (95%)	7 (4%)	1 (1%)	25	43
1	E	168/218 (77%)	163 (97%)	5 (3%)	0	100	100
1	F	167/218 (77%)	157 (94%)	9 (5%)	1 (1%)	25	43
1	G	171/218 (78%)	165 (96%)	6 (4%)	0	100	100
All	All	1185/1526 (78%)	1137 (96%)	44 (4%)	4 (0%)	41	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	3	PRO
1	В	2	ILE
1	A	53	ASP
1	D	140	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	144/168 (86%)	132 (92%)	12 (8%)	11 22
1	В	151/168 (90%)	138 (91%)	13 (9%)	10 20
1	C	149/168 (89%)	142 (95%)	7 (5%)	26 49
1	D	151/168 (90%)	132 (87%)	19 (13%)	4 8
1	E	147/168 (88%)	131 (89%)	16 (11%)	6 12
1	F	146/168 (87%)	129 (88%)	17 (12%)	5 10
1	G	150/168 (89%)	139 (93%)	11 (7%)	14 27
All	All	1038/1176 (88%)	943 (91%)	95 (9%)	9 18

5 of 95 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	Е	73	LEU
1	F	31	THR
1	Е	103	SER
1	Е	172	ASP
1	F	107	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	173	ASN
1	G	173	ASN
1	G	40	ASN
1	Е	87	GLN
1	F	148	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ} {>} 2$		$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	159/218 (72%)	-0.03	5 (3%) 49 5	52	15, 27, 51, 60	0
1	В	166/218 (76%)	-0.11	5 (3%) 50 5	53	15, 25, 50, 59	0
1	С	164/218 (75%)	-0.08	3 (1%) 68 7	71	13, 22, 41, 53	0
1	D	166/218 (76%)	0.07	5 (3%) 50 5	53	21, 30, 52, 59	0
1	E	162/218 (74%)	0.16	7 (4%) 35 3	38	27, 36, 56, 66	0
1	F	161/218 (73%)	-0.06	5 (3%) 49 5	52	20, 31, 56, 69	0
1	G	165/218 (75%)	0.15	10 (6%) 21	22	19, 32, 58, 72	0
All	All	1143/1526 (74%)	0.01	40 (3%) 44	47	13, 29, 54, 72	0

The worst 5 of 40 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	9	THR	5.9
1	D	8	GLN	4.6
1	D	10	SER	4.5
1	D	9	THR	4.0
1	F	1	MET	3.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	CA	F	1003	1/1	0.71	0.09	61,61,61,61	0
2	CA	A	1001	1/1	0.86	0.10	60,60,60,60	0
2	CA	Ε	1007	1/1	0.88	0.13	60,60,60,60	0
2	CA	F	1008	1/1	0.89	0.18	59,59,59,59	0
2	CA	В	1004	1/1	0.96	0.06	52,52,52,52	0
2	CA	G	1005	1/1	0.97	0.04	53,53,53,53	0
2	CA	С	1002	1/1	0.98	0.18	56,56,56,56	0
2	CA	С	1006	1/1	0.98	0.20	35,35,35,35	0

6.5 Other polymers (i)

There are no such residues in this entry.

