



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 4, 2023 – 07:12 am GMT

PDB ID : 2CK3  
Title : Azide inhibited bovine F1-ATPase  
Authors : Bowler, M.W.; Montgomery, M.G.; Leslie, A.G.W.; Walker, J.E.  
Deposited on : 2006-04-10  
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	510	<div style="display: flex; align-items: center;"> <div style="width: 12%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 85%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>
1	B	510	<div style="display: flex; align-items: center;"> <div style="width: 14%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 88%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div>
1	C	510	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>
2	D	482	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 87%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>
2	E	482	<div style="display: flex; align-items: center;"> <div style="width: 18%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>
2	F	482	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 90%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>
3	G	272	<div style="display: flex; align-items: center;"> <div style="width: 25%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 61%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 33%; height: 10px; background-color: grey;"></div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	H	146	
5	I	50	

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 26509 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP SYNTHASE SUBUNIT ALPHA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	487	3738	2359	658	709	12	0	4	0
1	B	480	3667	2310	649	696	12	0	1	0
1	C	490	3735	2353	659	711	12	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	481	GLY	SER	cloning artifact	UNP P19483
B	481	GLY	SER	cloning artifact	UNP P19483
C	481	GLY	SER	cloning artifact	UNP P19483

- Molecule 2 is a protein called ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	467	3550	2252	602	685	11	0	2	0
2	E	458	3486	2209	596	671	10	0	2	0
2	F	466	3541	2248	600	682	11	0	3	0

- Molecule 3 is a protein called ATP SYNTHASE SUBUNIT GAMMA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	G	182	1397	881	250	260	6	0	0	1


- Molecule 4 is a protein called ATP SYNTHASE SUBUNIT DELTA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	H	83	Total 620	C 391	N 102	O 126	S 1	0	0	0

- Molecule 5 is a protein called ATP SYNTHASE SUBUNIT EPSILON, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	I	25	Total 203	C 130	N 38	O 34	S 1	0	0	0

- Molecule 6 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
6	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
6	B	1	Total 31	C 10	N 6	O 12	P 3	0	0
6	C	1	Total 31	C 10	N 6	O 12	P 3	0	0
6	F	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
7	A	1	Total 1	Mg 1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total Mg 1 1	0	0
7	C	1	Total Mg 1 1	0	0
7	D	1	Total Mg 1 1	0	0
7	F	1	Total Mg 1 1	0	0

- Molecule 8 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	D	1	Total C N O P 27 10 5 10 2	0	0

- Molecule 9 is AZIDE ION (three-letter code: AZI) (formula:  $N_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	D	1	Total N 3 3	0	0

- Molecule 10 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	E	1	Total O P 5 4 1	0	0

- Molecule 11 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	380	Total O 380 380	0	0
11	B	409	Total O 409 409	0	0
11	C	378	Total O 378 378	0	0
11	D	383	Total O 383 383	0	0
11	E	246	Total O 246 246	0	0
11	F	484	Total O 484 484	0	0
11	G	113	Total O 113 113	0	0
11	H	8	Total O 8 8	0	0
11	I	7	Total O 7 7	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP SYNTHASE SUBUNIT ALPHA, MITOCHONDRIAL




- Molecule 1: ATP SYNTHASE SUBUNIT ALPHA, MITOCHONDRIAL





- Molecule 1: ATP SYNTHASE SUBUNIT ALPHA, MITOCHONDRIAL









• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL




• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL




• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL








● Molecule 3: ATP SYNTHASE SUBUNIT GAMMA, MITOCHONDRIAL



● Molecule 4: ATP SYNTHASE SUBUNIT DELTA, MITOCHONDRIAL



● Molecule 5: ATP SYNTHASE SUBUNIT EPSILON, MITOCHONDRIAL



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	261.16Å 105.22Å 122.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 43.53 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.00-1.95) 98.4 (43.53-1.95)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.76 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.197 , 0.226 0.208 , 0.234	Depositor DCC
$R_{free}$ test set	12057 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.0	Xtrriage
Anisotropy	0.258	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 61.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	26509	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AZI, PO4, ADP, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	2/3798 (0.1%)	0.55	0/5123
1	B	0.38	0/3718	0.52	0/5016
1	C	0.35	0/3775	0.51	0/5095
2	D	0.36	0/3611	0.55	0/4901
2	E	0.35	0/3547	0.52	0/4811
2	F	0.39	0/3607	0.54	0/4895
3	G	0.38	0/1406	0.47	0/1880
4	H	0.39	0/623	0.62	0/840
5	I	0.41	0/207	0.59	0/279
All	All	0.37	2/24292 (0.0%)	0.53	0/32840

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	422[A]	VAL	CA-CB	5.47	1.66	1.54
1	A	422[B]	VAL	CA-CB	5.47	1.66	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3738	0	3850	43	0
1	B	3667	0	3776	22	0
1	C	3735	0	3835	32	0
2	D	3550	0	3608	43	0
2	E	3486	0	3542	44	0
2	F	3541	0	3609	31	0
3	G	1397	0	1481	16	0
4	H	620	0	614	27	0
5	I	203	0	205	3	0
6	A	31	0	13	0	0
6	B	31	0	13	0	0
6	C	31	0	13	0	0
6	F	31	0	13	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
7	F	1	0	0	0	0
8	D	27	0	12	1	0
9	D	3	0	0	1	0
10	E	5	0	0	0	0
11	A	380	0	0	3	0
11	B	409	0	0	4	0
11	C	378	0	0	6	0
11	D	383	0	0	2	0
11	E	246	0	0	3	0
11	F	484	0	0	6	0
11	G	113	0	0	3	0
11	H	8	0	0	0	0
11	I	7	0	0	0	0
All	All	26509	0	24584	253	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 253 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:122:ALA:HA	4:H:123:ALA:CB	1.74	1.18
3:G:68:ILE:HB	3:G:69:ILE:HA	1.30	1.13
2:D:89:GLU:HG3	2:D:110:THR:HB	1.29	1.09
2:E:256:ASP:HA	2:E:257[B]:ASN:HB3	1.35	1.09

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:104:ASP:HA	4:H:105:LEU:HB2	1.31	1.09

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

#### 5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

#### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 5 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	ANP	F	600	7	29,33,33	1.81	8 (27%)	31,52,52	1.96	5 (16%)
8	ADP	D	600	7	24,29,29	1.01	2 (8%)	29,45,45	1.16	2 (6%)
9	AZI	D	1092	-	0,2,2	-	-	0,1,1	-	-
6	ANP	A	600	7	29,33,33	1.80	9 (31%)	31,52,52	1.94	5 (16%)
6	ANP	C	600	7	29,33,33	1.74	6 (20%)	31,52,52	2.01	7 (22%)
10	PO4	E	602	-	4,4,4	0.90	0	6,6,6	0.50	0
6	ANP	B	600	7	29,33,33	1.78	6 (20%)	31,52,52	1.78	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ANP	F	600	7	-	3/14/38/38	0/3/3/3
8	ADP	D	600	7	-	2/12/32/32	0/3/3/3
6	ANP	A	600	7	-	3/14/38/38	0/3/3/3
6	ANP	C	600	7	-	3/14/38/38	0/3/3/3
6	ANP	B	600	7	-	2/14/38/38	0/3/3/3

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	600	ANP	PB-N3B	4.27	1.74	1.63
6	B	600	ANP	PG-N3B	4.20	1.74	1.63
6	F	600	ANP	PB-N3B	4.18	1.74	1.63
6	C	600	ANP	PG-N3B	4.17	1.74	1.63
6	A	600	ANP	PB-N3B	4.17	1.74	1.63

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	F	600	ANP	O1G-PG-N3B	-7.75	100.36	111.77
6	C	600	ANP	O1G-PG-N3B	-7.67	100.47	111.77
6	A	600	ANP	O1G-PG-N3B	-7.49	100.75	111.77
6	B	600	ANP	O1G-PG-N3B	-6.10	102.79	111.77
6	B	600	ANP	O2B-PB-O1B	3.73	117.74	109.92

There are no chirality outliers.

5 of 13 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
6	A	600	ANP	PB-N3B-PG-O1G
6	A	600	ANP	PG-N3B-PB-O1B
6	B	600	ANP	PG-N3B-PB-O1B
6	B	600	ANP	PG-N3B-PB-O3A
6	C	600	ANP	PB-N3B-PG-O1G


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	D	600	ADP	1	0
9	D	1092	AZI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	487/510 (95%)	0.84	59 (12%) 4 7	12, 21, 45, 63	0
1	B	480/510 (94%)	0.83	72 (15%) 2 3	10, 20, 57, 82	0
1	C	490/510 (96%)	0.63	32 (6%) 18 27	12, 21, 38, 75	0
2	D	467/482 (96%)	0.55	26 (5%) 24 33	12, 19, 41, 67	0
2	E	458/482 (95%)	1.06	86 (18%) 1 1	12, 24, 64, 77	0
2	F	466/482 (96%)	0.45	26 (5%) 24 33	13, 19, 41, 62	0
3	G	182/272 (66%)	1.90	69 (37%) 0 0	10, 29, 48, 58	0
4	H	83/146 (56%)	2.75	51 (61%) 0 0	5, 26, 44, 49	0
5	I	25/50 (50%)	2.29	9 (36%) 0 0	8, 17, 43, 46	0
All	All	3138/3444 (91%)	0.86	430 (13%) 3 4	5, 21, 50, 82	0

The worst 5 of 430 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	21	THR	12.0
3	G	106	ILE	11.7
1	B	510	ALA	10.7
4	H	105	LEU	10.0
2	E	470	ALA	9.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
10	PO4	E	602	5/5	0.88	0.17	68,69,72,72	0
9	AZI	D	1092	3/3	0.92	0.16	17,17,19,25	0
7	MG	A	601	1/1	0.96	0.10	14,14,14,14	0
7	MG	B	601	1/1	0.96	0.09	19,19,19,19	0
6	ANP	B	600	31/31	0.97	0.10	13,19,27,32	0
6	ANP	C	600	31/31	0.98	0.11	12,16,19,19	0
7	MG	C	601	1/1	0.98	0.11	17,17,17,17	0
7	MG	D	601	1/1	0.98	0.07	16,16,16,16	0
7	MG	F	601	1/1	0.98	0.13	13,13,13,13	0
8	ADP	D	600	27/27	0.98	0.13	10,17,21,22	0
6	ANP	F	600	31/31	0.98	0.12	12,16,20,22	0
6	ANP	A	600	31/31	0.98	0.14	11,15,20,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ANP B 600:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around ANP C 600:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ADP D 600:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.