

Full wwPDB X-ray Structure Validation Report (i)

Mar 23, 2024 – 11:47 PM EDT

PDB ID : 2DVQ

Title: Crystal structure analysis of the N-terminal bromodomain of human BRD2

complexed with acetylated histone H4 peptide

Authors: Nakamura, Y.; Umehara, T.; Shirouzu, M.; Padmanabhan, B.; Yokoyama, S.;

RIKEN Structural Genomics/Proteomics Initiative (RSGI)

Deposited on : 2006-08-01

Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive | Similar resolution |
|-----------------------|-------------------------|--|
| Metric | $(\# \mathrm{Entries})$ | $(\# 	ext{Entries}, 	ext{ resolution range}(ext{Å}))$ |
| Clashscore | 141614 | 1773 (2.04-2.04) |
| Ramachandran outliers | 138981 | 1752 (2.04-2.04) |
| Sidechain outliers | 138945 | 1752 (2.04-2.04) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

| Mol | Chain | Length | Quality of chain | | | | |
|-----|-------|--------|------------------|-----|-----|-----|-------|
| 1 | A | 122 | | 69% | | 21% | • 7% |
| 1 | В | 122 | | 68% | | 19% | 5% 8% |
| 1 | С | 122 | | 69% | | 17% | • 12% |
| 2 | Р | 15 | 20% | 33% | | 47% | |
| 2 | Q | 15 | 33% | | 47% | | 20% |



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Bromodomain-containing protein 2.

| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf | Trace | | | |
|-----|-------|----------|-------|-----|-----|---------|---------|-------|---|---|---|
| 1 | Λ | 114 | Total | С | N | О | S | Se | 0 | 0 | 0 |
| 1 | A | 114 | 953 | 615 | 161 | 168 | 2 | 7 | 0 | 0 | 0 |
| 1 | D | 112 | Total | С | N | О | S | Se | 0 | 0 | 0 |
| 1 | Б | 112 | 935 | 605 | 158 | 163 | 2 | 7 | 0 | U | U |
| 1 | С | 107 | Total | С | N | О | S | Se | 0 | 0 | 0 |
| 1 | | 107 | 895 | 582 | 150 | 154 | 2 | 7 | Ü | 0 | U |

There are 21 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| A | 87 | MSE | MET | modified residue | UNP P25440 |
| A | 121 | MSE | MET | modified residue | UNP P25440 |
| A | 123 | MSE | MET | modified residue | UNP P25440 |
| A | 142 | MSE | MET | modified residue | UNP P25440 |
| A | 148 | MSE | MET | modified residue | UNP P25440 |
| A | 165 | MSE | MET | modified residue | UNP P25440 |
| A | 180 | MSE | MET | modified residue | UNP P25440 |
| В | 87 | MSE | MET | modified residue | UNP P25440 |
| В | 121 | MSE | MET | modified residue | UNP P25440 |
| В | 123 | MSE | MET | modified residue | UNP P25440 |
| В | 142 | MSE | MET | modified residue | UNP P25440 |
| В | 148 | MSE | MET | modified residue | UNP P25440 |
| В | 165 | MSE | MET | modified residue | UNP P25440 |
| В | 180 | MSE | MET | modified residue | UNP P25440 |
| С | 87 | MSE | MET | modified residue | UNP P25440 |
| С | 121 | MSE | MET | modified residue | UNP P25440 |
| С | 123 | MSE | MET | modified residue | UNP P25440 |
| С | 142 | MSE | MET | modified residue | UNP P25440 |
| С | 148 | MSE | MET | modified residue | UNP P25440 |
| С | 165 | MSE | MET | modified residue | UNP P25440 |
| С | 180 | MSE | MET | modified residue | UNP P25440 |

• Molecule 2 is a protein called histone H4.



| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf | Trace |
|-----|-------|----------|----------------------------|---------|---------|-------|
| 2 | Р | 8 | Total C N O 46 28 9 9 | 0 | 0 | 0 |
| 2 | Q | 15 | Total C N O 82 48 17 17 | 0 | 0 | 0 |

• Molecule 3 is water.

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|------------------|---------|---------|
| 3 | A | 68 | Total O 68 68 | 0 | 0 |
| 3 | В | 82 | Total O 82 82 | 0 | 0 |
| 3 | С | 77 | Total O 77 77 | 0 | 0 |
| 3 | Р | 7 | Total O 7 7 | 0 | 0 |
| 3 | Q | 16 | Total O 16 16 | 0 | 0 |




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Bromodomain-containing protein 2





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

| Property | Value | Source |
|--|---|-----------|
| Space group | C 1 2 1 | Depositor |
| Cell constants | 113.81Å 55.14Å 67.49Å | Depositor |
| a, b, c, α , β , γ | 90.00° 94.06° 90.00° | Depositor |
| Resolution (Å) | 20.00 - 2.04 | Depositor |
| % Data completeness | 95.9 (20.00-2.04) | Depositor |
| (in resolution range) | 30.3 (20.00-2.04) | Depositor |
| R_{merge} | 0.07 | Depositor |
| R_{sym} | (Not available) | Depositor |
| Refinement program | REFMAC 5.1.24 | Depositor |
| R, R_{free} | 0.188 , 0.248 | Depositor |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| Total number of atoms | 3161 | wwPDB-VP |
| Average B, all atoms (Å ²) | 31.0 | wwPDB-VP |



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ALY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond | lengths | Bond angles | | |
|-------|-------|------|----------|-------------|--------------------|--|
| IVIOI | Chain | RMSZ | # Z > 5 | RMSZ | # Z >5 | |
| 1 | A | 0.72 | 0/971 | 0.76 | 1/1301 (0.1%) | |
| 1 | В | 0.75 | 0/953 | 0.79 | $2/1277 \ (0.2\%)$ | |
| 1 | С | 0.71 | 0/913 | 0.78 | 1/1224 (0.1%) | |
| 2 | Р | 0.96 | 0/32 | 1.06 | 0/39 | |
| 2 | Q | 1.07 | 0/68 | 1.52 | 1/85 (1.2%) | |
| All | All | 0.74 | 0/2937 | 0.80 | 5/3926 (0.1%) | |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 2 | Q | 0 | 1 |

There are no bond length outliers.

All (5) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | \mathbf{Z} | $Observed(^o)$ | $Ideal(^{o})$ |
|-----|-------|-----|------|-----------|--------------|----------------|---------------|
| 1 | A | 160 | ASP | CB-CG-OD2 | 6.52 | 124.17 | 118.30 |
| 2 | Q | 7 | GLY | N-CA-C | -6.50 | 96.85 | 113.10 |
| 1 | С | 160 | ASP | CB-CG-OD1 | 5.37 | 123.13 | 118.30 |
| 1 | В | 122 | ASP | CB-CG-OD2 | 5.25 | 123.03 | 118.30 |
| 1 | В | 144 | ASP | CB-CG-OD2 | 5.16 | 122.94 | 118.30 |

There are no chirality outliers.

All (1) planarity outliers are listed below:



| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|---------|
| 2 | Q | 2 | GLY | Peptide |

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 953 | 0 | 946 | 29 | 0 |
| 1 | В | 935 | 0 | 932 | 35 | 0 |
| 1 | С | 895 | 0 | 895 | 14 | 0 |
| 2 | Р | 46 | 0 | 42 | 4 | 0 |
| 2 | Q | 82 | 0 | 77 | 25 | 0 |
| 3 | A | 68 | 0 | 0 | 8 | 0 |
| 3 | В | 82 | 0 | 0 | 8 | 0 |
| 3 | С | 77 | 0 | 0 | 6 | 1 |
| 3 | Р | 7 | 0 | 0 | 0 | 0 |
| 3 | Q | 16 | 0 | 0 | 4 | 0 |
| All | All | 3161 | 0 | 2892 | 85 | 1 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | $\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$ | Clash overlap (Å) |
|------------------|------------------|--|----------------------|
| 2:Q:1:SER:OG | 2:Q:2:GLY:HA2 | 1.37 | 1.24 |
| 1:B:118:LYS:HB2 | 2:Q:1:SER:HB3 | 1.42 | 0.98 |
| 1:C:76:THR:HA | 3:C:214:HOH:O | 1.67 | 0.94 |
| 1:B:164:LEU:HD13 | 3:B:216:HOH:O | 1.76 | 0.85 |
| 1:B:118:LYS:H | 2:Q:1:SER:HB3 | 1.43 | 0.83 |
| 1:B:118:LYS:CB | 2:Q:1:SER:HB3 | 2.11 | 0.80 |
| 1:A:159:THR:HG22 | 2:Q:3:ARG:O | 1.82 | 0.79 |
| 1:B:118:LYS:H | 2:Q:1:SER:CB | 1.96 | 0.78 |
| 2:Q:1:SER:CB | 2:Q:2:GLY:HA2 | 2.15 | 0.76 |
| 2:Q:5:LYS:HA | 3:Q:25:HOH:O | 1.85 | 0.76 |
| 1:A:182:GLN:HG3 | 1:B:180:MSE:O | 1.87 | 0.74 |
| 1:B:153:TYR:O | 2:Q:8:LYS:NZ | 2.21 | 0.73 |
| 1:C:160:ASP:CG | 1:C:162:ILE:HG22 | 2.09 | 0.72 |

Continued on next page...



 $Continued\ from\ previous\ page...$

| Continuea from prev | | Interatomic | Clash |
|---------------------|------------------|--------------------------------------|-------------|
| Atom-1 | Atom-2 | ${\rm distance}\ (\mathring{\rm A})$ | overlap (Å) |
| 3:B:253:HOH:O | 2:Q:10:LEU:HD23 | 1.91 | 0.70 |
| 2:Q:10:LEU:HG | 3:Q:24:HOH:O | 1.92 | 0.69 |
| 2:P:14:GLY:HA2 | 2:P:15:ALA:HB2 | 1.77 | 0.66 |
| 1:B:116:ILE:N | 1:B:116:ILE:HD13 | 2.12 | 0.65 |
| 1:B:161:ASP:OD1 | 2:Q:15:ALA:HB3 | 1.98 | 0.64 |
| 1:B:118:LYS:N | 2:Q:1:SER:HB3 | 2.13 | 0.63 |
| 1:A:77:ASN:N | 1:A:77:ASN:HD22 | 1.97 | 0.62 |
| 1:B:108:LEU:HD13 | 2:Q:12:ALY:HD3 | 1.82 | 0.62 |
| 1:A:101:GLN:HG3 | 3:A:250:HOH:O | 2.00 | 0.61 |
| 1:B:116:ILE:HD11 | 2:Q:5:LYS:CB | 2.32 | 0.59 |
| 1:B:101:GLN:HG2 | 3:B:246:HOH:O | 2.01 | 0.59 |
| 1:A:77:ASN:HD22 | 1:A:77:ASN:H | 1.51 | 0.59 |
| 1:C:156:ASN:OD1 | 1:C:162:ILE:HG21 | 2.04 | 0.58 |
| 1:A:97:TRP:CG | 1:A:98:PRO:HD3 | 2.38 | 0.58 |
| 1:B:183:GLU:HG3 | 3:B:256:HOH:O | 2.04 | 0.57 |
| 1:A:185:GLN:O | 1:A:186:GLU:HB2 | 2.05 | 0.56 |
| 1:B:116:ILE:O | 2:Q:1:SER:OG | 2.23 | 0.56 |
| 1:B:117:ILE:HA | 2:Q:1:SER:OG | 2.06 | 0.56 |
| 1:A:73:GLY:HA2 | 1:A:133:ASN:O | 2.07 | 0.55 |
| 1:A:182:GLN:H | 1:B:182:GLN:NE2 | 2.04 | 0.55 |
| 1:A:76:THR:HA | 1:A:186:GLU:HB2 | 1.89 | 0.55 |
| 1:B:116:ILE:N | 1:B:116:ILE:CD1 | 2.69 | 0.55 |
| 1:A:182:GLN:H | 1:B:182:GLN:HE21 | 1.55 | 0.55 |
| 1:B:97:TRP:CG | 1:B:98:PRO:HD3 | 2.43 | 0.54 |
| 1:B:76:THR:HG23 | 3:B:231:HOH:O | 2.08 | 0.54 |
| 2:Q:1:SER:OG | 2:Q:2:GLY:CA | 2.32 | 0.54 |
| 1:A:183:GLU:HA | 3:A:245:HOH:O | 2.07 | 0.53 |
| 1:A:171:LYS:HG2 | 1:B:143:GLN:OE1 | 2.09 | 0.53 |
| 1:C:97:TRP:CG | 1:C:98:PRO:HD3 | 2.43 | 0.53 |
| 1:C:95:PHE:CG | 1:C:165:MSE:HG2 | 2.44 | 0.53 |
| 1:B:154:ILE:HG21 | 2:Q:2:GLY:HA3 | 1.92 | 0.52 |
| 3:B:269:HOH:O | 1:C:162:ILE:HB | 2.10 | 0.51 |
| 1:A:75:VAL:O | 1:A:186:GLU:HB3 | 2.11 | 0.51 |
| 1:A:86:VAL:HG22 | 1:A:176:LYS:HB3 | 1.93 | 0.51 |
| 1:B:158:PRO:CD | 2:P:8:LYS:HA | 2.41 | 0.50 |
| 2:Q:1:SER:HG | 2:Q:2:GLY:HA2 | 1.68 | 0.49 |
| 2:Q:1:SER:CB | 2:Q:2:GLY:CA | 2.89 | 0.47 |
| 1:C:118:LYS:HD3 | 3:C:225:HOH:O | 2.13 | 0.47 |
| 1:B:110:LEU:HA | 3:B:253:HOH:O | 2.15 | 0.47 |
| 1:B:116:ILE:HD12 | 2:Q:5:LYS:H | 1.79 | 0.47 |
| 1:A:171:LYS:HG3 | 3:A:218:HOH:O | 2.14 | 0.47 |

Continued on next page...



 $Continued\ from\ previous\ page...$

| A | | Interatomic | Clash | |
|------------------|------------------|---------------------------------------|-------------|--|
| Atom-1 | Atom-2 | ${\rm distance} \ (\mathring{\rm A})$ | overlap (Å) | |
| 1:C:100:ARG:HD2 | 3:C:240:HOH:O | 2.15 | 0.46 | |
| 1:C:88:LYS:HE2 | 3:C:236:HOH:O | 2.14 | 0.46 | |
| 1:B:78:GLN:HB3 | 1:B:138:ALA:HB2 | 1.98 | 0.46 | |
| 1:A:75:VAL:HB | 3:A:226:HOH:O | 2.16 | 0.45 | |
| 1:A:101:GLN:HG2 | 3:A:214:HOH:O | 2.17 | 0.45 | |
| 1:A:167:GLN:O | 1:A:171:LYS:HG3 | 2.15 | 0.45 | |
| 1:B:90:LEU:HD12 | 1:B:145:PHE:CE1 | 2.52 | 0.45 | |
| 1:B:143:GLN:HG2 | 3:B:262:HOH:O | 2.15 | 0.45 | |
| 1:B:97:TRP:CD2 | 1:B:98:PRO:HD3 | 2.51 | 0.45 | |
| 1:B:115:LYS:HB2 | 1:B:116:ILE:HD13 | 1.98 | 0.45 | |
| 1:C:95:PHE:CD1 | 1:C:165:MSE:HG2 | 2.52 | 0.45 | |
| 1:B:118:LYS:HB2 | 2:Q:1:SER:CB | 2.30 | 0.45 | |
| 1:B:142:MSE:HE3 | 1:B:177:VAL:HG11 | 1.98 | 0.44 | |
| 1:A:184:GLU:HB2 | 3:C:215:HOH:O | 2.17 | 0.44 | |
| 2:Q:10:LEU:HB3 | 3:Q:24:HOH:O | 2.18 | 0.44 | |
| 1:A:74:ARG:HD2 | 1:A:136:TRP:CD2 | 2.53 | 0.43 | |
| 1:A:74:ARG:HD2 | 1:A:136:TRP:CE2 | 2.53 | 0.43 | |
| 1:A:128:ARG:HD3 | 3:A:256:HOH:O | 2.18 | 0.43 | |
| 1:A:154:ILE:O | 2:Q:8:LYS:HE3 | 2.19 | 0.42 | |
| 1:C:181:PRO:O | 1:C:182:GLN:HB2 | 2.21 | 0.41 | |
| 1:C:164:LEU:HA | 1:C:167:GLN:HE21 | 1.86 | 0.41 | |
| 1:C:164:LEU:HD13 | 3:C:226:HOH:O | 2.20 | 0.41 | |
| 1:A:97:TRP:CD2 | 1:A:98:PRO:HD3 | 2.56 | 0.41 | |
| 1:B:118:LYS:HE2 | 3:Q:23:HOH:O | 2.21 | 0.41 | |
| 1:A:164:LEU:HA | 1:A:167:GLN:HE21 | 1.85 | 0.41 | |
| 1:A:128:ARG:NE | 3:A:235:HOH:O | 2.55 | 0.41 | |
| 1:A:77:ASN:N | 1:A:77:ASN:ND2 | 2.67 | 0.40 | |
| 3:A:217:HOH:O | 2:P:10:LEU:HD12 | 2.21 | 0.40 | |
| 1:A:112:ASP:OD2 | 2:P:10:LEU:HB2 | 2.21 | 0.40 | |
| 1:B:118:LYS:HA | 1:B:118:LYS:HD2 | 1.69 | 0.40 | |
| 1:C:80:GLN:O | 1:C:84:LYS:HB2 | 2.20 | 0.40 | |

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

| Atom-1 | Atom-1 Atom-2 | | $egin{aligned} 	ext{Clash} \ 	ext{overlap } (ext{Å}) \end{aligned}$ | |
|---------------|----------------------|------|--|--|
| 3:C:208:HOH:O | 3:C:208:HOH:O[2_655] | 1.90 | 0.30 | |



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles |
|-----|-------|--------------------|-----------|---------|----------|-------------|
| 1 | A | $112/122 \ (92\%)$ | 111 (99%) | 0 | 1 (1%) | 17 8 |
| 1 | В | 110/122 (90%) | 109 (99%) | 0 | 1 (1%) | 17 8 |
| 1 | С | 105/122~(86%) | 104 (99%) | 1 (1%) | 0 | 100 100 |
| 2 | Р | 5/15 (33%) | 3 (60%) | 2 (40%) | 0 | 100 100 |
| 2 | Q | 12/15 (80%) | 7 (58%) | 3 (25%) | 2 (17%) | 0 0 |
| All | All | 344/396 (87%) | 334 (97%) | 6 (2%) | 4 (1%) | 13 5 |

All (4) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 184 | GLU |
| 2 | Q | 9 | GLY |
| 2 | Q | 10 | LEU |
| 1 | В | 75 | VAL |

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Rotameric Outliers | | Percentiles | | |
|-----|-------|---------------|-----------|--------------------|-----|-------------|--|--|
| 1 | A | 104/105 (99%) | 96 (92%) | 8 (8%) | 13 | 5 | | |
| 1 | В | 102/105 (97%) | 95 (93%) | 7 (7%) | 15 | 8 | | |
| 1 | С | 98/105 (93%) | 91 (93%) | 7 (7%) | 14 | 7 | | |
| 2 | Р | 1/5 (20%) | 1 (100%) | 0 | 100 | 100 | | |

Continued on next page...



Continued from previous page...

| Mol | Chain | Analysed | Analysed Rotameric Outliers | | Percentiles | | |
|-----|-------|---------------|-----------------------------|---------|-------------|--|--|
| 2 | Q | 3/5 (60%) | 3 (100%) | 0 | 100 100 | | |
| All | All | 308/325~(95%) | 286 (93%) | 22 (7%) | 14 7 | | |

All (22) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|---------|-----|------|
| 1 | A | 74 | ARG |
| 1 | A | 77 | ASN |
| 1 | A | 83 | HIS |
| 1 | A | 84 | LYS |
| 1 | A | 101 | GLN |
| 1 | A | 110 | LEU |
| 1 | A | 127 | LYS |
| 1 | A | 162 | ILE |
| 1 | В | 84 | LYS |
| 1 | В | 101 | GLN |
| 1 | В | 110 | LEU |
| 1 | В | 116 | ILE |
| 1 | В | 118 | LYS |
| 1 | В | 164 | LEU |
| 1 | В | 183 | GLU |
| 1 | С | 83 | HIS |
| 1 | C C C C | 84 | LYS |
| 1 | С | 92 | LYS |
| 1 | С | 101 | GLN |
| 1 | С | 108 | LEU |
| 1 | С | 110 | LEU |
| 1 | С | 157 | LYS |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 77 | ASN |
| 1 | A | 78 | GLN |
| 1 | A | 167 | GLN |
| 1 | В | 78 | GLN |
| 1 | В | 182 | GLN |
| 1 | С | 83 | HIS |
| 1 | С | 101 | GLN |
| 1 | С | 167 | GLN |



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol Type Chain | | Dag | Link | Bo | ond lengths | | Bond angles | | | |
|----------------|------|-------|---------|-------|-------------|------|-------------|---------|------|----------|
| MIOI | Type | Chain | ain Res | Lilik | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 2 | ALY | Р | 12 | 2 | 10,11,12 | 3.58 | 3 (30%) | 7,12,14 | 2.98 | 2 (28%) |
| 2 | ALY | Q | 12 | 2 | 10,11,12 | 3.54 | 3 (30%) | 7,12,14 | 3.94 | 5 (71%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|-------|
| 2 | ALY | Р | 12 | 2 | - | 7/9/10/12 | - |
| 2 | ALY | Q | 12 | 2 | - | 5/9/10/12 | - |

All (6) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | \mathbf{Z} | $\operatorname{Observed}(\mathring{A})$ | $Ideal(\AA)$ |
|-----|-------|-----|------|-------|--------------|---|--------------|
| 2 | Р | 12 | ALY | ОН-СН | 8.70 | 1.42 | 1.23 |
| 2 | Q | 12 | ALY | ОН-СН | 8.59 | 1.42 | 1.23 |
| 2 | Q | 12 | ALY | O-C | 5.44 | 1.41 | 1.19 |
| 2 | Р | 12 | ALY | O-C | 5.23 | 1.40 | 1.19 |
| 2 | Р | 12 | ALY | CH-NZ | 4.40 | 1.46 | 1.34 |
| 2 | Q | 12 | ALY | CH-NZ | 4.05 | 1.45 | 1.34 |

All (7) bond angle outliers are listed below:



| Mol | Chain | Res | Type | Atoms | Z | $Observed(^o)$ | $Ideal(^{o})$ |
|-----|-------|-----|------|-----------|-------|----------------|---------------|
| 2 | Р | 12 | ALY | ОН-СН-СН3 | -6.99 | 109.07 | 122.06 |
| 2 | Q | 12 | ALY | ОН-СН-СН3 | -6.48 | 110.02 | 122.06 |
| 2 | Q | 12 | ALY | CE-NZ-CH | -5.87 | 113.53 | 122.56 |
| 2 | Q | 12 | ALY | OH-CH-NZ | -3.79 | 111.06 | 121.74 |
| 2 | Q | 12 | ALY | CD-CE-NZ | 3.60 | 122.52 | 112.21 |
| 2 | Р | 12 | ALY | OH-CH-NZ | -2.37 | 115.06 | 121.74 |
| 2 | Q | 12 | ALY | CH3-CH-NZ | 2.02 | 119.66 | 116.09 |

There are no chirality outliers.

All (12) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|--------------|
| 2 | P | 12 | ALY | CH3-CH-NZ-CE |
| 2 | P | 12 | ALY | N-CA-CB-CG |
| 2 | P | 12 | ALY | C-CA-CB-CG |
| 2 | Q | 12 | ALY | N-CA-CB-CG |
| 2 | Q | 12 | ALY | C-CA-CB-CG |
| 2 | P | 12 | ALY | CG-CD-CE-NZ |
| 2 | Q | 12 | ALY | CH3-CH-NZ-CE |
| 2 | Q | 12 | ALY | CG-CD-CE-NZ |
| 2 | Р | 12 | ALY | CA-CB-CG-CD |
| 2 | Q | 12 | ALY | CA-CB-CG-CD |
| 2 | P | 12 | ALY | OH-CH-NZ-CE |
| 2 | P | 12 | ALY | CE-CD-CG-CB |

There are no ring outliers.

1 monomer is involved in 1 short contact:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|----------------------|------|---------|--------------|
| 2 | Q | 12 | ALY | 1 | 0 |

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

