



Full wwPDB X-ray Structure Validation Report ⓘ

May 28, 2020 – 09:34 pm BST

PDB ID : 2EDC
Title : CRYSTALLOGRAPHIC AND FLUORESCENCE STUDIES OF THE INTERACTION OF HALOALKANE DEHALOGENASE WITH HALIDE IONS: STUDIES WITH HALIDE COMPOUNDS REVEAL A HALIDE BINDING SITE IN THE ACTIVE SITE
Authors : Verschueren, K.H.G.; Dijkstra, B.W.
Deposited on : 1993-08-31
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	310	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2605 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HALOALKANE DEHALOGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	310	2479	1596	406	462	15	0	0	0

- Molecule 2 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total I 1 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	125	Total O 125 125	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: HALOALKANE DEHALOGENASE



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	94.90Å 73.10Å 41.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT	Depositor
R, R_{free}	0.177 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2605	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/2552	1.68	55/3470 (1.6%)

There are no bond length outliers.

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	300	ARG	NE-CZ-NH1	12.87	126.73	120.30
1	A	170	ASP	CB-CG-OD1	12.48	129.53	118.30
1	A	143	ARG	NE-CZ-NH1	11.74	126.17	120.30
1	A	27	ASP	CB-CG-OD1	11.64	128.78	118.30
1	A	124	ASP	CB-CG-OD2	-9.46	109.79	118.30
1	A	170	ASP	CB-CG-OD2	-9.20	110.02	118.30
1	A	116	ARG	NE-CZ-NH2	-8.70	115.95	120.30
1	A	116	ARG	NE-CZ-NH1	8.48	124.54	120.30
1	A	188	ASP	CB-CG-OD1	8.46	125.91	118.30
1	A	230	ASP	CB-CG-OD1	8.32	125.79	118.30
1	A	300	ARG	NE-CZ-NH2	-8.16	116.22	120.30
1	A	35	ARG	NE-CZ-NH2	8.14	124.37	120.30
1	A	266	ASP	CB-CG-OD2	-8.13	110.99	118.30
1	A	64	ARG	NE-CZ-NH1	8.08	124.34	120.30
1	A	124	ASP	CB-CG-OD1	7.75	125.28	118.30
1	A	45	ASP	CB-CG-OD2	-7.51	111.54	118.30
1	A	45	ASP	CB-CG-OD1	7.28	124.85	118.30
1	A	193	ARG	NE-CZ-NH1	7.11	123.85	120.30
1	A	112	ARG	NE-CZ-NH1	6.96	123.78	120.30
1	A	6	ARG	NE-CZ-NH2	6.95	123.78	120.30
1	A	76	ARG	NE-CZ-NH2	6.86	123.73	120.30
1	A	143	ARG	NE-CZ-NH2	-6.85	116.88	120.30
1	A	27	ASP	CB-CG-OD2	-6.82	112.16	118.30
1	A	114	ASP	CB-CG-OD2	-6.74	112.24	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	197	THR	CA-CB-CG2	-6.63	103.12	112.40
1	A	223	PRO	N-CA-CB	6.62	111.24	103.30
1	A	186	ARG	NE-CZ-NH2	6.60	123.60	120.30
1	A	220	ARG	NE-CZ-NH2	6.50	123.55	120.30
1	A	35	ARG	NE-CZ-NH1	-6.31	117.15	120.30
1	A	271	MET	CG-SD-CE	6.26	110.22	100.20
1	A	103	ARG	NE-CZ-NH1	-6.12	117.24	120.30
1	A	103	ARG	CD-NE-CZ	-6.11	115.05	123.60
1	A	147	MET	CG-SD-CE	6.09	109.94	100.20
1	A	300	ARG	CD-NE-CZ	6.07	132.10	123.60
1	A	229	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	A	114	ASP	CB-CG-OD1	5.99	123.69	118.30
1	A	93	ASP	CB-CG-OD2	-5.97	112.92	118.30
1	A	18	TYR	CB-CG-CD2	-5.96	117.43	121.00
1	A	225	MET	CG-SD-CE	5.93	109.68	100.20
1	A	235	ASP	CB-CG-OD1	5.83	123.54	118.30
1	A	184	ASP	CB-CG-OD2	-5.79	113.09	118.30
1	A	258	MET	CG-SD-CE	5.78	109.44	100.20
1	A	53	LEU	CB-CG-CD2	-5.75	101.23	111.00
1	A	26	ASP	CB-CG-OD2	-5.66	113.20	118.30
1	A	152	MET	CG-SD-CE	5.65	109.25	100.20
1	A	22	PRO	O-C-N	5.55	131.58	122.70
1	A	96	ASP	CB-CG-OD1	5.55	123.30	118.30
1	A	235	ASP	CB-CG-OD2	-5.51	113.34	118.30
1	A	48	ASP	CB-CG-OD2	-5.46	113.38	118.30
1	A	201	ALA	CB-CA-C	5.41	118.21	110.10
1	A	210	PHE	N-CA-CB	-5.33	101.01	110.60
1	A	26	ASP	CB-CG-OD1	5.20	122.98	118.30
1	A	260	ASP	CB-CG-OD1	5.18	122.96	118.30
1	A	229	ARG	NE-CZ-NH2	-5.08	117.76	120.30
1	A	237	SER	N-CA-CB	-5.06	102.91	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2479	0	2379	34	0
2	A	1	0	0	0	0
3	A	125	0	0	3	0
All	All	2605	0	2379	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:GLN:HE21	1:A:292:GLN:H	1.19	0.85
1:A:220:ARG:HD3	3:A:472:HOH:O	1.89	0.72
1:A:309:THR:HG22	1:A:309:THR:O	1.91	0.70
1:A:307:ALA:C	1:A:309:THR:H	1.95	0.68
1:A:142:LYS:HG2	1:A:143:ARG:HG3	1.77	0.67
1:A:309:THR:O	1:A:310:GLU:HG2	2.01	0.60
1:A:96:ASP:OD2	3:A:406:HOH:O	2.17	0.58
1:A:149:ALA:HA	1:A:263:LEU:HD13	1.87	0.56
1:A:112:ARG:HH11	1:A:112:ARG:HG3	1.71	0.54
1:A:153:THR:HG21	1:A:158:GLN:HB2	1.91	0.53
1:A:151:LEU:HD13	1:A:241:ILE:HG13	1.92	0.51
1:A:103:ARG:HD3	3:A:420:HOH:O	2.09	0.51
1:A:307:ALA:C	1:A:309:THR:N	2.64	0.51
1:A:296:GLU:O	1:A:300:ARG:HG3	2.13	0.48
1:A:31:TYR:N	1:A:32:PRO:CD	2.77	0.48
1:A:304:LYS:HE3	1:A:304:LYS:HB2	1.60	0.46
1:A:91:PRO:HD2	1:A:97:TYR:OH	2.15	0.46
1:A:284:ILE:HD12	1:A:291:VAL:HG13	1.99	0.45
1:A:112:ARG:NH1	1:A:112:ARG:HG3	2.32	0.44
1:A:134:PRO:HB2	1:A:243:PHE:CE2	2.53	0.44
1:A:157:THR:C	1:A:158:GLN:HG2	2.38	0.43
1:A:284:ILE:HD11	1:A:298:VAL:HG22	2.00	0.43
1:A:47:GLU:O	1:A:117:ASN:HB2	2.19	0.42
1:A:154:ASP:OD1	1:A:157:THR:CG2	2.67	0.42
1:A:187:LEU:HD23	1:A:187:LEU:HA	1.84	0.42
1:A:20:PHE:HB2	1:A:39:LEU:HD22	2.02	0.41
1:A:103:ARG:HH11	1:A:103:ARG:HD3	1.41	0.41
1:A:49:VAL:HG11	1:A:115:LEU:HD22	2.03	0.41
1:A:307:ALA:O	1:A:309:THR:N	2.50	0.41
1:A:208:ALA:N	1:A:209:PRO:CD	2.84	0.41
1:A:157:THR:O	1:A:158:GLN:HG2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:THR:CG2	1:A:158:GLN:HB2	2.51	0.40
1:A:208:ALA:HB3	1:A:209:PRO:HD3	2.03	0.40
1:A:56:GLU:HA	1:A:57:PRO:HA	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	308/310 (99%)	289 (94%)	17 (6%)	2 (1%)	25 31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	ASN
1	A	137	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	262/262 (100%)	246 (94%)	16 (6%)	18 25

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	PRO
1	A	14	ASN
1	A	15	LEU
1	A	27	ASP
1	A	73	SER
1	A	113	LEU
1	A	116	ARG
1	A	123	GLN
1	A	157	THR
1	A	222	PHE
1	A	246	ASN
1	A	266	ASP
1	A	271	MET
1	A	272	LYS
1	A	292	GLN
1	A	304	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	ASN
1	A	104	ASN
1	A	123	GLN
1	A	251	GLN
1	A	292	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.