



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2024 – 11:20 AM EST

PDB ID : 2F7K
Title : Crystal Structure of Human Pyridoxal Kinase
Authors : Jiang, T.; Cao, P.; Gong, Y.; Tang, L.
Deposited on : 2005-12-01
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	
1	B	327	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 5327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridoxal kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	323	2560	1608	458	478	16	0	0	0
1	B	323	2560	1608	458	478	16	0	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	SER	-	cloning artifact	UNP O00764
A	-13	TYR	-	cloning artifact	UNP O00764
A	-12	TYR	-	cloning artifact	UNP O00764
A	-11	HIS	-	expression tag	UNP O00764
A	-10	HIS	-	expression tag	UNP O00764
A	-9	HIS	-	expression tag	UNP O00764
A	-8	HIS	-	expression tag	UNP O00764
A	-7	HIS	-	expression tag	UNP O00764
A	-6	HIS	-	expression tag	UNP O00764
A	-5	HIS	-	expression tag	UNP O00764
A	-4	GLU	-	cloning artifact	UNP O00764
A	-3	GLY	-	cloning artifact	UNP O00764
A	-2	VAL	-	cloning artifact	UNP O00764
A	-1	ARG	-	cloning artifact	UNP O00764
A	0	THR	-	cloning artifact	UNP O00764
B	-14	SER	-	cloning artifact	UNP O00764
B	-13	TYR	-	cloning artifact	UNP O00764
B	-12	TYR	-	cloning artifact	UNP O00764
B	-11	HIS	-	expression tag	UNP O00764
B	-10	HIS	-	expression tag	UNP O00764
B	-9	HIS	-	expression tag	UNP O00764
B	-8	HIS	-	expression tag	UNP O00764
B	-7	HIS	-	expression tag	UNP O00764
B	-6	HIS	-	expression tag	UNP O00764
B	-5	HIS	-	expression tag	UNP O00764

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLU	-	cloning artifact	UNP O00764
B	-3	GLY	-	cloning artifact	UNP O00764
B	-2	VAL	-	cloning artifact	UNP O00764
B	-1	ARG	-	cloning artifact	UNP O00764
B	0	THR	-	cloning artifact	UNP O00764

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	96	Total O 96 96	0	0
2	B	111	Total O 111 111	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Pyridoxal kinase

Chain A: 



• Molecule 1: Pyridoxal kinase

Chain B: 



P249	M250	N251	L252	K253	C256	E257	R258	T259	V260	S261	T262	L263	H264	V266	L267	Q268	R269	T270	I271	Q272	C273	A274	A278	G279	E280	G281	V282	R283	P284	S285	P286	M287	Q288	L289	R292	M293	V294	Q295	S296	K297	R298	D299	I300	E301	D302	P303	E304	I305	V306	V307	Q308	A309	I310	V311	L312
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics i

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	52.50Å 52.50Å 301.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.80 7.99 – 2.80	Depositor EDS
% Data completeness (in resolution range)	81.0 (8.00-2.80) 81.0 (7.99-2.80)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.94 (at 2.78Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.228 , 0.271 0.213 , 0.255	Depositor DCC
R_{free} test set	709 reflections (4.59%)	wwPDB-VP
Wilson B-factor (Å ²)	68.3	Xtrriage
Anisotropy	0.089	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 71.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.190 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5327	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/2612	0.87	10/3540 (0.3%)
1	B	0.45	0/2612	0.83	3/3540 (0.1%)
All	All	0.45	0/5224	0.85	13/7080 (0.2%)

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	PRO	CA-N-CD	-7.94	100.39	111.50
1	A	149	PRO	CA-N-CD	-7.75	100.66	111.50
1	B	120	TRP	CA-CB-CG	6.54	126.12	113.70
1	A	-3	GLY	N-CA-C	-6.51	96.83	113.10
1	A	192	SER	N-CA-C	-5.85	95.22	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2560	0	2551	354	0
1	B	2560	0	2551	334	0
2	A	96	0	0	0	0
2	B	111	0	0	3	0
All	All	5327	0	5102	660	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 65.

The worst 5 of 660 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:LEU:HB2	1:A:223:ILE:HB	1.38	1.05
1:A:72:ASN:HD21	1:A:74:MET:HG3	1.22	1.04
1:A:221:MET:HG2	1:A:309:ALA:HA	1.39	1.02
1:A:147:ILE:CD1	1:A:147:ILE:H	1.74	1.00
1:A:147:ILE:H	1:A:147:ILE:HD13	1.24	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	321/327 (98%)	229 (71%)	65 (20%)	27 (8%)	1 2
1	B	321/327 (98%)	240 (75%)	65 (20%)	16 (5%)	2 6
All	All	642/654 (98%)	469 (73%)	130 (20%)	43 (7%)	1 3

5 of 43 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	-9	HIS
1	A	59	SER
1	A	109	VAL
1	A	195	GLY
1	A	210	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/289 (99%)	226 (79%)	59 (21%)	1	3
1	B	285/289 (99%)	223 (78%)	62 (22%)	1	3
All	All	570/578 (99%)	449 (79%)	121 (21%)	1	3

5 of 121 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	307	VAL
1	B	260	VAL
1	B	45	ASN
1	B	250	ASN
1	B	304	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	-10	HIS
1	B	21	ASN
1	B	209	ASN
1	B	-5	HIS
1	B	29	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/327 (98%)	-0.28	0 100 100	36, 44, 50, 56	0
1	B	323/327 (98%)	-0.28	0 100 100	35, 44, 50, 55	0
All	All	646/654 (98%)	-0.28	0 100 100	35, 44, 50, 56	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.