



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 07:01 pm BST

PDB ID : 2GD5
Title : Structural basis for budding by the ESCRTIII factor CHMP3
Authors : Muziol, T.M.; Pineda-Molina, E.; Ravelli, R.B.; Zamborlini, A.; Usami, Y.;
Gottlinger, H.; Weissenhorn, W.
Deposited on : 2006-03-15
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	179	 2% 35% 36% 7% • 21%
1	B	179	 2% 45% 29% 12% • 13%
1	C	179	 2% 42% 32% • • 21%
1	D	179	 45% 39% 7% 9%

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 4746 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Charged multivesicular body protein 3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	142	1134	710	207	205	1	11	0	0	0
1	B	156	1219	763	223	220	1	12	0	0	0
1	C	142	1123	703	204	204	1	11	0	0	0
1	D	162	1270	793	229	234	1	13	0	0	0

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	GLY	-	CLONING ARTIFACT	UNP Q9Y3E7
A	6	ALA	-	CLONING ARTIFACT	UNP Q9Y3E7
A	7	MSE	-	CLONING ARTIFACT	UNP Q9Y3E7
A	8	ALA	-	CLONING ARTIFACT	UNP Q9Y3E7
A	27	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	67	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	84	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	89	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	91	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	111	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	114	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	127	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	134	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	135	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	157	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
A	163	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	5	GLY	-	CLONING ARTIFACT	UNP Q9Y3E7
B	6	ALA	-	CLONING ARTIFACT	UNP Q9Y3E7
B	7	MSE	-	CLONING ARTIFACT	UNP Q9Y3E7
B	8	ALA	-	CLONING ARTIFACT	UNP Q9Y3E7
B	27	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	67	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	84	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	89	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	91	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	111	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	114	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	127	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	134	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	135	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	157	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
B	163	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	5	GLY	-	CLONING ARTIFACT	UNP Q9Y3E7
C	6	ALA	-	CLONING ARTIFACT	UNP Q9Y3E7
C	7	MSE	-	CLONING ARTIFACT	UNP Q9Y3E7
C	8	ALA	-	CLONING ARTIFACT	UNP Q9Y3E7
C	27	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	67	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	84	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	89	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	91	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	111	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	114	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	127	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	134	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	135	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	157	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
C	163	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	5	GLY	-	CLONING ARTIFACT	UNP Q9Y3E7
D	6	ALA	-	CLONING ARTIFACT	UNP Q9Y3E7
D	7	MSE	-	CLONING ARTIFACT	UNP Q9Y3E7
D	8	ALA	-	CLONING ARTIFACT	UNP Q9Y3E7
D	27	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	67	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	84	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	89	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	91	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	111	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	114	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	127	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	134	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	135	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7
D	157	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	163	MSE	MET	MODIFIED RESIDUE	UNP Q9Y3E7


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Charged multivesicular body protein 3



- Molecule 1: Charged multivesicular body protein 3





- Molecule 1: Charged multivesicular body protein 3





- Molecule 1: Charged multivesicular body protein 3

Chain D: 45% 39% 7% 9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	33.58Å 72.57Å 89.23Å 68.68° 83.61° 76.70°	Depositor
Resolution (Å)	20.00 – 2.80 33.09 – 2.80	Depositor EDS
% Data completeness (in resolution range)	95.7 (20.00-2.80) 95.6 (33.09-2.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.63 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.259 , 0.301 0.252 , 0.292	Depositor DCC
R_{free} test set	914 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	62.6	Xtrriage
Anisotropy	0.714	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4746	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/1128	0.74	0/1476
1	B	0.72	0/1213	0.76	0/1589
1	C	0.71	0/1116	0.70	0/1459
1	D	0.73	0/1265	0.74	0/1659
All	All	0.72	0/4722	0.74	0/6183

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1134	0	1232	78	0
1	B	1219	0	1325	78	0
1	C	1123	0	1218	68	0
1	D	1270	0	1365	68	0
All	All	4746	0	5140	273	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 273 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:169:LEU:O	1:D:173:THR:HG22	1.37	1.20
1:C:33:GLN:HE22	1:C:122:GLU:HG2	1.16	1.09
1:C:33:GLN:NE2	1:C:122:GLU:HG2	1.74	1.00
1:D:160:GLU:HB3	1:D:163:MSE:HE2	1.40	1.00
1:C:71:ARG:NH2	1:D:101:ALA:HB2	1.77	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/179 (77%)	118 (86%)	12 (9%)	8 (6%)	1	4
1	B	152/179 (85%)	121 (80%)	20 (13%)	11 (7%)	1	2
1	C	136/179 (76%)	108 (79%)	16 (12%)	12 (9%)	1	1
1	D	158/179 (88%)	143 (90%)	11 (7%)	4 (2%)	5	19
All	All	584/716 (82%)	490 (84%)	59 (10%)	35 (6%)	1	4

5 of 35 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	121	PRO
1	A	122	GLU
1	B	8	ALA
1	B	122	GLU
1	B	124	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/142 (89%)	112 (89%)	14 (11%)	6	19
1	B	132/142 (93%)	118 (89%)	14 (11%)	6	20
1	C	124/142 (87%)	112 (90%)	12 (10%)	8	24
1	D	138/142 (97%)	124 (90%)	14 (10%)	7	22
All	All	520/568 (92%)	466 (90%)	54 (10%)	7	21

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	162	GLU
1	C	105	GLN
1	D	157	MSE
1	B	164	GLU
1	B	179	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	38	GLN
1	B	94	GLN
1	D	83	HIS
1	A	124	GLN
1	D	17	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	131/179 (73%)	0.01	3 (2%) 60 51	23, 56, 91, 93	0
1	B	144/179 (80%)	0.08	4 (2%) 53 43	38, 58, 82, 90	0
1	C	131/179 (73%)	0.04	3 (2%) 60 51	34, 61, 92, 98	0
1	D	149/179 (83%)	-0.07	0 100 100	29, 55, 71, 82	0
All	All	555/716 (77%)	0.01	10 (1%) 68 61	23, 58, 88, 98	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	137	ALA	4.1
1	B	9	GLU	3.1
1	A	110	VAL	2.5
1	A	121	PRO	2.3
1	B	5	GLY	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.