

wwPDB NMR Structure Validation Summary Report (i)

Jun 3, 2023 – 05:06 AM EDT

PDB ID : 2H3K BMRB ID : 6759

Title: Solution Structure of the first NEAT domain of IsdH

Authors: Pilpa, R.M.; Fadeev, E.A.; Villareal, V.A.; Wong, M.A.; Phillips, M.; Clubb,

R.T.

Deposited on : 2006-05-22

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-ShiftChecker} &:& v1.2\\ BMRB \ Restraints \ Analysis &:& v1.2 \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 79%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$rac{ m NMR~archive}{ m (\#Entries)}$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain			
1	A	144	41%	45%	6%	8%



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: fewest violations, lowest energy, minimized average structure.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues				
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model				
1	A:3-A:37, A:45-A:141 (132)	0.58	5	

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	2, 4, 5, 6, 8, 9, 10, 14, 16, 18, 19, 20
2	1, 7, 11, 13, 15, 17
3	3, 12



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2319 atoms, of which 1135 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Haptoglobin-binding surface anchored protein.

Mol	Chain	Residues	Atoms			Trace		
1	Λ	1.4.4	Total	С	Н	N	О	0
1	A	144	2319	754	1135	189	241	U




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: Haptoglobin-binding surface anchored protein



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.

• Molecule 1: Haptoglobin-binding surface anchored protein





5 Refinement protocol and experimental data overview (i)



The models were refined using the following method: distance geometry.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: structures with acceptable covalent geometry, structures with favorable non-bond energy, structures with the least restraint violations, structures with the lowest energy.

The authors did not provide any information on software used for structure solution, optimization or refinement.

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1508
Number of shifts mapped to atoms	1508
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	79%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0 ± 0.0	3.0 ± 0.0
All	All	0	60

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	24	ARG	Sidechain	20
1	A	86	ARG	Sidechain	20
1	A	101	ARG	Sidechain	20

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1077	1043	1043	43±6
All	All	21540	20860	20860	857

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

5 of 254 unique clashes are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Clash(Å)	$Distance(\mathring{A})$	Models	
Atom-1	Atom-1 Atom-2 Cla		Distance(A)	Worst	Total
1:A:52:VAL:HG11	1:A:54:TYR:CZ	0.83	2.09	20	16
1:A:61:VAL:HG11	1:A:134:PHE:CZ	0.79	2.13	12	14
1:A:74:PHE:CE2	1:A:85:VAL:HG21	0.79	2.13	16	9
1:A:110:GLU:C	1:A:111:LEU:HD23	0.76	2.01	17	20
1:A:104:VAL:HG21	1:A:138:ILE:CG2	0.76	2.09	9	16

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	132/144 (92%)	105±3 (80±2%)	22±3 (17±2%)	5±2 (4±1%)	6	34
All	All	$2640/2880 \ (92\%)$	2099 (80%)	443 (17%)	98 (4%)	6	34

5 of 22 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	33	ASP	16
1	A	96	ASP	14
1	A	122	GLY	12
1	A	80	ASN	10
1	A	45	SER	7

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles			
1	A	122/133 (92%)	90±3 (74±3%)	32±3 (26±3%)	2 23			
All	All	2440/2660 (92%)	1807 (74%)	633 (26%)	2 23			



5 of 85 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	77	TYR	20
1	A	86	ARG	20
1	A	111	LEU	20
1	A	112	LYS	20
1	A	72	LYS	19

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 79% for the well-defined parts and 76% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1508
Number of shifts mapped to atoms	1508
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	139	0.49 ± 0.15	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	136	0.32 ± 0.14	None needed ($< 0.5 \text{ ppm}$)
¹³ C′	114	0.27 ± 0.15	None needed ($< 0.5 \text{ ppm}$)
^{15}N	122	0.03 ± 0.76	None needed ($< 0.5 \text{ ppm}$)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 79%, i.e. 1427 atoms were assigned a chemical shift out of a possible 1809. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	601/647 (93%)	247/259 (95%)	238/264 (90%)	116/124 (94%)
Sidechain	774/1024 (76%)	504/655 (77%)	270/332 (81%)	0/37 (0%)

Continued on next page...



Continued from previous page...

	Total	$^{1}{ m H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Aromatic	52/138 (38%)	$26/66 \ (39\%)$	$25/69 \ (36\%)$	1/3 (33%)
Overall	1427/1809 (79%)	777/980 (79%)	533/665 (80%)	117/164 (71%)

7.1.4 Statistically unusual chemical shifts (i)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	102	PHE	CE1	117.34	124.17 - 137.29	-10.2
1	A	71	TRP	CE3	131.04	111.58 - 129.41	5.9
1	A	71	TRP	CZ3	129.85	113.48 - 129.28	5.4
1	A	122	GLY	N	128.52	91.59 - 127.52	5.3

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1807
Intra-residue ($ i-j =0$)	369
Sequential ($ i-j =1$)	455
Medium range ($ i-j >1$ and $ i-j <5$)	232
Long range (i-j ≥5)	685
Inter-chain	0
Hydrogen bond restraints	66
Disulfide bond restraints	0
Total dihedral-angle restraints	140
Number of unmapped restraints	0
Number of restraints per residue	13.5
Number of long range restraints per residue ¹	5.2

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	2.5	0.17
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	$\operatorname{Max}(^{\circ})$
1.0-10.0 (Small)	2.1	2.3
10.0-20.0 (Medium)	None	None
>20.0 (Large)	4.0	105.1



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Doctroints type	Count	% ¹	Vio	lated	3	Consis	tentl	${ m y~Violated^4}$
Restraints type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	369	20.4	2	0.5	0.1	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	292	16.2	2	0.7	0.1	0	0.0	0.0
Sidechain-Sidechain	77	4.3	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	455	25.2	5	1.1	0.3	0	0.0	0.0
Backbone-Backbone	171	9.5	3	1.8	0.2	0	0.0	0.0
Backbone-Sidechain	249	13.8	2	0.8	0.1	0	0.0	0.0
Sidechain-Sidechain	35	1.9	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	232	12.8	1	0.4	0.1	1	0.4	0.1
Backbone-Backbone	46	2.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	101	5.6	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	85	4.7	1	1.2	0.1	1	1.2	0.1
Long range ($ i-j \ge 5$)	685	37.9	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	132	7.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	294	16.3	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	259	14.3	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	66	3.7	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1807	100.0	8	0.4	0.4	1	0.1	0.1
Backbone-Backbone	415	23.0	3	0.7	0.2	0	0.0	0.0
Backbone-Sidechain	936	51.8	4	0.4	0.2	0	0.0	0.0
Sidechain-Sidechain	456	25.2	1	0.2	0.1	1	0.2	0.1

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID		Nun	nber o	f viola	ations	3	Maan (Å)	Max (Å)	${ m SD}^6$ (Å)	Modian (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (A)	$SD^*(A)$	Median (Å)
1	0	2	1	0	0	3	0.12	0.14	0.01	0.12
2	0	0	1	0	0	1	0.13	0.13	0.0	0.13
3	2	0	1	0	0	3	0.15	0.16	0.01	0.15
4	1	1	1	0	0	3	0.16	0.17	0.01	0.17
5	0	0	1	0	0	1	0.14	0.14	0.0	0.14
6	1	1	1	0	0	3	0.14	0.16	0.02	0.14
7	1	1	1	0	0	3	0.15	0.17	0.02	0.16
8	1	0	1	0	0	2	0.15	0.15	0.0	0.15
9	1	0	1	0	0	2	0.13	0.14	0.01	0.13
10	0	0	1	0	0	1	0.12	0.12	0.0	0.12
11	1	1	1	0	0	3	0.13	0.16	0.02	0.13

Continued on next page...




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	${ m SD}^6$ (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	2	1	1	0	0	4	0.12	0.13	0.01	0.13
13	2	1	1	0	0	4	0.14	0.16	0.01	0.14
14	0	0	1	0	0	1	0.13	0.13	0.0	0.13
15	1	1	1	0	0	3	0.13	0.15	0.02	0.14
16	0	1	1	0	0	2	0.12	0.13	0.01	0.12
17	1	0	1	0	0	2	0.12	0.13	0.01	0.12
18	1	2	1	0	0	4	0.13	0.16	0.02	0.12
19	0	1	1	0	0	2	0.13	0.15	0.02	0.13
20	1	1	1	0	0	3	0.13	0.14	0.01	0.12

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1733(IR:367, SQ:450, MR:231, LR:685, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	Fraction of the ensemble			
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
0	2	0	0	0	2	1	5.0
0	1	0	0	0	1	2	10.0
1	0	0	0	0	1	3	15.0
0	1	0	0	0	1	4	20.0
0	0	0	0	0	0	5	25.0
0	1	0	0	0	1	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
1	0	0	0	0	1	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	1	0	0	1	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,198)	1:A:90:TYR:HE1	1:A:92:PRO:HB2	20	0.14	0.01	0.14
(1,198)	1:A:90:TYR:HE1	1:A:92:PRO:HB3	20	0.14	0.01	0.14
(1,198)	1:A:90:TYR:HE2	1:A:92:PRO:HB2	20	0.14	0.01	0.14
(1,198)	1:A:90:TYR:HE2	1:A:92:PRO:HB3	20	0.14	0.01	0.14
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD11	13	0.14	0.02	0.14
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD12	13	0.14	0.02	0.14
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD13	13	0.14	0.02	0.14
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD21	13	0.14	0.02	0.14
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD22	13	0.14	0.02	0.14
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD23	13	0.14	0.02	0.14
(1,1192)	1:A:104:VAL:HB	1:A:105:SER:H	6	0.13	0.01	0.14
(1,512)	1:A:135:ALA:H	1:A:136:LYS:HA	4	0.12	0.0	0.12
(1,1213)	1:A:88:VAL:HA	1:A:88:VAL:HB	3	0.15	0.01	0.15
(1,737)	1:A:45:SER:H	1:A:46:ILE:H	2	0.14	0.03	0.14


¹Number of violated models, ²Standard deviation



9.5 All violated distance restraints (i)

9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,737)	1:A:45:SER:H	1:A:46:ILE:H	7	0.17
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD11	4	0.17
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD12	4	0.17
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD13	4	0.17
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD21	4	0.17
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD22	4	0.17
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD23	4	0.17
(1,198)	1:A:90:TYR:HE1	1:A:92:PRO:HB2	4	0.17
(1,198)	1:A:90:TYR:HE1	1:A:92:PRO:HB3	4	0.17
(1,198)	1:A:90:TYR:HE2	1:A:92:PRO:HB2	4	0.17
(1,198)	1:A:90:TYR:HE2	1:A:92:PRO:HB3	4	0.17

Continued on next page...



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD11	3	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD12	3	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD13	3	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD21	3	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD22	3	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD23	3	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD11	7	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD12	7	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD13	7	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD21	7	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD22	7	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD23	7	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD11	18	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD12	18	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD13	18	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD21	18	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD22	18	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD23	18	0.16
(1,198)	1:A:90:TYR:HE1	1:A:92:PRO:HB2	6	0.16
(1,198)	1:A:90:TYR:HE1	1:A:92:PRO:HB3	6	0.16
(1,198)	1:A:90:TYR:HE2	1:A:92:PRO:HB2	6	0.16
(1,198)	1:A:90:TYR:HE2	1:A:92:PRO:HB3	6	0.16
(1,198)	1:A:90:TYR:HE1	1:A:92:PRO:HB2	11	0.16
(1,198)	1:A:90:TYR:HE1	1:A:92:PRO:HB3	11	0.16
(1,198)	1:A:90:TYR:HE2	1:A:92:PRO:HB2	11	0.16
(1,198)	1:A:90:TYR:HE2	1:A:92:PRO:HB3	11	0.16
(1,1213)	1:A:88:VAL:HA	1:A:88:VAL:HB	13	0.16
(1,590)	1:A:111:LEU:HA	1:A:111:LEU:HD11	8	0.15



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

A 1 - 4	Commit	% ¹	${f Violated}^3$			Consistently Violated ⁴		
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
CHI1	9	6.4	1	11.1	0.7	0	0.0	0.0
PSI	59	42.1	2	3.4	1.4	2	3.4	1.4
	4	2.9	4	100.0	2.9	4	100.0	2.9
PHI	68	48.6	1	1.5	0.7	0	0.0	0.0
Total	140	100.0	8	5.7	5.7	6	4.3	4.3

 $^{^1}$ percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Nui	$\overline{\mathbf{mber}}$	of	violati	ions	Mean (°)	Max (°)	SD (°)	Median (°)
Wiodei 1D	CHI1	PSI		PHI	Total	Mean ()	Max ()	SD ()	Median ()
1	0	2	4	0	6	54.27	91.6	38.05	72.8
2	0	2	4	0	6	56.33	96.4	40.75	71.15
3	0	2	4	0	6	56.62	97.0	39.4	75.35
4	0	2	4	0	6	56.65	102.1	40.18	71.6
5	0	2	4	0	6	57.58	102.1	41.58	72.25
6	0	2	4	0	6	53.68	90.2	38.13	69.35
7	0	2	4	0	6	57.85	94.2	40.24	78.65
8	0	2	4	0	6	54.45	94.1	38.54	70.95
9	0	2	4	0	6	54.7	91.3	38.67	72.75
10	0	2	4	0	6	53.75	91.6	37.52	71.05
11	0	2	4	0	6	58.43	104.6	41.32	76.8
12	0	2	4	0	6	58.6	99.4	41.97	76.7
13	0	2	4	0	6	56.08	88.5	38.89	78.2
14	0	2	4	0	6	56.92	93.5	39.46	78.3
15	0	2	4	1	7	49.8	91.1	42.14	74.5
16	1	2	4	0	7	50.71	105.1	43.56	70.0
17	0	2	4	0	6	54.13	91.5	37.93	70.05
18	0	2	4	0	6	54.17	99.0	40.58	66.65
19	0	2	4	0	6	55.28	92.6	38.75	71.65
20	0	2	4	0	6	53.05	88.3	37.37	69.9



10.2.1 Bar graph: Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Numb	oer of	vio	olated	restraints	Fraction	n of the ensemble
CHI1	PSI		PHI	Total	Count ¹	%
1	0	0	1	2	1	5.0
0	0	0	0	0	2	10.0
0	0	0	0	0	3	15.0
0	0	0	0	0	4	20.0
0	0	0	0	0	5	25.0
0	0	0	0	0	6	30.0
0	0	0	0	0	7	35.0
0	0	0	0	0	8	40.0
0	0	0	0	0	9	45.0
0	0	0	0	0	10	50.0
0	0	0	0	0	11	55.0

Continued on next page...




Continued from previous page...

Numb	oer of	vio	olated	restraints	Fractio	n of the ensemble
CHI1	PSI		PHI	Total	Count ¹	%
0	0	0	0	0	12	60.0
0	0	0	0	0	13	65.0
0	0	0	0	0	14	70.0
0	0	0	0	0	15	75.0
0	0	0	0	0	16	80.0
0	0	0	0	0	17	85.0
0	0	0	0	0	18	90.0
0	0	0	0	0	19	95.0
0	2	4	0	6	20	100.0

¹ Number of models with violations

10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram: Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models



in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	$\mid \mathbf{SD}^2 \mid$	Media
(1,3)	1:A:55:THR:N	1:A:55:THR:CA	1:A:55:THR:CB	1:A:55:THR:CG2	20	91.46	9.31	91.7
(1,5)	1:A:108:THR:N	1:A:108:THR:CA	1:A:108:THR:CB	1:A:108:THR:CG2	20	90.99	2.57	90.9
(1,4)	1:A:70:THR:N	1:A:70:THR:CA	1:A:70:THR:CB	1:A:70:THR:CG2	20	84.37	5.96	84.75
(1,6)	1:A:117:THR:N	1:A:117:THR:CA	1:A:117:THR:CB	1:A:117:THR:CG2	20	65.27	7.73	66.55
(1,101)	1:A:107:GLY:N	1:A:107:GLY:CA	1:A:107:GLY:C	1:A:108:THR:N	20	2.07	0.16	2.05
(1,79)	1:A:92:PRO:N	1:A:92:PRO:CA	1:A:92:PRO:C	1:A:93:VAL:N	20	1.68	0.24	1.6


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram: Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,3)	1:A:55:THR:N	1:A:55:THR:CA	1:A:55:THR:CB	1:A:55:THR:CG2	16	105.1
(1,3)	1:A:55:THR:N	1:A:55:THR:CA	1:A:55:THR:CB	1:A:55:THR:CG2	11	104.6
(1,3)	1:A:55:THR:N	1:A:55:THR:CA	1:A:55:THR:CB	1:A:55:THR:CG2	4	102.1
(1,3)	1:A:55:THR:N	1:A:55:THR:CA	1:A:55:THR:CB	1:A:55:THR:CG2	5	102.1
(1,3)	1:A:55:THR:N	1:A:55:THR:CA	1:A:55:THR:CB	1:A:55:THR:CG2	12	99.4
(1,4)	1:A:70:THR:N	1:A:70:THR:CA	1:A:70:THR:CB	1:A:70:THR:CG2	18	99.0
(1,3)	1:A:55:THR:N	1:A:55:THR:CA	1:A:55:THR:CB	1:A:55:THR:CG2	3	97.0
(1,5)	1:A:108:THR:N	1:A:108:THR:CA	1:A:108:THR:CB	1:A:108:THR:CG2	2	96.4
(1,5)	1:A:108:THR:N	1:A:108:THR:CA	1:A:108:THR:CB	1:A:108:THR:CG2	12	95.7
(1,3)	1:A:55:THR:N	1:A:55:THR:CA	1:A:55:THR:CB	1:A:55:THR:CG2	2	95.7

