



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 03:06 pm BST

PDB ID : 2J62
Title : Structure of a bacterial O-glcnacase in complex with glcnacstatin
Authors : Dorfmueller, H.C.; Borodkin, V.S.; Schimpl, M.; Shepherd, S.M.; Shpiro, N.A.;
van Aalten, D.M.F.
Deposited on : 2006-09-22
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	594	 2% 83% 13% ...
1	B	594	 2% 85% 12% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9774 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called O-GlcNAcase NagJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	585	4628	2913	756	942	17	0	0	0
1	B	585	4629	2913	757	942	17	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	388	ASP	ASN	conflict	UNP Q0TR53
B	388	ASP	ASN	conflict	UNP Q0TR53

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Cl	0	0
			2	2		
2	A	2	Total	Cl	0	0
			2	2		

- Molecule 3 is N-[(5R,6R,7R,8S)-6,7-DIHYDROXY-5-(HYDROXYMETHYL)-2-(2-PHENYLETHYL)-1,5,6,7,8,8A-HEXAHYDROIMIDAZO[1,2-A]PYRIDIN-8-YL]-2-METHYLPROPANAMIDE (three-letter code: GSZ) (formula: C₂₀H₂₈N₃O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	27	20	3	4	0	0
3	B	1	27	20	3	4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	198	198	198	0	0
4	B	261	261	261	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: O-GlcNAcase NagJ



- Molecule 1: O-GlcNAcase NagJ



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	130.25Å 145.92Å 152.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.26 20.00 – 2.26	Depositor EDS
% Data completeness (in resolution range)	99.6 (20.00-2.26) 99.6 (20.00-2.26)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.80 (at 2.26Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.179 , 0.219 0.178 , 0.217	Depositor DCC
R_{free} test set	1372 reflections (2.02%)	wwPDB-VP
Wilson B-factor (Å ²)	40.9	Xtrriage
Anisotropy	0.051	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.007 for -h,-l,-k	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9774	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GSZ, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	4/4724 (0.1%)	0.68	3/6413 (0.0%)
1	B	0.64	0/4725	0.70	6/6415 (0.1%)
All	All	0.67	4/9449 (0.0%)	0.69	9/12828 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	140	GLY	C-O	14.29	1.46	1.23
1	A	140	GLY	C-N	8.56	1.53	1.34
1	A	67	ASN	CG-ND2	7.71	1.52	1.32
1	A	67	ASN	CG-OD1	6.18	1.37	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	502	ARG	NE-CZ-NH1	9.11	124.85	120.30
1	B	584	GLU	N-CA-C	-7.61	90.45	111.00
1	B	502	ARG	NE-CZ-NH2	-7.04	116.78	120.30
1	A	502	ARG	NE-CZ-NH2	-6.53	117.03	120.30
1	A	98	ASP	C-N-CD	-5.47	108.57	120.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	502	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	B	284	LEU	CA-CB-CG	-5.41	102.86	115.30
1	B	555	LEU	CA-CB-CG	5.22	127.31	115.30
1	B	126	GLU	C-N-CA	5.19	134.67	121.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	99	PRO	Peptide
1	B	127	ASN	Peptide
1	B	583	TYR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4628	0	4416	60	0
1	B	4629	0	4418	46	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	27	0	28	0	0
3	B	27	0	28	0	0
4	A	198	0	0	4	0
4	B	261	0	0	1	2
All	All	9774	0	8890	104	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:PRO:CB	1:A:100:ASN:HB2	1.73	1.18
1:A:99:PRO:HB2	1:A:100:ASN:HB2	1.22	1.11

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:PRO:CB	1:A:100:ASN:CB	2.49	0.89
1:B:502:ARG:HD3	1:B:502:ARG:O	1.74	0.87
1:A:99:PRO:HB3	1:A:100:ASN:HB2	1.56	0.85
1:A:99:PRO:HB2	1:A:100:ASN:CB	2.04	0.82
1:A:515:LYS:HG2	1:A:515:LYS:O	1.77	0.81
1:A:99:PRO:HB3	1:A:100:ASN:CB	2.14	0.77
1:A:237:GLN:HG3	4:A:2046:HOH:O	1.84	0.76
1:B:47:ASN:HD21	1:B:446:ASP:HB2	1.53	0.72
1:B:359:PRO:HA	1:B:389:ARG:HH21	1.56	0.71
1:A:108:GLU:OE1	4:A:2009:HOH:O	2.08	0.69
1:A:41:VAL:HG13	1:A:59:GLY:HA3	1.78	0.65
1:A:502:ARG:O	1:A:502:ARG:HD3	1.97	0.64
1:A:99:PRO:HB3	1:A:100:ASN:CG	2.17	0.63
1:B:487:ASN:HD21	1:B:491:ALA:H	1.46	0.62
1:B:156:VAL:O	1:B:160:LYS:HG3	2.01	0.60
1:B:502:ARG:HD3	1:B:502:ARG:C	2.18	0.60
1:A:619:PHE:HD2	1:A:621:LEU:HD22	1.66	0.59
1:A:502:ARG:C	1:A:502:ARG:HD3	2.22	0.59
1:A:68:LEU:HG	1:A:71:GLU:HG3	1.82	0.59
1:A:619:PHE:CD2	1:A:621:LEU:CD2	2.86	0.58
1:B:47:ASN:ND2	1:B:446:ASP:HB2	2.20	0.57
1:A:99:PRO:HB3	1:A:100:ASN:OD1	2.05	0.56
1:A:80:ASN:O	1:A:84:GLU:HG3	2.04	0.56
1:A:107:GLY:O	1:A:145:GLU:HA	2.06	0.56
1:B:285:TYR:CZ	1:B:322:LYS:HD3	2.41	0.56
1:A:420:ASN:H	1:A:420:ASN:HD22	1.54	0.55
1:A:619:PHE:CD2	1:A:621:LEU:HD22	2.42	0.55
1:B:420:ASN:H	1:B:420:ASN:HD22	1.56	0.54
1:B:500:GLU:CD	1:B:500:GLU:H	2.11	0.54
1:A:52:ASN:O	1:A:173:ILE:HA	2.07	0.54
1:A:538:ASN:OD1	1:A:542:ASN:ND2	2.41	0.54
1:A:343:ASN:HB3	4:A:2076:HOH:O	2.08	0.54
1:A:387:TYR:HB3	1:A:389:ARG:HD2	1.89	0.53
1:B:368:PRO:HD2	1:B:372:THR:HG21	1.90	0.53
1:B:66:ILE:HG22	1:B:102:THR:HB	1.90	0.53
1:A:67:ASN:HB2	1:A:103:THR:HG23	1.91	0.53
1:A:42:LEU:H	1:A:42:LEU:HD12	1.72	0.53
1:B:126:GLU:N	1:B:127:ASN:HB3	2.25	0.52
1:B:285:TYR:CE1	1:B:322:LYS:HD3	2.44	0.52
1:A:504:LYS:HE2	1:A:524:GLU:OE1	2.10	0.52
1:B:126:GLU:N	1:B:127:ASN:CB	2.73	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:619:PHE:CD2	1:A:621:LEU:HD21	2.45	0.52
1:B:487:ASN:ND2	1:B:491:ALA:H	2.07	0.51
1:A:331:VAL:HG22	1:A:364:MET:HB2	1.92	0.51
1:A:392:ALA:HA	1:A:425:PHE:HB3	1.92	0.51
1:A:624:ILE:HD12	1:B:224:ARG:HH21	1.76	0.51
1:A:133:TYR:CE2	1:A:175:ASP:HB3	2.45	0.50
1:B:295:TYR:CD2	1:B:331:VAL:HG13	2.46	0.50
1:A:398:PRO:HD2	1:A:430:PRO:HA	1.93	0.50
1:B:557:GLU:HG2	1:B:604:ILE:HG22	1.93	0.50
1:A:65:SER:HB3	1:A:92:GLU:HB3	1.93	0.49
1:B:85:PHE:HB2	1:B:160:LYS:HD2	1.94	0.49
1:A:183:GLY:C	1:A:184:ILE:HD12	2.32	0.49
1:B:359:PRO:HA	1:B:389:ARG:NH2	2.23	0.49
1:B:183:GLY:O	1:B:427:THR:HA	2.13	0.49
1:B:364:MET:HA	1:B:392:ALA:O	2.12	0.48
1:B:315:ASN:O	1:B:320:LYS:HG3	2.13	0.48
1:A:183:GLY:O	1:A:427:THR:HA	2.13	0.48
1:A:295:TYR:CE2	1:A:331:VAL:HG23	2.48	0.48
1:B:295:TYR:CD2	1:B:331:VAL:CG1	2.98	0.47
1:B:387:TYR:HB3	1:B:389:ARG:HG2	1.97	0.46
1:A:85:PHE:CD1	1:A:160:LYS:HG2	2.51	0.46
1:B:387:TYR:HB3	1:B:389:ARG:HH11	1.80	0.46
1:B:154:TYR:CE1	1:B:209:LYS:HA	2.51	0.45
1:A:557:GLU:HG2	1:A:602:ALA:HB3	1.96	0.45
1:A:68:LEU:HG	1:A:71:GLU:CG	2.45	0.45
1:B:193:TRP:CZ3	1:B:201:GLN:HG3	2.51	0.45
1:B:184:ILE:HD11	1:B:441:ILE:HG23	1.98	0.45
1:A:505:MET:HG2	1:A:612:PHE:CD1	2.52	0.44
1:B:75:ASP:OD1	1:B:77:ASN:HB2	2.17	0.44
1:B:61:LYS:HE2	1:B:166:SER:HB2	2.00	0.44
1:A:99:PRO:HB2	1:A:100:ASN:CA	2.48	0.44
1:A:194:THR:O	1:A:197:ASP:HB2	2.17	0.44
1:A:98:ASP:HA	1:A:99:PRO:O	2.18	0.43
1:B:265:PHE:CD2	1:B:306:LYS:HD3	2.52	0.43
1:B:177:PRO:HB3	1:B:449:TRP:CE3	2.53	0.43
1:A:588:GLU:HG2	4:A:2048:HOH:O	2.17	0.43
1:A:515:LYS:HG3	1:B:300:GLN:OE1	2.19	0.43
1:B:337:THR:HB	1:B:368:PRO:HA	2.01	0.43
1:B:387:TYR:HD2	1:B:389:ARG:NH1	2.16	0.43
1:A:416:ASP:HB3	1:A:419:LEU:HD13	2.01	0.42
1:B:304:ALA:HB2	1:B:349:TYR:CD1	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:HIS:HD2	1:A:256:GLY:O	2.02	0.42
1:B:154:TYR:CD1	1:B:209:LYS:HG2	2.54	0.42
1:B:231:TYR:HB2	1:B:236:MET:SD	2.59	0.42
1:A:98:ASP:OD1	1:A:99:PRO:O	2.38	0.42
1:A:274:PHE:CE2	1:A:313:ARG:HD2	2.55	0.42
1:B:209:LYS:HD3	1:B:449:TRP:CH2	2.55	0.42
1:A:41:VAL:CG1	1:A:59:GLY:HA3	2.49	0.42
1:A:274:PHE:CZ	1:A:313:ARG:HD2	2.56	0.41
1:B:478:PHE:CE1	1:B:555:LEU:HD13	2.55	0.41
1:A:183:GLY:HA3	1:A:212:THR:O	2.21	0.41
1:A:337:THR:HB	1:A:368:PRO:HA	2.01	0.41
1:B:476:LYS:O	1:B:480:ASN:HB2	2.21	0.41
1:A:433:HIS:HB3	1:A:550:GLU:OE1	2.20	0.41
1:A:97:ASN:OD1	1:A:98:ASP:N	2.54	0.41
1:B:420:ASN:HD22	1:B:420:ASN:N	2.19	0.41
1:A:561:LEU:HD22	1:A:610:GLN:HA	2.04	0.40
1:A:114:PRO:O	1:A:118:GLU:HG2	2.21	0.40
1:A:165:GLU:O	1:A:166:SER:HB2	2.21	0.40
1:B:47:ASN:ND2	4:B:2004:HOH:O	2.54	0.40
1:B:557:GLU:CG	1:B:604:ILE:HG22	2.51	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:2241:HOH:O	4:B:2241:HOH:O[6_555]	1.97	0.23
4:B:2237:HOH:O	4:B:2241:HOH:O[6_555]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	583/594 (98%)	559 (96%)	21 (4%)	3 (0%)	29	29
1	B	583/594 (98%)	558 (96%)	21 (4%)	4 (1%)	22	21
All	All	1166/1188 (98%)	1117 (96%)	42 (4%)	7 (1%)	25	25

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	99	PRO
1	B	128	LEU
1	B	127	ASN
1	B	388	ASP
1	A	98	ASP
1	A	343	ASN
1	B	41	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	500/507 (99%)	469 (94%)	31 (6%)	18	17
1	B	500/507 (99%)	475 (95%)	25 (5%)	24	26
All	All	1000/1014 (99%)	944 (94%)	56 (6%)	21	21

All (56) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	VAL
1	A	42	LEU
1	A	66	ILE
1	A	72	GLU
1	A	86	LEU
1	A	98	ASP
1	A	129	LYS
1	A	180	SER
1	A	237	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	257	ILE
1	A	284	LEU
1	A	313	ARG
1	A	328	LEU
1	A	355	GLU
1	A	358	ASP
1	A	365	TRP
1	A	372	THR
1	A	389	ARG
1	A	394	TRP
1	A	409	LEU
1	A	420	ASN
1	A	482	SER
1	A	492	LYS
1	A	502	ARG
1	A	515	LYS
1	A	555	LEU
1	A	556	ASP
1	A	588	GLU
1	A	592	ASN
1	A	621	LEU
1	A	623	LEU
1	B	40	GLN
1	B	41	VAL
1	B	86	LEU
1	B	113	ILE
1	B	121	ASN
1	B	123	THR
1	B	128	LEU
1	B	174	THR
1	B	233	GLU
1	B	284	LEU
1	B	313	ARG
1	B	328	LEU
1	B	331	VAL
1	B	365	TRP
1	B	409	LEU
1	B	420	ASN
1	B	458	LYS
1	B	487	ASN
1	B	492	LYS
1	B	515	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	555	LEU
1	B	580	THR
1	B	581	GLU
1	B	619	PHE
1	B	624	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	ASN
1	A	201	GLN
1	A	223	HIS
1	A	390	ASN
1	A	420	ASN
1	A	610	GLN
1	B	47	ASN
1	B	77	ASN
1	B	121	ASN
1	B	420	ASN
1	B	487	ASN
1	B	610	GLN
1	B	614	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GSZ	B	1627	-	25,29,29	1.18	1 (4%)	28,41,41	1.33	1 (3%)
3	GSZ	A	1626	-	25,29,29	1.11	2 (8%)	28,41,41	1.37	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GSZ	B	1627	-	-	1/14/35/35	0/2/3/3
3	GSZ	A	1626	-	-	5/14/35/35	0/2/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1627	GSZ	CAQ-NAE	-3.21	1.33	1.38
3	A	1626	GSZ	CAQ-NAE	-2.96	1.33	1.38
3	A	1626	GSZ	CAC-NAK	2.22	1.50	1.46

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1627	GSZ	CAQ-NAE-CAD	-5.03	106.21	109.31
3	A	1626	GSZ	CAQ-NAE-CAD	-5.02	106.22	109.31
3	A	1626	GSZ	CAB-CAA-CAF	-2.13	107.76	111.37

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1627	GSZ	CAD-CAC-NAK-CAL
3	A	1626	GSZ	CAD-CAC-NAK-CAL
3	A	1626	GSZ	NAK-CAL-CAN-CAZ

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	1626	GSZ	OAM-CAL-CAN-CAZ
3	A	1626	GSZ	OAM-CAL-CAN-CBA
3	A	1626	GSZ	NAK-CAL-CAN-CBA

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	585/594 (98%)	-0.10	11 (1%) 66 69	35, 44, 54, 62	10 (1%)
1	B	585/594 (98%)	-0.09	14 (2%) 59 62	32, 44, 55, 66	8 (1%)
All	All	1170/1188 (98%)	-0.09	25 (2%) 63 66	32, 44, 54, 66	18 (1%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	41	VAL	18.6
1	B	144	ILE	4.8
1	B	122	GLY	3.9
1	B	121	ASN	3.7
1	A	112	ASP	3.3
1	A	624	ILE	3.2
1	A	58	ASP	3.1
1	B	343	ASN	3.0
1	B	118	GLU	3.0
1	B	143	ALA	2.9
1	B	99	PRO	2.9
1	A	409	LEU	2.8
1	A	106	ILE	2.8
1	A	517	ASP	2.7
1	B	142	ILE	2.6
1	A	104	LEU	2.5
1	B	90	ASN	2.4
1	A	580	THR	2.2
1	B	127	ASN	2.2
1	B	58	ASP	2.2
1	A	144	ILE	2.1
1	A	444	ALA	2.1
1	A	98	ASP	2.1
1	B	135	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	344	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	B	1626	1/1	0.81	0.15	67,67,67,67	0
2	CL	A	1625	1/1	0.91	0.08	75,75,75,75	0
2	CL	A	1627	1/1	0.91	0.04	67,67,67,67	0
3	GSZ	B	1627	27/27	0.94	0.16	37,44,62,64	0
3	GSZ	A	1626	27/27	0.96	0.12	37,42,53,54	0
2	CL	B	1625	1/1	0.96	0.06	67,67,67,67	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GSZ B 1627:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GSZ A 1626:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.