

wwPDB NMR Structure Validation Summary Report (i)

Jun 5, 2023 – 07:59 AM EDT

PDB ID	:	2M2S
BMRB ID	:	18931
Title	:	Solution structure of the antimicrobial peptide [Aba5,7,12,14]BTD-2
Authors	:	Conibear, A.C.; Rosengren, K.; Daly, N.L.; Troiera Henriques, S.; Craik, D.J.
Deposited on	:	2013-01-02

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 66%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f NMR} { m archive} \ (\#{ m Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain			
1	А	18	78%	22%		



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues						
Well-defined core	Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1	A:1-A:4, A:6-A:6, A:8-A:11,	0.83	10			
	A:13-A:13, A:15-A:18 (14)					

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	2, 4, 10, 11, 14, 15, 17, 18, 19
2	1, 5, 6, 8
3	3, 9, 12, 16
4	7, 13, 20



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 294 atoms, of which 156 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called [Aba5,7,12,14]BTD-2.

Mol	Chain	Residues	Atoms					Trace		
1 Λ	1 A		10	Total	С	Н	Ν	0	S	0
1			294	82	156	36	18	2	0	



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: [Aba5,7,12,14]BTD-2

Chain A:	78%	22%
G1 A5 A7 A7 A12 A12 A13 A14 A18 A18		

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 10. Colouring as in section 4.1 above.

• Molecule 1: [Aba5,7,12,14]BTD-2

Chain A: 72% 6% 22%



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing, torsion angle dynamics.*

Of the 50 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version	
CYANA	structure solution	3.0	
CNS	refinement	2.1	
CYANA	refinement	3.0	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	185
Number of shifts mapped to atoms	185
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	66%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ABA

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	114	128	128	0 ± 0
All	All	2280	2560	2560	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models		
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total	
1:A:8:ARG:NE	1:A:8:ARG:HA	0.59	2.13	8	1	
1:A:9:ARG:NE	1:A:9:ARG:HA	0.48	2.24	19	1	
1:A:18:ARG:HA	1:A:18:ARG:NE	0.43	2.29	11	1	
1:A:3:CYS:HA	1:A:15:VAL:O	0.41	2.16	20	1	

All unique clashes are listed below, sorted by their clash magnitude.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	12/18~(67%)	$11 \pm 1 (95 \pm 7\%)$	$1\pm1 (5\pm7\%)$	0±0 (0±0%)	100 100
All	All	240/360~(67%)	229~(95%)	11 (5%)	0 (0%)	100 100

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

There are no Ramachandran outliers.

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the side chain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	12/12~(100%)	$12\pm0~(100\pm2\%)$	0±0 (0±2%)	91 98
All	All	240/240~(100%)	239~(100%)	1 (0%)	91 98

All 1 unique residues with a non-rotameric sidechain are listed below.

Mol	Chain	Res	Type	Models (Total)
1	А	13	ARG	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.



Mol	Type	Chain	Res	Link	Bond		ngths
WIOI	туре	Ullalli	nes	LIIIK	Counts	RMSZ	#Z>2
1	ABA	А	5	1	4,5,6	$0.91{\pm}0.11$	0±0 (0±0%)
1	ABA	А	14	1	4,5,6	$0.82 {\pm} 0.06$	0±0 (0±0%)
1	ABA	А	7	1	4,5,6	$0.99 {\pm} 0.15$	0 ± 0 (6±10%)
1	ABA	А	12	1	4,5,6	$0.83 {\pm} 0.09$	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
	туре	Ullaili	nes	LIIIK	Counts	RMSZ	#Z>2
1	ABA	А	5	1	$1,\!5,\!7$	$0.28 {\pm} 0.09$	0±0 (0±0%)
1	ABA	А	14	1	$1,\!5,\!7$	0.25 ± 0.11	0±0 (0±0%)
1	ABA	А	7	1	$1,\!5,\!7$	0.29 ± 0.21	0±0 (0±0%)
1	ABA	А	12	1	$1,\!5,\!7$	$0.33 {\pm} 0.20$	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ABA	А	14	1	-	$0\pm 0,3,4,6$	-
1	ABA	А	12	1	-	$0\pm 0,3,4,6$	-
1	ABA	А	7	1	-	$0\pm0,3,4,6$	-
1	ABA	А	5	1	-	$0\pm0,3,4,6$	-

All unique bond outliers are listed below.

Mol	Chain	ain Res	Type	Atoms Z		$Observed(\lambda)$	Ideal(Å)	Moo	lels
	Ullalli		es Type	Atoms		Observed(A)	Iucai(A)	Worst	Total
1	А	7	ABA	CA-N	2.27	1.41	1.48	13	5

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 66% for the well-defined parts and 66% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	185
Number of shifts mapped to atoms	185
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 66%, i.e. 149 atoms were assigned a chemical shift out of a possible 226. 0 out of 4 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	43/72~(60%)	29/30~(97%)	14/28~(50%)	0/14~(0%)
Sidechain	106/154~(69%)	74/98~(76%)	32/38~(84%)	0/18~(0%)
Overall	149/226~(66%)	103/128~(80%)	46/66~(70%)	0/32~(0%)

7.1.4 Statistically unusual chemical shifts (i)


There are no statistically unusual chemical shifts.



7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	108
Intra-residue $(i-j =0)$	52
Sequential (i-j =1)	28
Medium range ($ i-j >1$ and $ i-j <5$)	6
Long range $(i-j \ge 5)$	10
Inter-chain	0
Hydrogen bond restraints	12
Disulfide bond restraints	0
Total dihedral-angle restraints	32
Number of unmapped restraints	3
Number of restraints per residue	7.8
Number of long range restraints per residue ¹	1.2

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	0.5	0.15
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	Max $(^{\circ})$
1.0-10.0 (Small)	0.1	1.9
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destructures to me	Count	$\%^1$	Vic	lated	3	Consis	tently	y Violated ⁴
Restraints type	Count	701	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$
Intra-residue (i-j =0)	52	48.1	2	3.8	1.9	0	0.0	0.0
Backbone-Backbone	2	1.9	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	49	45.4	2	4.1	1.9	0	0.0	0.0
Sidechain-Sidechain	1	0.9	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	28	25.9	1	3.6	0.9	0	0.0	0.0
Backbone-Backbone	18	16.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	10	9.3	1	10.0	0.9	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j > 1 \& i-j < 5$)	6	5.6	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	2	1.9	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	2	1.9	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	2	1.9	0	0.0	0.0	0	0.0	0.0
Long range $(i-j \ge 5)$	10	9.3	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	9	8.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	1	0.9	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	12	11.1	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	108	100.0	3	2.8	2.8	0	0.0	0.0
Backbone-Backbone	43	39.8	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	62	57.4	3	4.8	2.8	0	0.0	0.0
Sidechain-Sidechain	3	2.8	0	0.0	0.0	0	0.0	0.0

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID		Nun	nber o	f viola	ations	5	Mean (Å)	Mor (Å)	SD^6 (Å)	Median (Å)
Model ID	IR^{1}	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Mean (A)	Max (Å)	$SD^{*}(A)$	Median (A)
1	0	0	0	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	1	0	0	0	0	1	0.13	0.13	0.0	0.13
4	0	0	0	0	0	0	0.0	0.0	0.0	0.0
5	0	1	0	0	0	1	0.14	0.14	0.0	0.14
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	1	0	0	0	1	0.14	0.14	0.0	0.14
8	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	1	0	0	0	0	1	0.11	0.11	0.0	0.11
10	0	1	0	0	0	1	0.14	0.14	0.0	0.14
11	0	0	0	0	0	0	0.0	0.0	0.0	0.0

Continued on next page...




Madal ID			nber o		ations	5				
Model ID	IR^1	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Mean (Å)	Max (Å)	\mathbf{SD}^{6} (Å)	Median (Å)
12	0	1	0	0	0	1	0.11	0.11	0.0	0.11
13	0	0	0	0	0	0	0.0	0.0	0.0	0.0
14	1	0	0	0	0	1	0.15	0.15	0.0	0.15
15	0	0	0	0	0	0	0.0	0.0	0.0	0.0
16	0	1	0	0	0	1	0.15	0.15	0.0	0.15
17	0	0	0	0	0	0	0.0	0.0	0.0	0.0
18	0	0	0	0	0	0	0.0	0.0	0.0	0.0
19	0	1	0	0	0	1	0.14	0.14	0.0	0.14
20	0	1	0	0	0	1	0.13	0.13	0.0	0.13

Continued from previous page..

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation





The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right




9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 93(IR:50, SQ:27, MR:6, LR:10, IC:0) restraints are not violated in the ensemble.

Nu		of vio		restra	aints	Fractio	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC ⁵	Total	Count^6	%
1	0	0	0	0	1	1	5.0
1	0	0	0	0	1	2	10.0
0	0	0	0	0	0	3	15.0
0	0	0	0	0	0	4	20.0
0	0	0	0	0	0	5	25.0
0	0	0	0	0	0	6	30.0
0	1	0	0	0	1	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations






9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,36)	1:A:6:VAL:HB	1:A:7:ABA:H	7	0.14	0.01	0.14
(1,15)	1:A:7:ABA:H	1:A:7:ABA:HB2	2	0.13	0.02	0.13


¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,36)	1:A:6:VAL:HB	1:A:7:ABA:H	16	0.15
(1,15)	1:A:7:ABA:H	1:A:7:ABA:HB2	14	0.15
(1,36)	1:A:6:VAL:HB	1:A:7:ABA:H	5	0.14
(1,36)	1:A:6:VAL:HB	1:A:7:ABA:H	7	0.14
(1,36)	1:A:6:VAL:HB	1:A:7:ABA:H	10	0.14
(1,36)	1:A:6:VAL:HB	1:A:7:ABA:H	19	0.14
(1,36)	1:A:6:VAL:HB	1:A:7:ABA:H	20	0.13
(1,14)	1:A:7:ABA:H	1:A:7:ABA:HB3	3	0.13
(1,36)	1:A:6:VAL:HB	1:A:7:ABA:H	12	0.11
(1,15)	1:A:7:ABA:H	1:A:7:ABA:HB2	9	0.11



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	$\%^1$	$_1$ Violated ³			Consistently Violated ⁴			
Angle type	Count	70	Count	$\%^2$	$\%^{1}$	Count	$\%^2$	$\%^1$	
PSI	12	37.5	1	8.3	3.1	0	0.0	0.0	
PHI	16	50.0	2	12.5	6.2	0	0.0	0.0	
CHI1	4	12.5	0	0.0	0.0	0	0.0	0.0	
Total	32	100.0	3	9.4	9.4	0	0.0	0.0	

 1 percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Nui	nber (of viola	tions	Mean (°)	M_{OV} (°)	SD (°)	Median (°)
Model ID	PSI	PHI	CHI1	Total	Mean ()	$Max (^{\circ})$		Median ()
1	0	1	0	1	1.1	1.1	0.0	1.1
2	0	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	0	0.0	0.0	0.0	0.0
5	0	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0	0.0	0.0	0.0	0.0
7	1	0	0	1	1.9	1.9	0.0	1.9
8	0	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0	0.0	0.0	0.0	0.0
10	0	0	0	0	0.0	0.0	0.0	0.0
11	0	0	0	0	0.0	0.0	0.0	0.0
12	0	1	0	1	1.3	1.3	0.0	1.3
13	0	0	0	0	0.0	0.0	0.0	0.0
14	0	0	0	0	0.0	0.0	0.0	0.0
15	0	0	0	0	0.0	0.0	0.0	0.0
16	0	0	0	0	0.0	0.0	0.0	0.0
17	0	0	0	0	0.0	0.0	0.0	0.0
18	0	0	0	0	0.0	0.0	0.0	0.0
19	0	0	0	0	0.0	0.0	0.0	0.0
20	0	0	0	0	0.0	0.0	0.0	0.0



10.2.1 Bar graph : Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Nun	nber o	f violat	ted restraints	Fractio	n of the ensemble
PSI	PHI	CHI1	Total	Count^1	%
1	2	0	3	1	5.0
0	0	0	0	2	10.0
0	0	0	0	3	15.0
0	0	0	0	4	20.0
0	0	0	0	5	25.0
0	0	0	0	6	30.0
0	0	0	0	7	35.0
0	0	0	0	8	40.0
0	0	0	0	9	45.0
0	0	0	0	10	50.0
0	0	0	0	11	55.0

Continued on next page...





<u>9</u>	195
211	120

	0	1	ted restraints	Fractio	n of the ensemble
PSI	PHI	CHI1	Total	Count^1	%
0	0	0	0	12	60.0
0	0	0	0	13	65.0
0	0	0	0	14	70.0
0	0	0	0	15	75.0
0	0	0	0	16	80.0
0	0	0	0	17	85.0
0	0	0	0	18	90.0
0	0	0	0	19	95.0
0	0	0	0	20	100.0

Continued from previous page...

 1 Number of models with violations





10.4 Most violated dihedral-angle restraints in the ensemble (i)

No violations found



10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.

Data insufficient to plot histogram

10.5.2 Table: All violated dihedral-angle restraints (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,30)	1:A:17:ARG:N	1:A:17:ARG:CA	1:A:17:ARG:C	1:A:18:ARG:N	7	1.9
(1,6)	1:A:3:CYS:C	1:A:4:ARG:N	1:A:4:ARG:CA	1:A:4:ARG:C	12	1.3
(1,12)	1:A:6:VAL:C	1:A:7:ABA:N	1:A:7:ABA:CA	1:A:7:ABA:C	1	1.1

