



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 11, 2021 – 07:50 PM EDT

PDB ID : 2NTI
Title : Crystal structure of PCNA123 heterotrimer.
Authors : Hlinkova, V.; Ling, H.
Deposited on : 2006-11-07
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	 5% 74% 20% 5%
1	D	249	 3% 73% 22% 5%
1	G	249	 6% 64% 31% 5%
2	B	246	 2% 78% 18% 5%
2	E	246	 3% 77% 19% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	246	
3	C	244	
3	F	244	
3	I	244	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BR	D	3039	-	-	X	-
4	BR	F	3024	-	-	X	-
4	BR	I	3032	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 18488 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase sliding clamp B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	249	Total 1928	C 1227	N 310	O 382	S 9	0	0	0
1	A	248	Total 1914	C 1218	N 309	O 378	S 9	0	0	0
1	G	249	Total 1924	C 1225	N 310	O 380	S 9	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	2	VAL	PHE	engineered mutation	UNP P57766
A	2	VAL	PHE	engineered mutation	UNP P57766
G	2	VAL	PHE	engineered mutation	UNP P57766

- Molecule 2 is a protein called DNA polymerase sliding clamp C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	244	Total 1931	C 1241	N 302	O 383	S 5	0	0	0
2	B	245	Total 1935	C 1245	N 303	O 382	S 5	0	0	0
2	H	246	Total 1944	C 1249	N 304	O 386	S 5	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	1	MET	-	initiating methionine	UNP Q97Z84
B	1	MET	-	initiating methionine	UNP Q97Z84
H	1	MET	-	initiating methionine	UNP Q97Z84

- Molecule 3 is a protein called DNA polymerase sliding clamp A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	244	Total	C	N	O	S	0	0	0
			1934	1229	312	389	4			
3	C	244	Total	C	N	O	S	0	0	0
			1934	1229	312	389	4			
3	I	242	Total	C	N	O	S	0	0	0
			1917	1218	310	385	4			


- Molecule 4 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	6	Total	Br	0	0
			6	6		
4	E	8	Total	Br	0	0
			8	8		
4	F	10	Total	Br	0	0
			10	10		
4	A	5	Total	Br	0	0
			5	5		
4	B	5	Total	Br	0	0
			5	5		
4	C	7	Total	Br	0	0
			7	7		
4	G	2	Total	Br	0	0
			2	2		
4	H	5	Total	Br	0	0
			5	5		
4	I	6	Total	Br	0	0
			6	6		

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	1	Total	K	0	0
			1	1		

- Molecule 6 is 2,5,8,11,14,17,20,23-OCTAOXAPENTACOSAN-25-OL (three-letter code: 7PG) (formula: C₁₇H₃₆O₉).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	1	Total	C O	0	0
			26	17 9		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	111	Total	O	0	0
			111	111		
7	E	178	Total	O	0	0
			178	178		
7	F	112	Total	O	0	0
			112	112		
7	A	82	Total	O	0	0
			82	82		
7	B	168	Total	O	0	0
			168	168		
7	C	85	Total	O	0	0
			85	85		
7	G	67	Total	O	0	0
			67	67		
7	H	171	Total	O	0	0
			171	171		
7	I	72	Total	O	0	0
			72	72		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase sliding clamp B



- Molecule 1: DNA polymerase sliding clamp B





- Molecule 1: DNA polymerase sliding clamp B








- Molecule 2: DNA polymerase sliding clamp C





- Molecule 2: DNA polymerase sliding clamp C



- Molecule 2: DNA polymerase sliding clamp C




- Molecule 3: DNA polymerase sliding clamp A




- Molecule 3: DNA polymerase sliding clamp A





- Molecule 3: DNA polymerase sliding clamp A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	148.14Å 222.33Å 80.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.50 29.20 – 2.50	Depositor EDS
% Data completeness (in resolution range)	95.7 (30.00-2.50) 96.0 (29.20-2.50)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 2.51Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.214 , 0.251 0.210 , 0.246	Depositor DCC
R_{free} test set	1796 reflections (2.03%)	wwPDB-VP
Wilson B-factor (Å ²)	37.8	Xtrriage
Anisotropy	0.053	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18488	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, BR, 7PG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	1/1941 (0.1%)	0.56	0/2614
1	D	0.47	0/1955	0.58	0/2632
1	G	0.47	0/1951	0.56	0/2627
2	B	0.48	0/1970	0.62	1/2662 (0.0%)
2	E	0.47	0/1965	0.59	0/2652
2	H	0.47	0/1979	0.59	0/2673
3	C	0.46	0/1960	0.60	0/2641
3	F	0.44	0/1960	0.57	0/2641
3	I	0.51	1/1942 (0.1%)	0.56	0/2615
All	All	0.47	2/17623 (0.0%)	0.58	1/23757 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	105	GLU	CD-OE2	6.30	1.32	1.25
3	I	243	LYS	CE-NZ	5.87	1.63	1.49

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	73	LEU	CA-CB-CG	-5.15	103.46	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	171	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1914	0	1971	48	0
1	D	1928	0	1987	49	0
1	G	1924	0	1983	79	0
2	B	1935	0	1937	46	0
2	E	1931	0	1926	39	0
2	H	1944	0	1941	42	0
3	C	1934	0	1945	43	0
3	F	1934	0	1945	37	0
3	I	1917	0	1927	67	0
4	A	5	0	0	1	0
4	B	5	0	0	1	0
4	C	7	0	0	2	0
4	D	6	0	0	5	0
4	E	8	0	0	2	0
4	F	10	0	0	5	0
4	G	2	0	0	2	0
4	H	5	0	0	0	0
4	I	6	0	0	5	0
5	H	1	0	0	0	0
6	H	26	0	36	2	0
7	A	82	0	0	0	0
7	B	168	0	0	2	0
7	C	85	0	0	3	0
7	D	111	0	0	4	0
7	E	178	0	0	3	0
7	F	112	0	0	6	0
7	G	67	0	0	4	0
7	H	171	0	0	6	0
7	I	72	0	0	3	0
All	All	18488	0	17598	419	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 419 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:122:PRO:HB2	2:E:123:PRO:CD	1.79	1.13
2:E:122:PRO:HB2	2:E:123:PRO:HD3	1.30	1.10
1:G:89:ILE:HD11	1:G:100:ILE:HD11	1.34	1.05
2:B:17:ARG:HH11	2:B:17:ARG:HG2	1.23	1.03
1:G:56:VAL:HG11	1:G:238:GLY:HA3	1.40	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/249 (99%)	236 (96%)	8 (3%)	2 (1%)	19	35
1	D	247/249 (99%)	239 (97%)	7 (3%)	1 (0%)	34	54
1	G	247/249 (99%)	237 (96%)	10 (4%)	0	100	100
2	B	243/246 (99%)	237 (98%)	6 (2%)	0	100	100
2	E	240/246 (98%)	232 (97%)	6 (2%)	2 (1%)	19	35
2	H	244/246 (99%)	237 (97%)	7 (3%)	0	100	100
3	C	242/244 (99%)	227 (94%)	11 (4%)	4 (2%)	9	16
3	F	242/244 (99%)	231 (96%)	11 (4%)	0	100	100
3	I	238/244 (98%)	228 (96%)	10 (4%)	0	100	100
All	All	2189/2217 (99%)	2104 (96%)	76 (4%)	9 (0%)	34	54

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	126	LEU
3	C	179	ASP
2	E	122	PRO
1	A	128	PRO
3	C	124	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/220 (99%)	200 (92%)	18 (8%)	11	22
1	D	220/220 (100%)	201 (91%)	19 (9%)	10	20
1	G	219/220 (100%)	195 (89%)	24 (11%)	6	12
2	B	219/220 (100%)	197 (90%)	22 (10%)	7	15
2	E	218/220 (99%)	200 (92%)	18 (8%)	11	22
2	H	220/220 (100%)	199 (90%)	21 (10%)	8	17
3	C	220/220 (100%)	195 (89%)	25 (11%)	5	11
3	F	220/220 (100%)	206 (94%)	14 (6%)	17	33
3	I	218/220 (99%)	194 (89%)	24 (11%)	6	12
All	All	1972/1980 (100%)	1787 (91%)	185 (9%)	8	17

5 of 185 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	229	ASN
2	H	33	GLU
1	G	38	ARG
1	G	130	VAL
2	H	85	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	184	GLN
1	G	135	ASN
3	C	194	ASN
3	C	229	ASN
3	I	100	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 56 ligands modelled in this entry, 55 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	7PG	H	2001	5	25,25,25	0.72	0	24,24,24	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	7PG	H	2001	5	-	12/23/23/23	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	H	2001	7PG	C6-C5-O2-C4
6	H	2001	7PG	C15-C16-O8-C17
6	H	2001	7PG	O2-C5-C6-O3
6	H	2001	7PG	C13-C14-O7-C15
6	H	2001	7PG	C14-C13-O6-C12

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	H	2001	7PG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	248/249 (99%)	0.14	13 (5%) 27 29	24, 43, 66, 84	0
1	D	249/249 (100%)	0.13	7 (2%) 53 56	22, 38, 58, 73	0
1	G	249/249 (100%)	0.33	16 (6%) 19 20	24, 44, 67, 77	0
2	B	245/246 (99%)	-0.19	6 (2%) 59 62	18, 32, 47, 65	0
2	E	244/246 (99%)	0.08	8 (3%) 46 50	18, 30, 48, 67	0
2	H	246/246 (100%)	-0.07	11 (4%) 33 36	19, 31, 50, 78	0
3	C	244/244 (100%)	0.35	22 (9%) 9 9	27, 42, 65, 80	0
3	F	244/244 (100%)	0.37	19 (7%) 13 13	27, 40, 66, 88	0
3	I	242/244 (99%)	0.36	17 (7%) 16 16	29, 44, 64, 84	0
All	All	2211/2217 (99%)	0.17	119 (5%) 25 27	18, 38, 63, 88	0

The worst 5 of 119 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	82	LYS	8.2
2	H	125	VAL	7.0
3	I	244	VAL	6.7
3	C	123	GLU	6.3
2	H	246	ASP	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	K	H	4001	1/1	0.68	0.21	122,122,122,122	0
6	7PG	H	2001	26/26	0.82	0.21	43,57,65,66	0
4	BR	I	3032	1/1	0.84	0.10	63,63,63,63	1
4	BR	E	3010	1/1	0.86	0.17	50,50,50,50	1
4	BR	A	3043	1/1	0.89	0.10	52,52,52,52	1
4	BR	I	3048	1/1	0.89	0.16	60,60,60,60	1
4	BR	D	3037	1/1	0.90	0.18	52,52,52,52	1
4	BR	A	3046	1/1	0.91	0.14	64,64,64,64	1
4	BR	H	3020	1/1	0.91	0.07	49,49,49,49	1
4	BR	D	3028	1/1	0.91	0.14	58,58,58,58	1
4	BR	F	3041	1/1	0.92	0.10	50,50,50,50	1
4	BR	I	3019	1/1	0.92	0.14	56,56,56,56	1
4	BR	E	3050	1/1	0.92	0.11	60,60,60,60	1
4	BR	F	3021	1/1	0.92	0.11	63,63,63,63	1
4	BR	G	3045	1/1	0.92	0.15	52,52,52,52	1
4	BR	G	3052	1/1	0.92	0.12	48,48,48,48	1
4	BR	B	3036	1/1	0.93	0.13	52,52,52,52	1
4	BR	H	3004	1/1	0.93	0.07	50,50,50,50	1
4	BR	F	3006	1/1	0.94	0.11	42,42,42,42	1
4	BR	D	3039	1/1	0.94	0.14	66,66,66,66	1
4	BR	C	3054	1/1	0.94	0.15	60,60,60,60	1
4	BR	D	3016	1/1	0.94	0.09	51,51,51,51	1
4	BR	A	3033	1/1	0.94	0.25	52,52,52,52	1
4	BR	D	3027	1/1	0.94	0.08	53,53,53,53	1
4	BR	C	3018	1/1	0.95	0.11	57,57,57,57	1
4	BR	H	3051	1/1	0.95	0.09	56,56,56,56	1
4	BR	I	3003	1/1	0.95	0.12	50,50,50,50	1
4	BR	C	3035	1/1	0.95	0.11	50,50,50,50	1
4	BR	F	3023	1/1	0.95	0.10	37,37,37,37	1
4	BR	B	3031	1/1	0.95	0.12	63,63,63,63	1
4	BR	D	3015	1/1	0.95	0.09	40,40,40,40	1
4	BR	B	3038	1/1	0.95	0.12	40,40,40,40	1
4	BR	F	3024	1/1	0.96	0.06	40,40,40,40	1
4	BR	C	3042	1/1	0.96	0.06	52,52,52,52	1
4	BR	I	3011	1/1	0.96	0.07	53,53,53,53	1
4	BR	A	3047	1/1	0.96	0.13	47,47,47,47	1
4	BR	F	3012	1/1	0.96	0.07	43,43,43,43	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	BR	E	3017	1/1	0.96	0.07	56,56,56,56	1
4	BR	E	3049	1/1	0.96	0.15	56,56,56,56	1
4	BR	A	3044	1/1	0.96	0.25	56,56,56,56	1
4	BR	B	3026	1/1	0.97	0.15	65,65,65,65	1
4	BR	B	3053	1/1	0.97	0.08	51,51,51,51	1
4	BR	I	3007	1/1	0.97	0.07	44,44,44,44	1
4	BR	H	3025	1/1	0.98	0.06	49,49,49,49	1
4	BR	F	3030	1/1	0.98	0.06	53,53,53,53	1
4	BR	E	3040	1/1	0.98	0.06	65,65,65,65	1
4	BR	E	3022	1/1	0.98	0.10	42,42,42,42	1
4	BR	C	3008	1/1	0.98	0.05	40,40,40,40	1
4	BR	F	3002	1/1	0.99	0.04	40,40,40,40	1
4	BR	F	3005	1/1	0.99	0.05	44,44,44,44	1
4	BR	C	3009	1/1	0.99	0.05	53,53,53,53	1
4	BR	E	3013	1/1	0.99	0.09	46,46,46,46	1
4	BR	H	3014	1/1	0.99	0.04	39,39,39,39	1
4	BR	C	3029	1/1	0.99	0.09	50,50,50,50	1
4	BR	E	3001	1/1	0.99	0.08	36,36,36,36	1
4	BR	F	3034	1/1	0.99	0.08	51,51,51,51	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.