



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2023 – 10:58 AM EDT

PDB ID : 2P8I  
Title : Crystal structure of putative dioxygenase (YP\_555069.1) from Burkholderia xenovorans LB400 at 1.40 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-03-22  
Resolution : 1.40 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	117	 85% 15%
1	B	117	 89% 9% ..
1	C	117	 93% 6% .
1	D	117	 94% 6%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4786 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	117	970	623	174	170	3	0	5	0
1	B	117	972	620	177	173	2	0	4	0
1	C	116	958	615	169	171	3	0	5	0
1	D	117	952	607	174	169	2	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q13JM0
A	1	MSE	MET	modified residue	UNP Q13JM0
A	56	MSE	MET	modified residue	UNP Q13JM0
B	0	GLY	-	expression tag	UNP Q13JM0
B	1	MSE	MET	modified residue	UNP Q13JM0
B	56	MSE	MET	modified residue	UNP Q13JM0
C	0	GLY	-	expression tag	UNP Q13JM0
C	1	MSE	MET	modified residue	UNP Q13JM0
C	56	MSE	MET	modified residue	UNP Q13JM0
D	0	GLY	-	expression tag	UNP Q13JM0
D	1	MSE	MET	modified residue	UNP Q13JM0
D	56	MSE	MET	modified residue	UNP Q13JM0

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Cl 1	0	0
2	B	2	Total 2	Cl 2	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Cl 1 1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	D	1	Total O P 5 4 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0

- Molecule 5 is CITRIC ACID (three-letter code: CIT) (formula:  $C_6H_8O_7$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			13	6	7		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	204	Total	O	0	0
			204	204		
6	B	236	Total	O	0	0
			236	236		
6	C	204	Total	O	0	0
			204	204		
6	D	233	Total	O	0	0
			233	233		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.84Å 138.77Å 44.05Å 90.00° 98.43° 90.00°	Depositor
Resolution (Å)	30.00 – 1.40 30.00 – 1.40	Depositor EDS
% Data completeness (in resolution range)	84.3 (30.00-1.40) 84.3 (30.00-1.40)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.51 (at 1.40Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.144 , 0.173 0.152 , 0.180	Depositor DCC
$R_{free}$ test set	4342 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.2	Xtrriage
Anisotropy	0.189	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.032 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4786	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, EDO, CIT, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.79	1/1016 (0.1%)	0.85	2/1380 (0.1%)
1	B	0.73	2/1015 (0.2%)	1.26	6/1379 (0.4%)
1	C	0.73	0/1004	0.78	0/1365
1	D	0.66	0/986	0.77	1/1340 (0.1%)
All	All	0.73	3/4021 (0.1%)	0.94	9/5464 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	MSE	SE-CE	-8.54	1.45	1.95
1	B	49[A]	ARG	CZ-NH1	6.38	1.41	1.33
1	B	49[B]	ARG	CZ-NH1	6.38	1.41	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	49[A]	ARG	NE-CZ-NH2	-20.04	110.28	120.30
1	B	49[B]	ARG	NE-CZ-NH2	-20.04	110.28	120.30
1	B	49[A]	ARG	NE-CZ-NH1	15.69	128.14	120.30
1	B	49[B]	ARG	NE-CZ-NH1	15.69	128.14	120.30
1	B	30	ARG	NE-CZ-NH1	-6.69	116.96	120.30
1	A	95	ARG	NE-CZ-NH2	-5.70	117.45	120.30
1	B	30	ARG	NE-CZ-NH2	5.67	123.14	120.30
1	A	30	ARG	NE-CZ-NH1	5.43	123.01	120.30
1	D	98	ARG	NE-CZ-NH1	5.18	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	970	0	926	20	0
1	B	972	0	915	10	0
1	C	958	0	901	8	0
1	D	952	0	890	5	0
2	A	1	0	0	1	0
2	B	2	0	0	0	0
2	D	1	0	0	0	0
3	A	5	0	0	1	0
3	B	5	0	0	0	0
3	C	5	0	0	1	0
3	D	5	0	0	0	0
4	A	8	0	12	0	0
4	C	8	0	12	0	0
4	D	4	0	6	0	0
5	C	13	0	5	0	0
6	A	204	0	0	5	0
6	B	236	0	0	2	0
6	C	204	0	0	4	0
6	D	233	0	0	1	0
All	All	4786	0	3667	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43[B]:LEU:CD2	1:A:61[B]:LEU:HD13	2.09	0.81
1:A:112:LEU:HD11	6:A:248:HOH:O	1.84	0.77
1:A:43[B]:LEU:CD2	1:A:61[B]:LEU:CD1	2.71	0.69
1:B:40:LYS:O	1:B:67[A]:GLN:NE2	2.27	0.68
1:C:109[A]:GLU:CD	6:C:633:HOH:O	2.36	0.63
2:A:117:CL:CL	1:B:79:HIS:NE2	2.69	0.61
1:C:3:PHE:CE2	1:D:32:GLN:HG2	2.35	0.61
1:A:112:LEU:CD1	6:A:248:HOH:O	2.46	0.60

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:HIS:HB3	1:B:37:TRP:CD1	2.38	0.59
1:A:43[B]:LEU:HD21	1:A:61[B]:LEU:HD13	1.82	0.59
1:B:30:ARG:HD3	6:B:424:HOH:O	2.03	0.58
1:A:112:LEU:CG	6:A:248:HOH:O	2.55	0.54
1:A:43[B]:LEU:HD23	1:A:61[B]:LEU:CD1	2.36	0.54
1:A:85:PHE:HB2	1:A:103:TRP:CZ3	2.43	0.53
1:D:64:THR:H	1:D:67[A]:GLN:HE21	1.55	0.53
1:A:43[A]:LEU:HD12	1:A:61[A]:LEU:HD23	1.92	0.51
1:C:85:PHE:HB2	1:C:103:TRP:CZ3	2.47	0.49
1:A:43[B]:LEU:HD23	1:A:61[B]:LEU:HD12	1.95	0.49
1:A:112:LEU:HG	6:A:248:HOH:O	2.11	0.48
1:A:43[B]:LEU:HD21	1:A:61[B]:LEU:CD1	2.42	0.48
1:A:42:GLN:HG3	1:C:27:TRP:CD1	2.48	0.47
1:C:43:LEU:HD12	6:C:471:HOH:O	2.15	0.47
1:A:47:HIS:NE2	3:A:118:PO4:O4	2.47	0.45
1:A:43[A]:LEU:CD1	1:A:61[A]:LEU:HD23	2.47	0.45
1:B:85:PHE:HB2	1:B:103:TRP:CZ3	2.52	0.45
1:C:43:LEU:CD1	6:C:471:HOH:O	2.65	0.45
1:B:16:VAL:O	1:B:58:SER:HA	2.17	0.44
1:D:64:THR:H	1:D:67[A]:GLN:NE2	2.15	0.44
1:A:16:VAL:O	1:A:58:SER:HA	2.18	0.43
1:B:36:HIS:HB2	6:B:329:HOH:O	2.18	0.43
1:B:49[A]:ARG:CZ	1:B:51:VAL:HG12	2.49	0.42
1:A:56[B]:MSE:SE	1:A:108:HIS:CE1	3.23	0.42
1:D:103:TRP:CD2	1:D:107:SER:HA	2.55	0.42
1:C:47:HIS:NE2	3:C:117:PO4:O4	2.44	0.41
1:A:32[B]:GLN:HG3	1:B:3:PHE:CZ	2.55	0.41
1:B:40:LYS:O	1:B:67[A]:GLN:CD	2.60	0.41
1:D:49:ARG:CZ	6:D:810:HOH:O	2.69	0.41
1:A:49:ARG:NE	6:A:307:HOH:O	2.54	0.40
1:C:109[A]:GLU:CG	6:C:633:HOH:O	2.69	0.40
1:A:103:TRP:CD2	1:A:107:SER:HA	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/117 (103%)	118 (98%)	2 (2%)	0	100	100
1	B	119/117 (102%)	117 (98%)	2 (2%)	0	100	100
1	C	119/117 (102%)	117 (98%)	2 (2%)	0	100	100
1	D	116/117 (99%)	114 (98%)	2 (2%)	0	100	100
All	All	474/468 (101%)	466 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	100/93 (108%)	100 (100%)	0	100	100
1	B	99/93 (106%)	97 (98%)	2 (2%)	55	23
1	C	98/93 (105%)	98 (100%)	0	100	100
1	D	96/93 (103%)	96 (100%)	0	100	100
All	All	393/372 (106%)	391 (100%)	2 (0%)	92	74

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	49[A]	ARG
1	B	49[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	36	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	32	GLN
1	D	116	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	C	117	-	4,4,4	1.12	0	6,6,6	2.85	3 (50%)
3	PO4	D	118	-	4,4,4	1.28	1 (25%)	6,6,6	1.16	0
5	CIT	C	118	-	12,12,12	1.20	0	17,17,17	1.80	5 (29%)
4	EDO	C	119	-	3,3,3	0.26	0	2,2,2	1.03	0
3	PO4	B	119	-	4,4,4	2.01	1 (25%)	6,6,6	1.20	1 (16%)
4	EDO	C	120	-	3,3,3	0.56	0	2,2,2	0.12	0
4	EDO	A	119	-	3,3,3	0.58	0	2,2,2	0.44	0
4	EDO	A	120	-	3,3,3	0.25	0	2,2,2	0.71	0
4	EDO	D	119	-	3,3,3	0.39	0	2,2,2	0.50	0
3	PO4	A	118	-	4,4,4	1.18	0	6,6,6	1.24	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CIT	C	118	-	-	2/16/16/16	-
4	EDO	C	119	-	-	0/1/1/1	-
4	EDO	C	120	-	-	0/1/1/1	-
4	EDO	A	119	-	-	0/1/1/1	-
4	EDO	A	120	-	-	1/1/1/1	-
4	EDO	D	119	-	-	0/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	119	PO4	P-O1	3.41	1.58	1.50
3	D	118	PO4	P-O3	-2.08	1.48	1.54

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	117	PO4	O4-P-O2	-4.64	93.09	107.97
5	C	118	CIT	O6-C6-C3	3.84	119.71	113.05
3	C	117	PO4	O4-P-O1	-3.42	98.39	110.89
3	C	117	PO4	O2-P-O1	3.35	123.14	110.89
5	C	118	CIT	O5-C6-C3	-3.16	117.78	122.25
5	C	118	CIT	O2-C1-C2	2.46	122.25	114.35
3	A	118	PO4	O4-P-O1	-2.32	102.41	110.89
5	C	118	CIT	C4-C3-C6	-2.16	105.46	110.11
3	B	119	PO4	O4-P-O2	-2.04	101.43	107.97
5	C	118	CIT	O4-C5-C4	2.02	120.85	114.35

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	120	EDO	O1-C1-C2-O2
5	C	118	CIT	O1-C1-C2-C3
5	C	118	CIT	O2-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	117	PO4	1	0
3	A	118	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	115/117 (98%)	-0.36	1 (0%) 84 82	9, 12, 21, 37	0
1	B	115/117 (98%)	-0.29	3 (2%) 56 55	9, 12, 25, 33	0
1	C	114/117 (97%)	-0.46	1 (0%) 84 82	8, 11, 20, 35	0
1	D	115/117 (98%)	-0.30	3 (2%) 56 55	9, 12, 26, 34	0
All	All	459/468 (98%)	-0.35	8 (1%) 70 69	8, 12, 25, 37	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	0	GLY	7.2
1	D	0	GLY	4.6
1	A	0	GLY	3.6
1	B	0	GLY	3.6
1	B	38	SER	3.4
1	D	36	HIS	2.6
1	D	38	SER	2.3
1	B	40	LYS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PO4	A	118	5/5	0.92	0.17	20,28,32,44	0
3	PO4	B	119	5/5	0.93	0.16	15,23,29,29	0
4	EDO	A	120	4/4	0.94	0.08	18,19,24,28	0
4	EDO	C	119	4/4	0.94	0.09	18,19,24,24	0
5	CIT	C	118	13/13	0.95	0.15	16,22,39,49	0
3	PO4	C	117	5/5	0.96	0.12	9,22,34,41	0
3	PO4	D	118	5/5	0.96	0.17	15,23,25,27	0
4	EDO	C	120	4/4	0.96	0.07	12,14,15,16	0
4	EDO	A	119	4/4	0.96	0.08	14,16,17,17	0
4	EDO	D	119	4/4	0.97	0.06	17,18,19,21	0
2	CL	A	117	1/1	0.97	0.12	24,24,24,24	0
2	CL	B	118	1/1	0.99	0.10	23,23,23,23	0
2	CL	D	117	1/1	0.99	0.08	19,19,19,19	0
2	CL	B	117	1/1	1.00	0.05	18,18,18,18	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.