



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 21, 2023 – 02:17 PM EDT

PDB ID : 2QE0
Title : Thioacylenzyme Intermediate of GAPN from *S. Mutans*, New Data Integration and Refinement.
Authors : Corbier, C.; Didierjean, C.; Bricogne, G.; Branlant, G.; D'Ambrosio, K.; Vonrhein, C.
Deposited on : 2007-06-22
Resolution : 2.19 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	 79% 18% .
1	B	475	 79% 19% .
1	C	475	 83% 14% .
1	D	475	 77% 21% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	G3H	A	800	-	X	-	-
2	G3H	B	800	-	X	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 16008 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NADP-dependent glyceraldehyde-3-phosphate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	474	3598	2287	601	698	12	0	1	0
1	B	474	3598	2287	601	698	12	0	1	0
1	C	474	3598	2287	601	698	12	0	1	0
1	D	474	3598	2287	601	698	12	0	1	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	58	ALA	SER	variant	UNP Q59931
A	85	ILE	VAL	variant	UNP Q59931
A	250	ALA	GLU	engineered mutation	UNP Q59931
A	347	THR	ALA	variant	UNP Q59931
B	58	ALA	SER	variant	UNP Q59931
B	85	ILE	VAL	variant	UNP Q59931
B	250	ALA	GLU	engineered mutation	UNP Q59931
B	347	THR	ALA	variant	UNP Q59931
C	58	ALA	SER	variant	UNP Q59931
C	85	ILE	VAL	variant	UNP Q59931
C	250	ALA	GLU	engineered mutation	UNP Q59931
C	347	THR	ALA	variant	UNP Q59931
D	58	ALA	SER	variant	UNP Q59931
D	85	ILE	VAL	variant	UNP Q59931
D	250	ALA	GLU	engineered mutation	UNP Q59931
D	347	THR	ALA	variant	UNP Q59931

- Molecule 2 is GLYCERALDEHYDE-3-PHOSPHATE (three-letter code: G3H) (formula: C₃H₇O₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	A	1	10	3	6	1	0	0
2	B	1	10	3	6	1	0	0
2	C	1	10	3	6	1	0	0
2	D	1	10	3	6	1	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	48	21	7	17	3	0	0
3	B	1	48	21	7	17	3	0	0
3	C	1	48	21	7	17	3	0	0
3	D	1	48	21	7	17	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	368	368	368	0	0
4	B	300	300	300	0	0
4	C	403	403	403	0	0
4	D	313	313	313	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NADP-dependent glyceraldehyde-3-phosphate dehydrogenase




- Molecule 1: NADP-dependent glyceraldehyde-3-phosphate dehydrogenase





- Molecule 1: NADP-dependent glyceraldehyde-3-phosphate dehydrogenase





● Molecule 1: NADP-dependent glyceraldehyde-3-phosphate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	142.47Å 155.21Å 113.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.78 – 2.19 19.78 – 2.22	Depositor EDS
% Data completeness (in resolution range)	96.5 (19.78-2.19) 99.8 (19.78-2.22)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 2.21Å)	Xtrriage
Refinement program	BUSTER-TNT 2.1.1	Depositor
R, R_{free}	0.184 , 0.230 0.187 , 0.232	Depositor DCC
R_{free} test set	12452 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtrriage
Anisotropy	0.380	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 62.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16008	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.04 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.9128e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G3H, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/3656	0.73	2/4945 (0.0%)
1	B	0.55	0/3656	0.72	5/4945 (0.1%)
1	C	0.65	0/3656	0.78	3/4945 (0.1%)
1	D	0.61	0/3656	0.77	9/4945 (0.2%)
All	All	0.60	0/14624	0.75	19/19780 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	103	ARG	NE-CZ-NH1	8.77	124.69	120.30
1	D	103	ARG	NE-CZ-NH2	-7.93	116.34	120.30
1	C	283	ARG	NE-CZ-NH2	6.66	123.63	120.30
1	D	283	ARG	NE-CZ-NH2	6.65	123.62	120.30
1	C	456	GLY	N-CA-C	-6.56	96.70	113.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	438	GLY	Mainchain
1	D	438	GLY	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3598	0	3651	78	0
1	B	3598	0	3651	71	0
1	C	3598	0	3651	64	1
1	D	3598	0	3651	78	0
2	A	10	0	4	2	0
2	B	10	0	4	4	0
2	C	10	0	4	1	0
2	D	10	0	4	1	0
3	A	48	0	25	8	0
3	B	48	0	25	7	0
3	C	48	0	24	7	0
3	D	48	0	25	9	0
4	A	368	0	0	14	0
4	B	300	0	0	4	0
4	C	403	0	0	10	0
4	D	313	0	0	9	0
All	All	16008	0	14719	272	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 272 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:230:GLY:HA2	3:D:4476:NAP:H51N	1.31	1.07
1:D:439:THR:HG23	1:D:441:ASN:OD1	1.62	0.98
1:C:439:THR:HG23	1:C:441:ASN:OD1	1.66	0.94
1:A:439:THR:HG23	1:A:441:ASN:OD1	1.68	0.92
1:B:439:THR:HG23	1:B:441:ASN:OD1	1.70	0.91

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:3:LYS:NZ	1:C:195:GLU:OE1[2_355]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	473/475 (100%)	463 (98%)	9 (2%)	1 (0%)	47	55
1	B	473/475 (100%)	456 (96%)	15 (3%)	2 (0%)	34	37
1	C	473/475 (100%)	461 (98%)	12 (2%)	0	100	100
1	D	473/475 (100%)	454 (96%)	17 (4%)	2 (0%)	34	37
All	All	1892/1900 (100%)	1834 (97%)	53 (3%)	5 (0%)	41	46

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	438	GLY
1	D	438	GLY
1	B	439	THR
1	D	439	THR
1	A	378	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	378/378 (100%)	350 (93%)	28 (7%)	13	14
1	B	378/378 (100%)	356 (94%)	22 (6%)	20	23
1	C	378/378 (100%)	354 (94%)	24 (6%)	18	20
1	D	378/378 (100%)	354 (94%)	24 (6%)	18	20
All	All	1512/1512 (100%)	1414 (94%)	98 (6%)	17	19

5 of 98 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	270	LYS
1	C	448	LYS
1	C	329	LYS
1	C	379	PHE
1	D	17	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	430	HIS
1	D	455	GLN
1	C	109	ASN
1	C	227	ASN
1	C	430	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAP	B	2476	-	45,52,52	1.70	11 (24%)	56,80,80	1.95	13 (23%)
3	NAP	C	3476	-	45,52,52	2.24	14 (31%)	56,80,80	2.50	26 (46%)
2	G3H	D	800	-	8,9,9	2.88	4 (50%)	10,12,12	4.08	6 (60%)
2	G3H	C	800	-	8,9,9	3.14	4 (50%)	10,12,12	4.32	5 (50%)
2	G3H	A	800	-	8,9,9	3.00	5 (62%)	10,12,12	3.73	6 (60%)
3	NAP	A	1476	-	45,52,52	1.77	9 (20%)	56,80,80	1.83	12 (21%)
3	NAP	D	4476	-	45,52,52	1.64	10 (22%)	56,80,80	2.01	12 (21%)
2	G3H	B	800	-	8,9,9	2.98	4 (50%)	10,12,12	4.28	7 (70%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	B	2476	-	-	9/31/67/67	0/5/5/5
3	NAP	C	3476	-	-	16/31/67/67	0/5/5/5
2	G3H	D	800	-	-	1/7/8/8	-
2	G3H	C	800	-	-	1/7/8/8	-
2	G3H	A	800	-	-	1/7/8/8	-
3	NAP	A	1476	-	-	6/31/67/67	0/5/5/5
3	NAP	D	4476	-	-	7/31/67/67	0/5/5/5
2	G3H	B	800	-	-	1/7/8/8	-

The worst 5 of 61 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	3476	NAP	C2N-N1N	8.07	1.44	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	800	G3H	P-O1P	5.57	1.78	1.60
3	B	2476	NAP	O4B-C1B	5.43	1.48	1.41
3	D	4476	NAP	O4D-C1D	5.31	1.48	1.41
2	C	800	G3H	P-O1P	5.15	1.76	1.60

The worst 5 of 87 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	800	G3H	O1P-P-O2P	-9.18	80.71	106.47
3	D	4476	NAP	O3D-C3D-C2D	-8.28	85.05	111.82
2	C	800	G3H	O4P-P-O1P	8.21	128.59	106.73
3	C	3476	NAP	C3N-C7N-N7N	8.08	127.44	117.75
2	D	800	G3H	O1P-P-O2P	-7.94	84.20	106.47

There are no chirality outliers.

5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	800	G3H	C3-O1P-P-O2P
3	A	1476	NAP	C5D-O5D-PN-O3
3	A	1476	NAP	C5D-O5D-PN-O1N
3	A	1476	NAP	C5D-O5D-PN-O2N
3	A	1476	NAP	C2D-C1D-N1N-C2N


There are no ring outliers.


8 monomers are involved in 39 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	2476	NAP	7	0
3	C	3476	NAP	7	0
2	D	800	G3H	1	0
2	C	800	G3H	1	0
2	A	800	G3H	2	0
3	A	1476	NAP	8	0
3	D	4476	NAP	9	0
2	B	800	G3H	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	474/475 (99%)	-0.23	5 (1%) 80 79	18, 29, 49, 79	0
1	B	474/475 (99%)	-0.10	10 (2%) 63 61	21, 33, 53, 87	0
1	C	474/475 (99%)	-0.38	3 (0%) 89 88	13, 23, 43, 63	0
1	D	474/475 (99%)	-0.10	11 (2%) 60 58	19, 31, 51, 73	0
All	All	1896/1900 (99%)	-0.20	29 (1%) 73 72	13, 29, 50, 87	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	THR	6.5
1	A	2	THR	5.5
1	B	29	ALA	3.6
1	A	17	GLU	3.4
1	A	258	ILE	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAP	B	2476	48/48	0.85	0.20	33,39,41,43	48
2	G3H	D	800	10/10	0.86	0.23	30,36,38,40	10
2	G3H	A	800	10/10	0.86	0.21	29,34,35,36	10
3	NAP	D	4476	48/48	0.87	0.20	34,41,45,47	48
2	G3H	B	800	10/10	0.90	0.22	31,38,39,41	10
3	NAP	A	1476	48/48	0.90	0.19	33,38,41,43	48
3	NAP	C	3476	48/48	0.92	0.17	23,30,36,38	48
2	G3H	C	800	10/10	0.94	0.19	24,29,29,32	10

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around NAP D 4476:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP A 1476:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.