



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2023 – 04:16 AM EDT

PDB ID : 2QIN
Title : Stenotrophomonas maltophilia L1 Metallo-beta-Lactamase Asp-120 Cys mutant
Authors : Spencer, J.
Deposited on : 2007-07-05
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	 3% 94% 2% 1%
1	B	269	 2% 96% 1% 1%
1	C	269	 9% 94% 5% 2%
1	D	269	 0% 97% 2% 1%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9238 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metallo-beta-lactamase L1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	263	1977	1245	356	367	9	0	0	0
1	B	264	1986	1249	358	370	9	0	1	0
1	C	265	1990	1252	359	370	9	0	0	0
1	D	265	1984	1249	356	370	9	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	120	CYS	ASP	engineered mutation	UNP P52700
B	120	CYS	ASP	engineered mutation	UNP P52700
C	120	CYS	ASP	engineered mutation	UNP P52700
D	120	CYS	ASP	engineered mutation	UNP P52700

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Zn 2	0	0
2	B	3	Total 3	Zn 3	0	0
2	C	2	Total 2	Zn 2	0	0
2	D	2	Total 2	Zn 2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	311	Total O 311 311	0	0
4	B	352	Total O 352 352	0	0
4	C	250	Total O 250 250	0	0
4	D	377	Total O 377 377	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Metallo-beta-lactamase L1



- Molecule 1: Metallo-beta-lactamase L1



- Molecule 1: Metallo-beta-lactamase L1



- Molecule 1: Metallo-beta-lactamase L1



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.14Å 112.84Å 78.35Å 90.00° 113.35° 90.00°	Depositor
Resolution (Å)	26.84 – 1.76 26.84 – 1.76	Depositor EDS
% Data completeness (in resolution range)	94.3 (26.84-1.76) 94.4 (26.84-1.76)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.01 (at 1.76Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.173 , 0.212 0.173 , 0.211	Depositor DCC
R_{free} test set	4969 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	17.0	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9238	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/2027	0.50	0/2769
1	B	0.34	0/2039	0.52	0/2783
1	C	0.32	0/2040	0.48	0/2786
1	D	0.37	0/2034	0.52	0/2779
All	All	0.34	0/8140	0.51	0/11117

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1977	0	1938	4	0
1	B	1986	0	1949	6	0
1	C	1990	0	1952	6	0
1	D	1984	0	1941	2	0
2	A	2	0	0	0	0
2	B	3	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	1	0	0	0	0
4	A	311	0	0	0	0
4	B	352	0	0	1	0
4	C	250	0	0	1	0
4	D	377	0	0	2	0
All	All	9238	0	7780	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:30:ARG:HH11	1:B:30:ARG:HG2	1.24	1.00
1:A:100:LYS:HA	1:A:100:LYS:HE2	1.75	0.69
1:B:30:ARG:HH11	1:B:30:ARG:CG	2.03	0.66
1:C:49:ALA:HA	1:C:209:ARG:HD3	1.78	0.64
1:B:30:ARG:HG2	1:B:30:ARG:NH1	2.05	0.64
1:C:75:THR:HG22	1:C:185:ILE:HD12	1.82	0.62
1:C:24:VAL:N	4:C:2166:HOH:O	2.36	0.58
1:B:245:ARG:HD2	4:B:4119:HOH:O	2.07	0.55
1:A:75:THR:HG22	1:A:185:ILE:HD12	1.91	0.52
1:B:162:ILE:HG13	1:B:162:ILE:O	2.11	0.51
1:C:310:ALA:HA	1:C:311:GLY:HA2	1.60	0.45
1:D:209:ARG:NH2	4:D:4273:HOH:O	2.47	0.43
1:B:30:ARG:CG	1:B:30:ARG:NH1	2.71	0.43
1:C:212:LYS:HA	1:C:212:LYS:HE2	2.01	0.42
1:A:78:ASP:HB2	1:A:183:GLY:CA	2.50	0.41
1:D:229:GLN:HG3	4:D:4334:HOH:O	2.19	0.41
1:C:139:ASN:HB3	1:C:191:PHE:CZ	2.56	0.41
1:A:41:PRO:HB3	1:A:94:HIS:CE1	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/269 (97%)	251 (96%)	10 (4%)	0	100	100
1	B	263/269 (98%)	254 (97%)	9 (3%)	0	100	100
1	C	263/269 (98%)	252 (96%)	11 (4%)	0	100	100
1	D	263/269 (98%)	254 (97%)	9 (3%)	0	100	100
All	All	1050/1076 (98%)	1011 (96%)	39 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/202 (98%)	194 (98%)	5 (2%)	47	25
1	B	200/202 (99%)	196 (98%)	4 (2%)	55	34
1	C	200/202 (99%)	197 (98%)	3 (2%)	65	49
1	D	199/202 (98%)	197 (99%)	2 (1%)	76	63
All	All	798/808 (99%)	784 (98%)	14 (2%)	59	40

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	LYS
1	A	120	CYS
1	A	236	TYR
1	A	245	ARG
1	A	309	THR
1	B	30	ARG
1	B	120	CYS
1	B	212	LYS
1	B	236	TYR
1	C	78	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	120	CYS
1	C	236	TYR
1	D	67	ASP
1	D	236	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	HIS
1	B	304	GLN
1	D	238	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/269 (97%)	0.18	7 (2%) 54 60	10, 18, 28, 41	0
1	B	264/269 (98%)	-0.10	6 (2%) 60 67	9, 14, 22, 36	1 (0%)
1	C	265/269 (98%)	0.58	24 (9%) 9 11	12, 23, 33, 40	0
1	D	265/269 (98%)	-0.09	4 (1%) 73 80	8, 12, 22, 31	0
All	All	1057/1076 (98%)	0.14	41 (3%) 39 45	8, 16, 29, 41	1 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	24	VAL	6.2
1	C	311	GLY	5.3
1	C	211	GLY	5.3
1	C	210	ASN	4.7
1	A	309	THR	4.6
1	C	24	VAL	4.4
1	C	310	ALA	4.2
1	C	209	ARG	3.8
1	C	184	GLY	3.8
1	A	24	VAL	3.8
1	C	212	LYS	3.7
1	C	272	ALA	3.5
1	B	310	ALA	3.5
1	C	273	ALA	3.3
1	C	76	PRO	3.1
1	D	25	PRO	3.1
1	C	268	ASN	3.0
1	B	311	GLY	3.0
1	C	107	ARG	3.0
1	C	286	LYS	2.9
1	A	306	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	285	ALA	2.9
1	C	30	ARG	2.7
1	B	276	ARG	2.6
1	C	299	GLN	2.5
1	C	276	ARG	2.5
1	A	211	GLY	2.4
1	C	309	THR	2.4
1	C	213	PRO	2.3
1	A	212	LYS	2.2
1	B	256	CYS	2.2
1	C	51	HIS	2.2
1	A	35	ASP	2.2
1	D	241	GLU	2.1
1	A	245	ARG	2.1
1	C	183	GLY	2.1
1	C	245	ARG	2.1
1	D	210	ASN	2.1
1	B	162	ILE	2.1
1	B	307	LYS	2.0
1	C	50	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	B	3001	1/1	0.73	0.25	10,10,10,10	1
2	ZN	C	2001	1/1	0.98	0.04	24,24,24,24	0
3	MG	B	4002	1/1	0.98	0.10	17,17,17,17	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	D	4001	1/1	0.98	0.09	23,23,23,23	0
2	ZN	A	2002	1/1	0.99	0.05	17,17,17,17	0
2	ZN	C	2002	1/1	0.99	0.07	20,20,20,20	0
2	ZN	D	2001	1/1	0.99	0.04	18,18,18,18	0
2	ZN	B	2001	1/1	0.99	0.04	16,16,16,16	0
2	ZN	A	2001	1/1	0.99	0.04	21,21,21,21	0
2	ZN	B	2002	1/1	1.00	0.04	13,13,13,13	0
2	ZN	D	2002	1/1	1.00	0.04	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.