



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2023 – 11:38 AM EDT

PDB ID : 2UBP
Title : STRUCTURE OF NATIVE UREASE FROM BACILLUS PASTEURII
Authors : Benini, S.; Rypniewski, W.R.; Wilson, K.S.; Ciurli, S.; Mangani, S.
Deposited on : 1998-11-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	101	 % 86% 12% •
2	B	122	 88% 12%
3	C	570	 2% 87% 10% ••

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6943 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (UREASE GAMMA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	101	781	494	133	148	6	8	0	0

- Molecule 2 is a protein called PROTEIN (UREASE BETA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	122	951	589	171	190	1	36	0	0

- Molecule 3 is a protein called PROTEIN (UREASE ALPHA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	570	4323	2714	743	843	23	44	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	19	GLU	ARG	variant	UNP P41020
C	28	TRP	-	insertion	UNP P41020
C	29	ILE	GLY	variant	UNP P41020
C	36	THR	TYR	variant	UNP P41020
C	37	THR	TYR	variant	UNP P41020
C	38	TYR	LEU	variant	UNP P41020
C	220	KCX	LYS	modified residue	UNP P41020
C	263	LEU	VAL	variant	UNP P41020
C	420	ILE	MET	conflict	UNP P41020

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total O S 5 4 1	0	0

- Molecule 5 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	2	Total Ni 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	124	Total O 124 124	0	0
6	B	179	Total O 179 179	0	0
6	C	578	Total O 578 578	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: PROTEIN (UREASE GAMMA SUBUNIT)



- Molecule 2: PROTEIN (UREASE BETA SUBUNIT)



- Molecule 3: PROTEIN (UREASE ALPHA SUBUNIT)



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.36Å 131.36Å 189.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.00 19.88 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.7 (20.00-2.00) 96.8 (19.88-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	7.60	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.01Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.160 , 0.200 0.163 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtrriage
Anisotropy	0.312	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6943	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, SO4, ACE, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.16	2/790 (0.3%)	1.52	6/1065 (0.6%)
2	B	1.02	6/963 (0.6%)	1.33	7/1296 (0.5%)
3	C	0.59	0/4392	1.19	22/5955 (0.4%)
All	All	0.76	8/6145 (0.1%)	1.26	35/8316 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	1	1
3	C	0	5
All	All	1	7

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	0	ACE	C-N	24.97	1.91	1.34
2	B	126	GLU	CA-CB	15.89	1.89	1.53
1	A	99	ILE	C-N	-13.81	1.02	1.34
2	B	5	ASN	N-CA	-11.16	1.24	1.46
2	B	6	TYR	N-CA	7.94	1.62	1.46
2	B	5	ASN	CA-CB	-7.83	1.32	1.53
2	B	5	ASN	CA-C	6.26	1.69	1.52
2	B	5	ASN	C-N	6.10	1.48	1.34

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	0	ACE	O-C-N	-25.11	82.53	122.70
2	B	5	ASN	N-CA-CB	20.83	148.09	110.60
1	A	99	ILE	O-C-N	-15.02	98.66	122.70
3	C	339	ARG	NE-CZ-NH2	-11.96	114.32	120.30
3	C	5	ARG	NE-CZ-NH2	-11.83	114.38	120.30
3	C	448	ASP	CB-CG-OD2	10.07	127.36	118.30
1	A	99	ILE	CA-C-N	9.60	138.33	117.20
3	C	305	ARG	NE-CZ-NH1	8.80	124.70	120.30
2	B	66	ARG	CD-NE-CZ	8.17	135.04	123.60
3	C	388	ARG	NE-CZ-NH2	-8.15	116.22	120.30
2	B	126	GLU	CB-CA-C	-8.01	94.39	110.40
2	B	126	GLU	CA-C-O	7.91	136.71	120.10
3	C	318	MET	CA-CB-CG	7.90	126.73	113.30
3	C	566	ARG	NE-CZ-NH2	-7.33	116.64	120.30
3	C	388	ARG	NE-CZ-NH1	7.18	123.89	120.30
3	C	536	ASP	CB-CG-OD2	-6.99	112.01	118.30
2	B	13	ARG	NE-CZ-NH2	6.94	123.77	120.30
1	A	22	ARG	NE-CZ-NH2	6.94	123.77	120.30
3	C	5	ARG	NH1-CZ-NH2	6.61	126.67	119.40
3	C	234	ARG	NE-CZ-NH2	-6.42	117.09	120.30
3	C	264	ARG	NE-CZ-NH2	-6.39	117.10	120.30
3	C	403	LEU	CA-CB-CG	-6.39	100.61	115.30
3	C	320	MET	CG-SD-CE	6.38	110.41	100.20
3	C	551	GLU	CA-CB-CG	6.06	126.72	113.40
2	B	31	ARG	NE-CZ-NH2	5.88	123.24	120.30
2	B	111	GLU	CA-CB-CG	5.78	126.11	113.40
1	A	100	SER	CA-C-O	-5.70	108.12	120.10
3	C	62	ARG	NE-CZ-NH2	-5.62	117.49	120.30
3	C	1	MET	CG-SD-CE	5.57	109.12	100.20
1	A	0	ACE	C-N-CA	5.43	135.27	121.70
3	C	86	ASP	CB-CG-OD1	5.41	123.17	118.30
3	C	21	ARG	NE-CZ-NH1	-5.22	117.69	120.30
3	C	4	ASN	CA-CB-CG	5.17	124.78	113.40
3	C	317	ASP	CB-CG-OD1	-5.06	113.75	118.30
3	C	302	ASN	N-CA-C	5.04	124.60	111.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	5	ASN	CA

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	99	ILE	Mainchain
2	B	101	ASP	Mainchain
3	C	15	THR	Mainchain
3	C	191	PRO	Mainchain
3	C	321	VAL	Mainchain
3	C	503	GLY	Mainchain
3	C	522	ASN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	781	0	805	17	0
2	B	951	0	937	5	0
3	C	4323	0	4290	38	1
4	C	5	0	0	0	0
5	C	2	0	0	0	0
6	A	124	0	0	2	0
6	B	179	0	0	5	0
6	C	578	0	0	12	5
All	All	6943	0	6032	59	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:0:ACE:C	1:A:1:MET:H2	1.27	1.38
1:A:0:ACE:CH3	1:A:0:ACE:O	1.66	1.32
1:A:0:ACE:O	1:A:0:ACE:H3	1.23	1.14
1:A:79:ASP:HB3	6:A:155:HOH:O	1.63	0.96
1:A:0:ACE:C	1:A:0:ACE:H1	1.28	0.89
1:A:0:ACE:CH3	1:A:1:MET:H2	1.81	0.88
1:A:0:ACE:C	1:A:0:ACE:H2	1.28	0.88
1:A:0:ACE:C	1:A:0:ACE:H3	1.28	0.86
1:A:0:ACE:CH3	1:A:1:MET:N	2.40	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:507:LYS:HD3	6:C:1364:HOH:O	1.77	0.83
3:C:367:MET:SD	6:C:1273:HOH:O	2.41	0.77
3:C:507:LYS:CD	6:C:1364:HOH:O	2.33	0.74
3:C:320:MET:HA	3:C:325:LEU:HD12	1.70	0.74
2:B:116:ARG:HA	6:B:300:HOH:O	1.89	0.73
3:C:322:CYS:SG	6:C:1154:HOH:O	2.39	0.72
2:B:53:LYS:HD2	6:B:228:HOH:O	1.89	0.71
6:B:276:HOH:O	3:C:1:MET:HG3	1.92	0.70
1:A:0:ACE:C	1:A:0:ACE:CH3	0.57	0.66
1:A:0:ACE:C	1:A:1:MET:H1	1.94	0.65
1:A:12:GLN:HG3	6:A:194:HOH:O	2.00	0.61
3:C:317:ASP:O	3:C:321:VAL:HG13	2.03	0.58
3:C:513:ARG:NH1	6:C:1293:HOH:O	2.16	0.58
6:B:240:HOH:O	3:C:4:ASN:HB2	2.04	0.56
1:A:0:ACE:O	1:A:1:MET:N	2.26	0.56
3:C:513:ARG:HD2	6:C:1293:HOH:O	2.05	0.56
3:C:314:GLU:OE2	6:C:1088:HOH:O	2.18	0.53
3:C:70:LEU:HD11	3:C:86:ASP:HB3	1.88	0.53
3:C:513:ARG:NH1	6:C:1433:HOH:O	2.41	0.52
3:C:470:THR:N	3:C:471:PRO:CD	2.71	0.52
3:C:249:HIS:CE1	3:C:281:GLY:HA3	2.46	0.51
3:C:206:ALA:HB3	3:C:207:PRO:HD3	1.93	0.51
2:B:95:GLU:O	3:C:104:ASP:HB3	2.11	0.50
3:C:362:THR:O	3:C:368:GLY:HA3	2.15	0.47
3:C:390:PRO:HA	3:C:402:ARG:HH12	1.78	0.47
3:C:138:VAL:O	3:C:159:GLY:HA3	2.14	0.46
3:C:302:ASN:N	3:C:303:PRO:HD3	2.31	0.46
3:C:274:PHE:O	3:C:275:HIS:C	2.54	0.46
3:C:302:ASN:N	3:C:303:PRO:CD	2.79	0.46
3:C:522:ASN:HB2	6:C:1107:HOH:O	2.16	0.45
1:A:0:ACE:CH3	1:A:1:MET:H1	2.27	0.45
3:C:507:LYS:NZ	6:C:1364:HOH:O	2.49	0.44
3:C:287:ILE:HD12	3:C:290:MET:HG3	1.99	0.44
3:C:303:PRO:HG3	3:C:368:GLY:HA2	2.00	0.44
1:A:73:VAL:N	1:A:74:PRO:CD	2.81	0.43
3:C:132:GLY:HA3	3:C:155:THR:OG1	2.18	0.43
3:C:468:ILE:HB	3:C:469:PRO:CD	2.48	0.43
3:C:452:LYS:HG3	6:C:1186:HOH:O	2.18	0.43
3:C:220:KCX:CX	3:C:222:HIS:HD2	2.32	0.42
3:C:137:HIS:CE1	3:C:274:PHE:CD2	3.07	0.42
3:C:201:HIS:CE1	3:C:226:GLY:O	2.73	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:53:LYS:NZ	6:B:228:HOH:O	2.51	0.42
3:C:129:VAL:HG22	3:C:437:LEU:HG	2.02	0.42
3:C:158:PHE:CZ	3:C:195:GLY:HA3	2.55	0.41
3:C:303:PRO:HG2	3:C:367:MET:O	2.21	0.41
1:A:72:GLY:O	1:A:76:MET:HG3	2.20	0.41
1:A:32:TYR:HB3	1:A:33:PRO:HD3	2.03	0.41
3:C:385:LYS:HB3	6:C:1333:HOH:O	2.20	0.40
3:C:540:ASN:HA	3:C:541:PRO:HD2	1.86	0.40
2:B:78:GLU:HB3	2:B:79:PRO:HD2	2.03	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:C:1380:HOH:O	6:C:1380:HOH:O[11_555]	1.58	0.62
6:C:1389:HOH:O	6:C:1389:HOH:O[2_665]	1.64	0.56
6:C:1409:HOH:O	6:C:1409:HOH:O[12_565]	1.75	0.45
6:C:1432:HOH:O	6:C:1432:HOH:O[11_555]	2.00	0.20
3:C:327:GLN:O	3:C:327:GLN:OE1[7_556]	2.02	0.18
6:C:1223:HOH:O	6:C:1228:HOH:O[2_665]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/101 (98%)	99 (100%)	0	0	100	100
2	B	120/122 (98%)	114 (95%)	5 (4%)	1 (1%)	19	13
3	C	567/570 (100%)	543 (96%)	21 (4%)	3 (0%)	29	23
All	All	786/793 (99%)	756 (96%)	26 (3%)	4 (0%)	29	23

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	99	ILE
3	C	396	ASN
3	C	275	HIS
3	C	367	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	84 (99%)	1 (1%)	71	76
2	B	101/101 (100%)	100 (99%)	1 (1%)	76	81
3	C	460/460 (100%)	446 (97%)	14 (3%)	41	41
All	All	646/646 (100%)	630 (98%)	16 (2%)	47	49

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LEU
2	B	33	SER
3	C	1	MET
3	C	7	GLN
3	C	77	LEU
3	C	158	PHE
3	C	249	HIS
3	C	253	LEU
3	C	264	ARG
3	C	317	ASP
3	C	321	VAL
3	C	324	HIS
3	C	326	LYS
3	C	396	ASN
3	C	398	SER
3	C	536	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	267	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	KCX	C	220	3,5	9,11,12	3.14	2 (22%)	5,12,14	2.25	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KCX	C	220	3,5	-	0/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	220	KCX	OQ1-CX	6.65	1.34	1.21
3	C	220	KCX	CX-NZ	6.41	1.46	1.35

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	220	KCX	CE-NZ-CX	-3.18	116.78	121.89
3	C	220	KCX	CD-CE-NZ	-2.80	104.22	112.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	220	KCX	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	C	900	-	4,4,4	1.71	1 (25%)	6,6,6	0.55	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	900	SO4	O4-S	-3.15	1.22	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	0:ACE	C	1:MET	N	1.91
1	A	99:ILE	C	100:SER	N	1.02

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/101 (99%)	-0.78	1 (1%) 82 81	12, 17, 26, 30	2 (2%)
2	B	120/122 (98%)	-0.61	0 100 100	14, 20, 27, 32	5 (4%)
3	C	569/570 (99%)	-0.71	12 (2%) 63 62	11, 16, 34, 57	17 (2%)
All	All	789/793 (99%)	-0.71	13 (1%) 72 70	11, 17, 30, 57	24 (3%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	328	ASN	4.4
1	A	100	SER	4.2
3	C	397	GLY	3.9
3	C	322	CYS	3.9
3	C	329	ILE	3.5
3	C	392	ALA	3.5
3	C	550	GLY	2.5
3	C	391	LEU	2.4
3	C	324	HIS	2.3
3	C	327	GLN	2.2
3	C	321	VAL	2.1
3	C	390	PRO	2.1
3	C	396	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	KCX	C	220	12/13	0.97	0.07	10,14,17,18	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	C	900	5/5	0.97	0.16	37,38,39,41	0
5	NI	C	901	1/1	0.99	0.02	19,19,19,19	0
5	NI	C	902	1/1	1.00	0.02	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.