



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 30, 2024 – 10:44 pm BST

PDB ID : 2VCC  
Title : Family 89 Glycoside Hydrolase from Clostridium perfringens  
Authors : Ficko-Blean, E.; Stubbs, K.A.; Berg, O.; Vocadlo, D.J.; Boraston, A.B.  
Deposited on : 2007-09-19  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	891	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-N-ACETYLGLUCOSAMINIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	887	7228	4582	1194	1423	29	59	7	1

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	554	Total O 578 578	0	24

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ALPHA-N-ACETYLGLUCOSAMINIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.71Å 90.71Å 252.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.00 45.36 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.5 (20.00-2.00) 66.1 (45.36-2.00)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.38 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.210 , 0.255 0.198 , 0.242	Depositor DCC
$R_{free}$ test set	2675 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.4	Xtrriage
Anisotropy	0.187	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.050 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7830	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	6/7401 (0.1%)	0.56	0/10015

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	284	GLU	CD-OE1	-15.69	1.08	1.25
1	A	750	GLU	CG-CD	8.71	1.65	1.51
1	A	448	GLU	CB-CG	-7.92	1.37	1.52
1	A	213	ASP	CG-OD2	-5.72	1.12	1.25
1	A	835	GLU	CG-CD	5.40	1.60	1.51
1	A	911	LEU	C-N	-5.27	1.23	1.33

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	278	THR	CB

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7228	0	6825	71	0
2	A	18	0	24	0	0
3	A	5	0	0	0	0
4	A	1	0	0	0	0
5	A	578	0	0	8	0
All	All	7830	0	6849	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:371:ASN:HD21	1:A:482:HIS:HD2	1.13	0.94
1:A:605:THR:HG21	5:A:2384:HOH:O	1.65	0.94
1:A:790:LYS:CB	5:A:2503:HOH:O	2.22	0.87
1:A:790:LYS:CA	5:A:2503:HOH:O	2.28	0.80
1:A:790:LYS:N	5:A:2503:HOH:O	2.16	0.77
1:A:200:LYS:HE2	1:A:212:LYS:O	1.84	0.77
1:A:646[B]:GLU:OE2	5:A:2426:HOH:O	2.02	0.77
1:A:642[B]:LYS:HD2	1:A:642[B]:LYS:H	1.49	0.76
1:A:371:ASN:HD21	1:A:482:HIS:CD2	2.02	0.76
1:A:336:GLN:HE22	1:A:465:GLN:HE21	1.33	0.75
1:A:85:LYS:H	1:A:146:ASN:HD21	1.34	0.74
1:A:674:ARG:HD2	1:A:868:GLU:OE1	1.88	0.72
1:A:659:ARG:HD2	1:A:663:TYR:CE2	2.25	0.71
1:A:408:GLN:H	1:A:465:GLN:HE22	1.39	0.68
1:A:295:ARG:HD2	1:A:326:GLY:O	1.94	0.68
1:A:659:ARG:HD3	1:A:661:ASP:OD2	1.95	0.65
1:A:85:LYS:H	1:A:146:ASN:ND2	1.95	0.64
1:A:790:LYS:HB3	5:A:2503:HOH:O	1.90	0.63
1:A:790:LYS:O	1:A:791:ASP:HB2	2.00	0.61
1:A:642[B]:LYS:H	1:A:642[B]:LYS:CD	2.08	0.60
1:A:220:ARG:HD3	1:A:221:ASP:O	2.03	0.57
1:A:266:ASN:ND2	5:A:3360[B]:HOH:O	1.92	0.57

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338:GLU:OE2	1:A:341:ARG:NH1	2.38	0.56
1:A:211:PHE:HA	1:A:214:LYS:HD3	1.89	0.54
1:A:459:ASP:OD1	1:A:506:HIS:HE1	1.92	0.53
1:A:605:THR:HG23	1:A:606:ASN:OD1	2.09	0.52
1:A:572:MET:H	1:A:727:GLN:NE2	2.07	0.52
1:A:231:VAL:HG22	1:A:286:VAL:HG22	1.92	0.51
1:A:316:GLU:OE2	1:A:397:ARG:NH2	2.42	0.51
1:A:77:LEU:O	1:A:126:ASN:HA	2.12	0.50
1:A:565:ASN:HD21	1:A:568:GLY:H	1.60	0.50
1:A:263:VAL:HG13	1:A:272:ASN:O	2.11	0.50
1:A:383:TRP:CE2	1:A:387:ARG:HG3	2.47	0.49
1:A:659:ARG:CD	1:A:661:ASP:OD2	2.61	0.49
1:A:773:ARG:HB3	1:A:775:GLU:OE2	2.12	0.48
1:A:426:GLN:HB3	1:A:443:LYS:HD2	1.95	0.47
1:A:642[A]:LYS:CD	1:A:642[A]:LYS:H	2.26	0.47
1:A:811:TRP:CE3	1:A:822:LEU:HD13	2.49	0.47
1:A:605:THR:CG2	1:A:606:ASN:OD1	2.62	0.47
1:A:277:GLU:H	1:A:277:GLU:CD	2.17	0.46
1:A:564:HIS:HE1	1:A:605:THR:HG22	1.81	0.46
1:A:583:ILE:HB	1:A:584:PRO:HD3	1.98	0.46
1:A:361:PRO:HA	1:A:364:PHE:CE2	2.51	0.46
1:A:370:GLN:HG2	1:A:413:MET:SD	2.56	0.46
1:A:298:LEU:HD21	1:A:318:PHE:CZ	2.51	0.45
1:A:239:VAL:HG21	1:A:256:TYR:CZ	2.51	0.45
1:A:642[A]:LYS:H	1:A:642[A]:LYS:HD2	1.80	0.45
1:A:65:LYS:HB3	1:A:145:VAL:HG21	1.98	0.45
1:A:523:ASN:ND2	1:A:550:ARG:HE	2.14	0.45
1:A:833:LEU:O	1:A:837:TYR:HB3	2.17	0.44
1:A:494:LYS:NZ	5:A:2349:HOH:O	2.20	0.44
1:A:40:ASN:HA	1:A:59:GLU:O	2.17	0.44
1:A:262:ASN:OD1	1:A:634:GLU:HG2	2.18	0.43
1:A:523:ASN:HD21	1:A:550:ARG:HE	1.65	0.43
1:A:572:MET:H	1:A:727:GLN:HE21	1.65	0.43
1:A:665:GLY:O	1:A:687:HIS:HE1	2.02	0.43
1:A:319:LEU:HD11	1:A:394:MET:HG3	2.01	0.43
1:A:292:TYR:HB3	1:A:595:GLY:HA2	2.01	0.43
1:A:568:GLY:HA2	1:A:602:ALA:HB3	2.00	0.43
1:A:792:SER:O	1:A:797:LYS:NZ	2.51	0.43
1:A:92:TYR:HB3	1:A:136:VAL:HA	2.02	0.42
1:A:581:THR:HA	1:A:622:GLN:HG2	2.02	0.42
1:A:507:ASP:HB3	1:A:510:ALA:HB2	2.03	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:547:ASP:OD1	1:A:550:ARG:HD3	2.21	0.41
1:A:598:ILE:HB	1:A:600:PRO:HD3	2.01	0.41
1:A:138:GLN:HG2	1:A:139:ASN:N	2.35	0.41
1:A:79:LYS:HB2	1:A:153:PHE:HB2	2.02	0.40
1:A:659:ARG:HD2	1:A:663:TYR:CD2	2.56	0.40
1:A:596:ILE:HD11	1:A:617:ALA:HB2	2.03	0.40
1:A:775:GLU:H	1:A:775:GLU:CD	2.24	0.40
1:A:539:ASP:O	1:A:561:ASN:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	891/891 (100%)	861 (97%)	29 (3%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	298	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	769/767 (100%)	734 (95%)	35 (5%)	27 23

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	LEU
1	A	125	ASP
1	A	165	LYS
1	A	166	LYS
1	A	182	GLU
1	A	209	ARG
1	A	220	ARG
1	A	223	LEU
1	A	231	VAL
1	A	247	VAL
1	A	262	ASN
1	A	278	THR
1	A	295	ARG
1	A	341	ARG
1	A	379	LEU
1	A	397	ARG
1	A	442	LEU
1	A	471	ASP
1	A	529	LEU
1	A	541	PHE
1	A	556	LEU
1	A	565	ASN
1	A	605	THR
1	A	642[A]	LYS
1	A	642[B]	LYS
1	A	645	LEU
1	A	659	ARG
1	A	706	LYS
1	A	709	ASP
1	A	713	ASP
1	A	721	PHE
1	A	811	TRP
1	A	823	LYS
1	A	853	LEU
1	A	894	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	ASN
1	A	118	ASN
1	A	146	ASN
1	A	151	ASN
1	A	192	ASN
1	A	201	ASN
1	A	254	ASN
1	A	325	ASN
1	A	370	GLN
1	A	426	GLN
1	A	455	GLN
1	A	465	GLN
1	A	482	HIS
1	A	506	HIS
1	A	518	GLN
1	A	523	ASN
1	A	534	GLN
1	A	565	ASN
1	A	687	HIS
1	A	727	GLN
1	A	745	ASN
1	A	847	ASN
1	A	888	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	A	1914	-	5,5,5	0.42	0	5,5,5	0.12	0
3	SO4	A	1916	-	4,4,4	0.13	0	6,6,6	0.11	0
2	GOL	A	1915	-	5,5,5	0.35	0	5,5,5	0.45	0
2	GOL	A	1913	-	5,5,5	0.37	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1914	-	-	4/4/4/4	-
2	GOL	A	1915	-	-	3/4/4/4	-
2	GOL	A	1913	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1914	GOL	O1-C1-C2-C3
2	A	1915	GOL	C1-C2-C3-O3
2	A	1915	GOL	O2-C2-C3-O3
2	A	1913	GOL	O1-C1-C2-C3
2	A	1913	GOL	C1-C2-C3-O3
2	A	1914	GOL	C1-C2-C3-O3
2	A	1914	GOL	O1-C1-C2-O2
2	A	1914	GOL	O2-C2-C3-O3
2	A	1913	GOL	O1-C1-C2-O2
2	A	1915	GOL	O1-C1-C2-C3
2	A	1913	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	886/891 (99%)	-0.21	14 (1%) 72 70	20, 31, 44, 57	20 (2%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	857	ALA	5.9
1	A	26	GLY	3.4
1	A	909	LYS	3.3
1	A	855	GLY	3.0
1	A	858	LYS	2.7
1	A	911	LEU	2.6
1	A	859	ALA	2.2
1	A	748	ASN	2.2
1	A	807	LEU	2.2
1	A	754	PHE	2.2
1	A	746	ASN	2.1
1	A	42	GLU	2.1
1	A	749	GLY	2.1
1	A	305	TYR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	A	1913	6/6	0.74	0.27	59,61,61,61	0
2	GOL	A	1914	6/6	0.83	0.16	48,50,50,51	0
3	SO4	A	1916	5/5	0.90	0.24	94,94,94,94	0
2	GOL	A	1915	6/6	0.91	0.27	41,42,43,44	0
4	CA	A	1917	1/1	0.99	0.05	29,29,29,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.